

Minimally invasive surgery plus rt-PA for ICH evacuation

Submission date 29/04/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 29/04/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 10/09/2019	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Study website
<http://www.braininjuryoutcomes.com>

Contact information

Type(s)
Scientific

Contact name
Dr Barbara Gregson

Contact details
Neurosurgical Trials Unit
Newcastle University
3-4 Claremont Terrace
Newcastle upon Tyne
United Kingdom
NE2 4AE

Additional identifiers

EudraCT/CTIS number
2007-006006-22

IRAS number

ClinicalTrials.gov number
NCT00224770

Secondary identifying numbers

6469

Study information

Scientific Title

Minimally invasive surgery plus rt-PA for ICH evacuation

Acronym

MISTIE

Study objectives

The purpose of this trial is to determine the safety of using a combination of minimally invasive surgery and clot lysis with rt-PA to remove ICH.

Ethics approval required

Old ethics approval format

Ethics approval(s)

14/01/2009, ref: 08/H0906/158

Study design

Randomised interventional treatment trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Topic: Generic Health Relevance and Cross Cutting Themes, Stroke Research Network; Subtopic: Generic Health Relevance (all Subtopics); Disease: Surgery

Interventions

Stage 1 Dose Finding: 60 patients with ICH (45 surgical and 15 medical, 3:1 randomisation). Tiers 1 and 2: MISTIE + rt-PA versus medical.

Stage 2 Safety: 50 patients with ICH (25 surgical and 25 medical, 1:1 randomisation)

The neurosurgeon will review the stability CT scan to determine the burr hole location and trajectory to be used during the operative procedure to place the catheter. The Surgical Center personnel will review whether the proposed burr hole location and trajectory is appropriate or

that a different location/trajectory is recommended. The catheter placement will be performed in either the operating room or the ICU. Careful hematoma aspiration is performed free hand using a 10 cc syringe.

Follow up length: 6 months

Study entry: single randomisation only

Intervention Type

Other

Phase

Phase II/III

Primary outcome measure

Mortality, 30 days and procedure related.

Secondary outcome measures

1. Cerebritis, meningitis
2. Clot size reduction, post-operative and by day 4 - 5
3. Glasgow Outcome Scale (GOS), extended GOS (eGOS), Rankin, Stroke Impact Scale (SIS), measured at 30, 90, 180, 270, 365 days
4. Symptomatic rebleeding

Overall study start date

01/04/2010

Completion date

01/12/2010

Eligibility

Key inclusion criteria

1. Aged 18 - 80 years, either sex
2. Glasgow Coma Scale (GCS) less than 14 or a National Institutes of Health Stroke Scale (NIHSS) (including the use of distal hand measures) greater than 6
3. Spontaneous supratentorial ICH greater than 20 cc
4. Symptoms less than 12 hours prior to diagnostic CT scan (an unknown time of symptom onset is exclusionary)
5. Intention to initiate surgery within 48 hours after diagnostic CT
6. First dose can be given within 54 hours of diagnostic CT
7. Six-hour clot size equal to the most previous clot size + 5 cc (as determined by additional CT scan at least 6 hours after the initial stability scan ($A*B*C$)/2 method)
8. Systolic blood pressure (SBP) less than 200 mmHg sustained for 6 hours recorded closest to the time of randomisation
9. Historical Rankin score of 0 or 1
10. Negative pregnancy test

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

Planned Sample Size: 110

Total final enrolment

141

Key exclusion criteria

1. Infratentorial hemorrhage including brainstem (any involvement of the midbrain or lower brainstem as demonstrated by radiograph or complete third nerve palsy)
2. Patients with platelet count less than 100,000, international normalised ratio (INR) greater than 1.7, or an elevated prothrombin time (PT) or activated partial thromboplastin time (APTT) (reversal of coumadin is permitted but the patient must not require coumadin during the acute hospitalisation). Irreversible coagulopathy either due to medical condition or prior to randomisation (patient must have a sustained INR less than 1.7 using short- and long-acting procoagulants [Novoseven, FFP, and/or vitamin K]).
3. Clotting disorders
4. Any concurrent serious illness that would interfere with the safety assessments including hepatic, renal, gastroenterologic, respiratory, cardiovascular, endocrinologic, immunologic, and haematologic disease
5. Patients with a mechanical valve
6. Patients with unstable mass or evolving intracranial compartment syndrome
7. Ruptured aneurysm, arteriovenous malformation (AVM), vascular anomaly
8. Greater than 80 years (higher incidence of amyloid)
9. Under 18 years of age (high incidence of occult vascular malformation)
10. Pregnant (positive pregnancy test) or lactating females (likelihood of altered coagulation function associated with the high oestrogen/progesterone state)
11. Irreversibly impaired brainstem function (bilateral fixed, dilated pupils and extensor motor posturing), GCS less than or equal to 4
12. Historical Rankin score greater than or equal to 2
13. Intraventricular haemorrhage requiring external ventricular drainage
14. Internal bleeding, involving retroperitoneal sites, or the gastrointestinal, genitourinary, or respiratory tracts
15. Superficial or surface bleeding, observed mainly at vascular puncture and access sites (e.g., venous cutdowns, arterial punctures) or site of recent surgical intervention
16. Known risk for embolisation, including history of left heart thrombus, mitral stenosis with atrial fibrillation, acute pericarditis, and subacute bacterial endocarditis
17. In the investigator's opinion, the patient is unstable and would benefit from a specific intervention rather than supportive care plus or minus endoscopic or MIS+rtPA
18. Prior enrolment in the study
19. Any other condition that the investigator believes would pose a significant hazard to the subject if the investigational therapy were initiated
20. Participation in another simultaneous trial of ICH treatment

Date of first enrolment

01/04/2010

Date of final enrolment

01/12/2010

Locations

Countries of recruitment

England

Germany

United Kingdom

Study participating centre

Neurosurgical Trials Unit

Newcastle upon Tyne

United Kingdom

NE2 4AE

Sponsor information

Organisation

Johns Hopkins University (USA)

Sponsor details

1550 Orleans Street,

3M50 South,

Baltimore, Maryland

United States of America

21231

Sponsor type

University/education

Website

<http://www.braininjuryoutcomes.com>

ROR

<https://ror.org/00za53h95>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research (NIHR) (UK)

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results	results	01/03/2013	10/09/2019	No	No
Results article			10/09/2019	Yes	No
HRA research summary			28/06/2023	No	No