A research study in Uganda to test the acceptability and feasibility of an intervention called DIALOG+, designed to improve care for people in primary care with poor physical and mental health

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
27/11/2020		[X] Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
02/12/2020		[X] Results		
Last Edited	Condition category	[] Individual participant data		
16/11/2023	Other			

Plain English summary of protocol

Background and study aims

DIALOG+ is an intervention delivered on a tablet or smartphone using an app. It is designed to help mental health professionals to improve the structure of their routine meetings with patients. It also helps to improve communication with patients during these meetings. Patients are first asked about how satisfied they are with eight areas of their lives (e.g. physical health, family relationships, leisure activities) and three areas of the treatment they are receiving (e.g. practical help, meetings), which is called the DIALOG scale. The patient then chooses up to three areas to discuss in more depth with their health professional. The clinician then discusses each area chosen by patients, using four steps that focus on solutions to the identified problems. This study aims to find out whether DIALOG+ can be adapted to be used to improve care for people in primary care with physical conditions and poor quality of life in Uganda. More specifically, the researchers want to find out how patients and health professionals experience DIALOG+ when it is used during their routine meetings. They also want to find out if DIALOG+ improves outcomes like quality of life and other symptoms in order to plan for larger definitive trials if indicated.

Who can participate?

Adults in primary care who have at least one common noncommunicable chronic condition (such as diabetes, hypertension, chronic obstructive pulmonary disease, cardiovascular disease, etc) and who have a low quality of life

What does the study involve?

The initial stage of this study involves a suggested adaptation of DIALOG+ from other trials that have been conducted. Potential minor adaptations may include suggested modifications to the number of domains addressed, item wording and training examples. DIALOG+ will be adapted to primary care settings to enable wider use with patients with chronic physical and mental health conditions with assessments before and 3 months after the intervention. The adaptation

process will be through stakeholder consultation based on actual experience of using the intervention. The results will provide insight into refining the training process and implementing the intervention across healthcare settings.

What are the possible benefits and risks of participating?

DIALOG+ is a resource-oriented and evidence-based intervention which makes use of existing personal and social resources to improve the quality of life of patients with mental illness. The intervention is low cost using routine meetings between patients and clinicians without the need for additional referrals or services. The researchers will adapt DIALOG+ for primary care to improve reach and enable a larger patient population to benefit. Additionally, the study focuses on a new and vulnerable population – those with comorbid (simultaneously present) noncommunicable chronic physical and mental health problems. Research has shown that comorbid physical and mental health problems are common, resource-intensive and result in a poor quality of life for billions of people worldwide.

A potential benefit for all participants involved in the study is that their suggestions and experiences might be incorporated into further adaptations, which will tailor the intervention to the needs of patients and clinicians in the context of the primary care system in Uganda. Additionally, for patients who will be involved in testing the DIALOG+, this might lead to improved quality of life, social functioning, and symptoms. The study will also benefit clinicians involved in terms of the training they will receive. Clinicians involved in the DIALOG+ study will be provided with training and supervision to enable them to implement the intervention. The researchers do not foresee any significant ethical, legal or management issues arising from this study. Within the assessments and interviews that will take place across both studies, questions will be raised with participants that might trigger feelings of distress or anxiety. Participants may experience anxiety in trying new interventions. Throughout the interventiontesting period, participants will continue to receive their routine care, including any medication, in addition to the intervention. The intervention can be stopped at any point. The use of DIALOG+ in mental health care settings has an evidence base for effectiveness and the researchers believe this is easily transferable across to physical health care too without any additional risks. To minimise the impact of potential risks, risk management strategies have been outlined in the study protocol.

Where is the study run from?
Outpatient clinic at Mityana and Masaka Hospitals (Uganda)

When is the study starting and how long is it expected to run for? November 2019 to June 2021

Who is funding the study? National Institute for Health Research (NIHR) (UK)

Who is the main contact?
Dr Francois van Loggerenberg, f.vanloggerenberg@qmul.ac.uk

Contact information

Type(s)Scientific

Contact name

Dr Francois van Loggerenberg

ORCID ID

https://orcid.org/0000-0001-5317-7983

Contact details

Unit for Social and Community Psychiatry Queen Mary University of London Newham Centre for Mental Health London United Kingdom E13 8SP +44 (0)207 540 4380 Ext: 2339 f.vanloggerenberg@qmul.ac.uk

Type(s)

Public

Contact name

Dr Francois van Loggerenberg

ORCID ID

https://orcid.org/0000-0001-5317-7983

Contact details

Unit for Social and Community Psychiatry Queen Mary University of London Newham Centre for Mental Health London United Kingdom E13 8SP +44 (0)207 540 4380 Ext: 2339 f.vanloggerenberg@qmul.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

16/137/97

Study information

Scientific Title

Testing the acceptability and feasibility of DIALOG+ in patients with common chronic conditions and poor mental health in primary care in Uganda: a non-randomized study with before and after design

Study objectives

The study aims to test the feasibility and acceptability of and adapted DIALOG+ intervention for patients with comorbid physical and mental health conditions in primary care.

The main research questions are:

- 1. Is the use of DIALOG+ feasible and acceptable in primary care for patients with non-communicable physical health conditions
- 2. How do we adapt the DIALOG+ intervention through a process of stakeholder consultation and use the adapted version to refine training materials?
- 3. How does the adapted DIALOG+ intervention affect patient outcomes?

Ethics approval required

Old ethics approval format

Ethics approval(s)

- 1. Approved 05/11/2020, Makerere University, College of Health Sciences School of Medicine Research and Ethics Committee (PO Box 7072 Kampala; +256 (0)414 533541; research@chs.mak. ac.uq), ref: 2020-195
- 2. Approved 18/12/2020, Uganda National Council of Science and Technology (PO Box 6884, Kampala; +256(0)41 4705500), ref: HS1116ES
- 3. Approved 27/01/2021, Queen Mary Ethics of Research Committee (Hazel Covill, Room W117, Finance Department, Queens' Building, Queen Mary University of London, Mile End Road, London, E1 4NS, UK; +44 (0)20 7882 7915; h.covill@qmul.ac.uk), ref: QMERC20.101

Study design

Non-randomized design using mixed methods and before and after assessments

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Chronic conditions such as diabetes, hypertension, chronic obstructive pulmonary disease (COPD), cardiovascular disease (CVD)

Interventions

DIALOG+ is a technology-mediated intervention, which involves a structured patient assessment covering satisfaction with eight life domains and three treatment domains (DIALOG scale) and a four-step solution-focused therapy approach to address patient concerns (DIALOG+). DIALOG+ aims to make routine meetings between clinicians and patients therapeutically effective.

Once enrolled, patients will receive DIALOG+ at their routine primary care appointments, around once per month. This will be delivered by the healthcare worker using an app on a tablet computer. The intervention period will be 3 months, during which patients will attend up to three DIALOG+ sessions.

Data collection with all participants will take place at baseline and following the 3-month intervention period. At baseline, the researchers will collect socio-demographic information from all participants, which, for patient participants, will include clinical characteristics.

A subset of participants will be invited to attend a qualitative interview after the end of the intervention (3 months) in order to capture the individual experience of the intervention; including barriers and facilitators of attending intervention sessions, suggested adaptations and the practical delivery of the intervention.

In addition to the research data collection, the research and clinical support staff keep a record of their intervention sessions. These reports will include the date, duration of interventions and the topic of discussion. This information is also captured within the DIALOG+ app and can be retrieved by the research staff. The aim is to record at least one session per patient, which will be scored for adherence.

Intervention Type

Behavioural

Primary outcome(s)

- 1. Quality of life measured using the Manchester Short Assessment (MANSA) at baseline and 3 months
- 2. Depression measured using the Patient Health Questionnaire (PHQ-9) at baseline and 3 months
- 3. Anxiety measured using Generalised Anxiety Disorder assessment (GAD-7) at baseline and 3 months
- 4. Objective social functioning measured using objective social outcomes index (SIX) at baseline and 3 months
- 5. Physical health symptoms measured using The Short Form (36) Health Survey (SF-36) at baseline and 3 months

Key secondary outcome(s))

There are no secondary outcome measures

Completion date

25/06/2021

Eligibility

Key inclusion criteria

- 1. Diagnosis of ≥1 common chronic non-communicable condition such as diabetes, hypertension, COPD. CVD. etc
- 2. Quality of life score measured on the MANSA ≤5
- 3. Capacity to provide informed consent assessed using the UBACC
- 4. Able to speak and understand the local language
- 5. Attending the outpatient clinic for at least 6 months
- 6. Live within a 20 km radius of the clinic
- 7. Aged 18 years and above

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

182

Key exclusion criteria

- 1. Unable or unwilling to provide informed consent
- 2. Does not meet inclusion criteria

Date of first enrolment

15/02/2021

Date of final enrolment

07/04/2021

Locations

Countries of recruitment

Uganda

Study participating centre Mityana Hospital

PO Box 52

Mityana

Uganda

Uganda

Study participating centre Masaka Regional Referral Hospital

PO Box 18

Masaka

Uganda

Uganda

Sponsor information

Organisation

Queen Mary University of London

ROR

https://ror.org/026zzn846

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets will be held at QMUL in anonymised form. Data sharing with external interests will be considered only after the publication of the findings that reflect the given data. The datasets will be available upon request from Stefan Priebe (s.priebe@qmul.ac.uk). The data collected will be both quantitative and qualitative. The duration of availability of data has not yet been decided. During the course of the study, data will be shared internally between the research group using an online data collection platform called REDCap. The method for sharing the data externally (if required) will be decided in due course. Informed consent will be obtained from all participants involved in the study. All participants are assigned a patient ID at the point of enrolment and all subsequent data collected will be linked to this ID, without any link to identifiable data following Good Clinical Practice.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		16/11/2023	16/11/2023	Yes	No
<u>Protocol article</u>		30/09/2021	04/10/2021	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes