# Treadmill training in sub-acute stroke

Submission date	<b>Recruitment status</b> No longer recruiting	Prospectively registered		
22/03/2017		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
19/04/2017	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
25/09/2017	Nervous System Diseases			

#### Plain English summary of protocol

Background and study aims

A stroke occurs when the blood supply to the brain is cut off. This can cause long term problems with mobility due to the injury to the brain. A major problem for people who have had a stroke is learning to walk again. Physiotherapy is a type of rehabilitation that helps restore movement and function to patients. Physiotherapists help stroke patients to walk using different types of exercises. Sometimes different equipment, such as using a treadmill, can be used to help patients to walk again. This study looks at the feasibility of using a treadmill and the outcomes of relearning how to walk for people who had had a stroke within the last three months. The main aims of this study is to see if a study like this is feasible, to see how to provide treadmill training for stroke patients, see if using a treadmill can help improve mobility when compared to standard physiotherapy patients and to create a larger study in the future.

#### Who can participate?

Adults over the age of 18 who have had a stroke.

#### What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group receive the standard physiotherapy treatment and gait (walking) training. Those in the second group receive the standard physiotherapy treatment three times a week over eight weeks. They also receive at least two gait training sessions that use a treadmill. After the eight weeks, participants continue with their standard physiotherapy treatments without using a treadmill. Participants are assessed before the treatment, right after the treatment and in a six month follow up to assess their mobility.

What are the possible benefits and risks of participating?

Participants may benefit from improving their mobility, independence and quality of life. There are minimal risks with participating, such as falling during physiotherapy; however participants are given support by physiotherapists to prevent this.

Where is the study run from?

This study is run by Queen Margaret University Edinburgh and takes place in Astley Ainslie Hospital (UK), St Johns Hospital (UK), the Royal Victoria Hospital (UK) and the Liberton Hospital (UK)

When is the study starting and how long is it expected to run for? October 2005 to December 2008

Who is funding the study? Chest Heart and Stroke Scotland (UK)

Who is the main contact? Dr Gillian Baer

# **Contact information**

# Type(s)

Scientific

#### Contact name

Dr Gillian Baer

#### Contact details

Physiotherapy School of Health Sciences Queen Margaret University Queen Margaret University Drive Edinburgh Edinburgh United Kingdom EH21 6UU

# Additional identifiers

#### Protocol serial number

R06/A99

# Study information

#### Scientific Title

Treadmill Training for people with Sub-Acute Stroke: A phase II randomised controlled trial

## Acronym

**STATT** 

#### Study objectives

The aim of this pilot study is to evaluate the effects and feasibility of using treadmill training in people with stroke, within the first 3 months after stroke.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Scotland A Multi-Centre Research Ethics Committee, 16/10/2006, 06/MRE00/82

#### Study design

Exploratory phase II single-blind feasibility multi-site randomised controlled trial

#### Primary study design

Interventional

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Sub-acute Stroke

#### **Interventions**

Participants are randomly allocated to either the intervention group or the control group. Randomisation is done after participants undergo baseline measurements using a computer generated minimisation method. Minimisation takes into account the side of the body that the stroke occurred on and whether the participants are ambulant without physical assistance (using the Functional Ambulation category 4-6) or the participants are non-ambulant or ambulant only with physical assistance (Functional Ambulation category 1-3).

Intervention Group (TT): This intervention occurs over eight weeks (or fewer if the participant is discharged from the unit before the eight weeks). Participants in this group receive at least three treatment sessions per week comprising of normal physiotherapy and gait training. At least two gait training sessions include the treadmill. The actual time on the treadmill is determined by the treating physiotherapist. All the time on the treadmill is recorded. After eight weeks, participants in this group reverted back to normal physiotherapy with no further access to the treadmill.

Control group (CON): This intervention occurs over eight weeks (or fewer if the participant is discharged from the unit before the eight weeks). Participant in this group receive at least three treatment sessions per week comprising of normal physiotherapy and gait training. Gait training has no access to a treadmill. The time for the normal physiotherapy is determined clinically.

Participants are assessed for their mobility at baseline, at the end of the interventions (eight weeks) and at a six month follow up. As this is a feasibility phase II randomised controlled trial, data from a battery of secondary outcome measures are collected.

#### **Intervention Type**

Other

#### Primary outcome(s)

Mobility is measured using the Rivermead Mobility Index at baseline, eight weeks and six months.

# Key secondary outcome(s))

- 1. Mobility and balance is measured using the Timed Up and Go (TUG) at baseline, eight weeks and six months
- 2. Gait speed is measured using a 10 metre walk (10mwt), at baseline, eight weeks and six months
- 3. Walking endurance is measured using a six minute walk test (6minwt) at baseline, eight weeks and six months
- 4. Confidence in walking is measured using a vertical 10cm Visual Analogue Scale (VAS), patient

interviews and following a standardised operating procedure at baseline, eight weeks and six months

- 5. Motor impairments and everyday motor function is measured using the Motor Assessment Scale (MAS) at baseline, eight weeks and six months
- 6. Dependence in Activities of Daily Living is measured using Barthel Index at baseline, eight weeks and six months
- 7. Self reported recovery and quality of life is measured using the Stroke Impact Scale v3.0 at baseline, eight weeks and six months

#### Completion date

19/12/2008

# Eligibility

### Key inclusion criteria

- 1. Aged over 18
- 2. Had a stroke as defined by WHO 1988
- 3. One minute standing balance (with or without support, this was required to allow donning a safety harness if supported treadmill training was required)
- 4. Medically stable
- 5. Within three months of stroke onset
- 6. Able to understand and follow verbal instructions; and
- 7. Able to give informed consent

#### Participant type(s)

Patient

## Healthy volunteers allowed

No

#### Age group

Adult

## Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

- 1. Co-existing non-stroke related neurological impairments
- 2. Co-morbidities precluding gait training (e.g. amputee)
- 3. Non-ambulant prior to stroke
- 4. Body weight greater than 138kg (Weight limit of equipment)
- 5. Unsafe to use treadmill
- 6. Unable to follow simple commands

#### Date of first enrolment

04/05/2007

# Date of final enrolment

20/06/2008

# Locations

#### Countries of recruitment

**United Kingdom** 

Scotland

# Study participating centre Astley Ainslie Hospital

Physiotherapy 33 Grange Loan Edinburgh United Kingdom EH9 2HL

# Study participating centre St Johns Hospital

Physiotherapy Howden Road West Howden Livingston United Kingdom EH54 6PP

# Study participating centre The Royal Victoria Hospital

Physiotherapy 13 Craigleith Road Edinburgh United Kingdom EH4 2DN

# Study participating centre Liberton Hosptial

Physiotherapy 113 Lasswade Road Edinburgh United Kingdom EH16 6UB

# Sponsor information

## Organisation

Queen Margaret University

#### **ROR**

https://ror.org/002g3cb31

# Funder(s)

#### Funder type

Charity

#### **Funder Name**

Chest Heart and Stroke Scotland

## Alternative Name(s)

Chest Heart & Stroke Scotland, CHSScotland, CHSS

#### **Funding Body Type**

Private sector organisation

#### **Funding Body Subtype**

Associations and societies (private and public)

#### Location

United Kingdom

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Gillian Baer (Principal Investigator) gbaer@qmu.ac.uk

# IPD sharing plan summary

Available on request

# **Study outputs**

Output type	<b>Details</b> results	Date created Date added Peer reviewed? Patient-facing?		
Results article		01/02/2018	Yes	No
Participant information sheet	version V3	01/11/2007 21/04/20	17 No	Yes