

The effect of the Oxford Head Elevating Laryngoscopy Pillow (OHELP), on spinal anaesthesia in elective caesarean sections

Submission date 22/11/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 31/01/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 31/01/2014	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Head Elevating Laryngoscopy Pillow (HELP) was introduced recently to the National Maternity Hospital and since then it has been a routine measure of safety in obese patient airway management; however, its impact on local anaesthetic effect post spinal anaesthesia is still unknown. The pillow is used routinely for obese mothers, and there are no obvious safety concerns with it.

The pillow is usually placed under the head and shoulders, and it changes the angle of the airway, making it easier to access the patients airway without moving the patient in the rare case that a general anaesthetic is needed.

Since the pillow changes the angle of the upper body, it might affect the spread of the local anaesthetic that is given for caesarean sections, and this initial study has been designed to investigate whether this might be the case. Since there is the possibility that use of the pillow will slow the spread of the anaesthetic, a catheter will be left in so that top-up doses can be administered easily.

Who can participate?

Expectant mothers, gestational age > 32 weeks and undergoing elective caesarean section.

What does the study involve?

Participants will be randomly allocated to one of two groups.

The intervention group will lie on an Oxford Head Elevating Laryngoscopy Pillow and a standard head pillow.

The control group will use only a standard head pillow. We will then compare the two groups of mothers to see whether they achieved a satisfactory level of anaesthetic block (numbness to touch) after 10 minutes.

We will also be investigating the number of women from each group that required conversion to general anaesthetic, and we will compare whether top-up doses of anaesthetic will be required.

What are the possible benefits and risks of participating?

The outcome of the study will hopefully be to establish whether the benefits of the pillow

easier airway management in case of general anaesthetic might not outweigh any disadvantages such as poor anaesthesia leading to discomfort or pain during the procedure, or even increasing the rate of requiring general anaesthetic. Since there is no evidence one way or the other at the moment, and the pillow is in routine use, it is important that this issue be investigated.

Where is the study run from?

The National Maternity Hospital, Ireland.

When is the study starting and how long is it expected to run for?

The study started in June 2013 and ran until October 2013.

Who is funding the study?

National Maternity Hospital, Ireland.

Who is the main contact?

Dr Hayat Elfil

Contact information

Type(s)

Scientific

Contact name

Dr Hayat Elfil

Contact details

Dept. of Anaesthesiology
National Maternity Hospital
Holles Street
Dublin
Ireland
2

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

The effect of the Oxford Head Elevating Laryngoscopy Pillow (OHELP), on subarachnoid local anesthetic spread in elective caesarean sections

Acronym

OHELP

Study objectives

That the Oxford Head Elevating Laryngoscopy Pillow slows the cephalad spread of local anaesthetic and therefore delays the onset of satisfactory sensory blockade (T6 or better) to perform caesarean section.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Research Ethics Committee - National Maternity Hospital, 30/04/2013

Study design

Single-centre prospective randomised controlled trial with two non-blinded parallel arms

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Extent of sensory block in females undergoing elective caesarean section for a singleton pregnancy, and receiving combined spinal-epidural (CSE) anaesthetic at L3/4

Interventions

Intervention group and a control group from consecutively eligible parturients.

The intervention group will lie on an Oxford Head Elevating Laryngoscopy Pillow and a standard head pillow .

The control group will use only a standard head pillow.

A seated position will be adopted for the neuraxial blockade performance. Combined Spinal Epidural will be administered at L3/4 (by palpating iliac crests). The operating theatre table will be held in the zero position (completely horizontal with no head elevation). The epidural space will be located using the loss of resistance to air (NOT saline) technique. The subarachnoid space will be located using the needle through needle technique.

Intrathecal injection will consist of 2.2 ml of hyperbaric Bupivacaine 0.5% and Fentanyl 15 micrograms + Morphine 100 micrograms. The epidural catheter will be left at 4 cm in the epidural space, but nothing will be injected via epidural catheter unless there is a block failure.

Left uterine displacement held at 20 degrees facilitated by Cardiff wedge support.

Block failure is defined as sensory block not reaching T6 (defined as level of xiphisternum) within 10 mins, in which case the patient will be given epidural boli of 5 ml Lignocaine 2% + Adrenaline 1:200,000 until block is satisfactory.

Patients were closely monitored post their caesarean sections for their sensory and motor block until they were able to move their legs and were haemodynamically stable. This took around one hour on average.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Sensory block will be assessed at 10 min post subarachnoid anesthesia induction (time 0) using loss of touch sensation to ethyl chloride sprayed bilaterally in the midclavicular line. The ethyl chloride spray container will be 5 cm from the patient's skin, and a continuous jet will be sprayed as the container is moved in a cranial direction.

Primary outcome measure is a binary variable defined as lack of sensory block on the right or the left, below T6, versus bilateral block at least as high as T6.

Measured at 10 min from the baseline.

Secondary outcome measures

1. Conversion to general anaesthetic for any reason
2. Administration of a top-up dose of anaesthetic of any sort

Measured at the baseline

Overall study start date

01/06/2013

Completion date

15/10/2013

Eligibility

Key inclusion criteria

All parturients agreeing to participate in the trial where these criteria are met:

1. Elective caesarean section where there is no fetal or maternal compromise
2. Singleton pregnancy
3. Gestational age > 32 weeks

Participant type(s)

Patient

Age group

Adult

Sex

Female

Target number of participants

100 (50 per arm)

Key exclusion criteria

1. Refusal to consent
2. Parturients opting for GA or with medical conditions preventing neuraxial anaesthesia
3. Emergency caesarean sections where there is fetal or maternal compromise (Lucas Grade 1 and 2)
4. Multiple pregnancy
5. Gestational age < 32 weeks
6. Body mass index (BMI) > 40

Date of first enrolment

01/06/2013

Date of final enrolment

15/10/2013

Locations**Countries of recruitment**

Ireland

Study participating centre

Dept. of Anaesthesiology

Dublin

Ireland

2

Sponsor information**Organisation**

The National Maternity Hospital (Ireland)

Sponsor details

Holles Street
Dublin
Ireland
2
+353 (0)1 637 3100
bodriscoll@nmh.ie

Sponsor type

Hospital/treatment centre

Website

<http://www.nmh.ie>

ROR

<https://ror.org/03jcx214>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

National Maternity Hospital (Ireland)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration