

Application of Multiple Cerebral Monitoring Severe Traumatic Brain Injury: a multicentre study

Submission date 31/08/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 10/07/2007	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 27/10/2021	Condition category Nervous System Diseases	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Wen-Ta Chiu

Contact details

No. 250 Wu-Hsing Street

Taipei

Taiwan

111

+886 (0)2 2739 0217

wtchiu@tmu.edu.tw

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

Application of Multiple Cerebral Monitoring Severe Traumatic Brain Injury: a multicentre study

Acronym

AMCMSTBI

Study objectives

Significant reduction in mortality and morbidity can be achieved by using intensive management protocols. Compared to early studies of 50% mortality, the results of efforts in investigating the proper procedures among each stages, reveals a reduction of mortality rate to average of 36% among severe head injured patients. Kiening et al. reported that the elevation of the Cerebral Perfusion Pressure (CPP) from 32 to 67 mmHg improved cerebral tissue oxygenation (PtO₂) by 62%. Recent studies also showed that the mortality significantly decreased in the group of PtO₂ of monitoring. However, there are still debates about the optimal CPP, especially in the patient with impaired pressure autoregulation. Nevertheless, optimal Inter-Cranial Pressure (ICP) and CPP management may not always prevent cerebral ischaemia, which adversely influences patient outcome.

The aims of this study are:

1. To evaluate the efficacy of multiple cerebral monitoring
2. To find the optimal CPP in different groups with autoregulation in the treatment of Severe Traumatic Brain Injury (STBI)
3. To investigate the outcome difference (Glasgow Outcome Score [GOS], Glasgow Outcome Score Extended [GOSE], survival analysis and Health-related Quality Of Life [HQOL]) among patients with different monitoring groups

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval gained from the local ethics committee on the 1st June 2006 (reference number: F950402).

Study design

Interventional, prospective, multicentred, randomised controlled study.

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Severe Traumatic Brain Injury

Interventions

The hospitals included in this series were Wan Fang Hospital, Veterans General Hospital, Taipei, National Taiwan University Hospital, Mackay Memorial Hospital, Shin Kong Wu Ho-Su Memorial Hospital and Taipei Tri-Service General Hospital. Twenty to thirty patients are provided from each medical centre.

All the subjects are randomised (block in four) and divided in to the study group (Group A) or control group (Group B):

Group A: multiple cerebral monitoring systems, ICP, PtO₂ and the TransCranial Doppler (TCD74)

Group B: ICP monitoring only

The major differences between the Group A and Group B are the treatment goals. In the Group A, PtO₂ of at least 25 - 30 mmHg is maintained, by the adjustment of CPP. In Group B, Mean Arterial Pressure (MAP) was kept at 90 mmHg or more and CPP was maintained at a minimum of 60 - 70 mmHg.

Cerebral monitoring:

1. The TransCranial Doppler (TCD) insonation was performed using a 2 MHz real time and pulse-Doppler transducer. For continuous monitoring, the 2 MHz probe was applied at a temporal window with a fixed head holder (Spenser Marco 600) connected to the TCD monitor (Nicolet /EME Companion II, Germany) to detect the Middle Cerebral Artery (MCA) at the depth of 55 - 65 cm and to record the Mean Flow Velocity (MFV)
2. An intraparenchymal ICP monitor (Codman electrode MicroSensor, Johnson & Johnson Medical, Ltd., USA) was used for monitoring the ICP. The ICP monitor was connected to a Hewlett Packard (HP) monitor, model 66s-M116A, via a pressure transducer (Codman neuromonitor interface control unit, 82-6605) and module
3. An intraparenchymal brain tissue combined Oxygen and Temperature probe (LICOX REFIT2, Integra NeuroSciences, Ltd., England) was used for monitoring the PtO₂. The PtO₂ probe monitor is placed in the healthy side of brain parenchyma. The signals are transmitted to a HP monitor, via a transducer box (LICOX REF POM.BOX, Integra NeuroSciences, Ltd., England) and monitor cable (LICOX REF NL950-MC-01, Integra NeuroSciences, Ltd., England)

Standard management:

In the intensive care unit, patients were positioned in bed with an approximately 30° head up. Monitoring involved in simultaneous and continuous assessment of MAP from a radial or brachial arterial line, electrocardiogram, pulse oximetry values, and End-tidal Carbon Dioxide (EtCO₂) by capnography. Central venous pressure and body temperature were assessed either continuously or intermittently at frequent intervals. Routine pharmacological or physical measures were adopted to avoid fever if body temperature levels were 37.5°C or higher. Mechanical ventilation was adjusted to keep EtCO₂ between 30 and 33 mmHg. Sedation with propofol titration is used for ICP control. Patients underwent complete neurological assessment by nursing staff hourly or more frequently if the condition was deteriorating.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

1. Application of multiple cerebral blood monitor in the treatment of STBI could result in reduced mortality rate and impaired quality of life
2. Patients with impaired autoregulation, the optimal CPP value could be changed

Secondary outcome measures

1. Through this study, the result could be contributed to the establishment of a new Clinical Practice Guideline of STBI
2. The result could be used as a guideline to train young physicians
3. In different groups, patients with multiple cerebral monitoring will have better GOS, GOSE, HRQL, and survival

Overall study start date

01/09/2006

Completion date

31/12/2011

Eligibility

Key inclusion criteria

Glasgow Coma Scale (GCS) score three to eight will be enrolled in this study.

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

320

Key exclusion criteria

1. Patients manifesting severe systemic injury with hypotension (systolic blood pressure less than 90 mmHg on admission)
2. GCS score three with fixed and dilated pupils after resuscitation
3. GCS score three to four with family refused aggressive treatment
4. Patients transferred from another institution longer than 24 hours after injury
5. Those who were dead on arrival

Date of first enrolment

01/09/2006

Date of final enrolment

31/12/2011

Locations

Countries of recruitment

Taiwan

Study participating centre

No. 250 Wu-Hsing Street

Taipei

Taiwan

111

Sponsor information

Organisation

Department of Health, Taipei City Government (Taiwan)

Sponsor details

1, No.1, Shifu Road

Xinyi District

Taipei

Taiwan

110

+886 (0)2 2728 7070

yuhhuey@health.gov.tw

Sponsor type

Government

Website

<http://english.taipei.gov.tw/health/index.jsp>

ROR

<https://ror.org/02eqw3q87>

Funder(s)

Funder type

Government

Funder Name

Department of Health, Taipei City Government (Taiwan)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration