Is toothpaste containing xylitol + fluoride more effective than toothpaste containing only fluoride in preventing dental caries among preschoolers?

Submission date	Recruitment status No longer recruiting	Prospectively registered		
18/11/2011		☐ Protocol		
Registration date 06/01/2012	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 03/11/2022	Condition category Oral Health	[] Individual participant data		

Plain English summary of protocol

Background and study aims:

Dental caries (tooth decay) is the most common disease among pre-schoolers. Low-income pre-schoolers are more likely to be at risk for tooth decay. Current public health strategies have failed to significantly control tooth decay in this age group. This is an important problem to address because untreated tooth decay leads to pain, infection, which can lead to poor academic outcomes, systemic health problems, cellulitis, and in rare cases, death.

Currently, fluoride toothpaste is used to prevent tooth decay in pre-schoolers. Studies suggest that xylitol, a naturally-occurring sweetener, also helps to prevent tooth decay. Based on the idea that adding xylitol to fluoride toothpaste leads to a more effective product, toothpaste manufacturers have developed a xylitol+fluoride toothpaste. However, until now, no studies have been conducted that compare fluoride toothpaste to xylitol+fluoride toothpaste as a way to prevent tooth decay in pre-schoolers. The Head Start Program is a national program targeted at low-income pre-schoolers ages 3-5 years. The aim of the program is to prepare vulnerable pre-schoolers for kindergarten by ensuring good overall health. Many Head Start Programme classrooms include a tooth brushing program that is overseen by the classroom teacher. Teachers are trained to ensure the pre-schoolers follow the correct tooth brushing method. Currently, all Head Start classrooms that have a tooth brushing program use fluoride toothpaste.

Who can participate?

Pre-schoolers are healthy and that consent is received for participation by the pre-schoolers caregiver.

What does the study involve?

This study will introduce xylitol-fluoride toothpaste to two Head Start classrooms and fluoride-toothpaste to two other Head Start classrooms. Our aim is to compare tooth decay rates for preschoolers in the two groups, from the start of the study to 9 months. We will also collect plaque and saliva samples at 9 months to see if the two groups differ in the levels of Streptococcus mutans, the bacteria which is responsible for plaque and tooth decay.

What are the possible benefits and risks of participating?

Pre-schoolers participating may be less likely to experience tooth decay. All tooth brushing activities were directly supervised by multiple classroom teachers, so the risks to the pre-schoolers were minimal.

Where is the study run from?

From four Head Start Programme classrooms in Majuro in the Republic of the Marshall Islands (50 pre-schoolers per classroom).

When is the study starting and how long is it expected to run for? The study took place during the 2010/2011 school year.

Who is funding the study?

United States Health Resources and Services Administration - Targeted Oral Health Services Systems.

Who is the main contact? Dr Ohnmar Tut ohnmar@u.washington.edu

Contact information

Type(s)

Scientific

Contact name

Dr Ohnmar Tut

Contact details

Republic of the Marshall Islands Ministry of Health Post Office Box 16 Majuro Marshall Islands 96960

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

A clinical trial comparing fluoride toothpaste and xylitol+fluoride toothpaste in reducing dental caries among pre-schoolers at increased risk for tooth decay: a randomised prospective trial

Study objectives

Xylitol+fluoride toothpaste is more effective than fluoride toothpaste at slowing the rates of tooth decay progression in pre-schoolers

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Single-center double-blinded classroom randomized prospective clinical trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Dental caries

Interventions

Arm 1: supervised in-class toothbrushing once/day with xylitol+1400ppm fluoride toothpaste Arm 2: supervised in-class toothbrushing once/day with 1400ppm fluoride toothpaste

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Total increase in the number of primary tooth surfaces with untreated dental caries from baseline to end of school year

Key secondary outcome(s))

Levels of intraoral plaque and salivary mutans streptococci (at end of school year)

Completion date

30/06/2011

Eligibility

Key inclusion criteria

- 1. Enrolled in one of four pre-schools in Majuro, Marshall Islands
- 2. Under age 5
- 3. Parent consent to participate in the trial

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Upper age limit

5 years

Sex

All

Total final enrolment

196

Key exclusion criteria

Allergy or sensitivity to fluoride toothpaste

Date of first enrolment

01/09/2010

Date of final enrolment

30/06/2011

Locations

Countries of recruitment

Marshall Islands

Study participating centre Republic of the Marshall Islands

Majuro Marshall Islands 96960

Sponsor information

Organisation

Republic of the Marshall Islands (Marshall Islands)

Funder(s)

Funder type

Government

Funder Name

United States Health Resources and Services Administration (HRSA) (USA) - Targeted Oral Health Services Systems TOHSS Grant ref: H47MC08647

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/01/2014	03/11/2022	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes