Short TEG: cessation of clopidogrel following insertion of drug-eluting coronary stents

Submission date	Recruitment status	Prospectively registered
23/04/2010	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
23/04/2010	Completed	[X] Results
Last Edited	Condition category Circulatory System	[] Individual participant data
22/07/2013	Circulatory System	

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Nicholas Curzen

Contact details

Southampton General Hospital Cardiology Research Wessex Cardiac Unit, Mail Point 46 Southampton United Kingdom SO16 6YD

Additional identifiers

Protocol serial number 7907

Study information

Scientific Title

An investigation into the effects of cessation of clopidogrel therapy on vascular inflammation and platelet reactivity in patients with drug-eluting coronary stents: is there a "rebound phenomenon"?

Study objectives

The hypothesis of this trial is that cessation of clopidogrel maintenance therapy after 12 months in patients receiving drug-eluting stents is associated with a pro-inflammatory and pro-thrombotic response that offers an explanation for the observed clustering of clinical events, including stent thrombosis, that have been described in the first 90 days after clopidogrel is stopped.

Antiplatelet agents (drugs used to prevent platelets from binding together and forming blood clots) are widely used in the prevention and treatment of cardiovascular disease. Aspirin and clopidogrel are essential antiplatelet drugs required in all patients undergoing percutaneous coronary intervention (PCI) where narrowed arteries are stretched open with balloons and stents to improve the blood supply to the heart. Previous studies have demonstrated a wide range of patient response to antiplatelet agents. Poor responders are at increased risk of complications including potentially fatal stent thrombosis (blood clot forming within the stent), particularly in the period immediately after cessation of clopidogrel. A large observational study has reported a clustering of events, including death and heart attacks within the first 90days of stopping clopidogrel, raising the possibility of a clopidogrel "rebound" effect. The reason for this effect is unknown and it has been proposed that it may be due to an increase in inflammatory markers and platelet activation following withdrawal of clopidogrel.

The aim of our study is to investigate whether clopidogrel cessation 12 months after PCI causes a rebound proinflammatory and prothrombotic effect. We will investigate the mechanism behind this phenomenon and whether it is related to the lack of synergistic effect of clopidogrel on responses to aspirin when clopidogrel treatment is withdrawn. We will also assess whether the response differs in diabetic patients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Southampton and South West Hampshire Research Ethics Committee (A) approved on the 5th November 2009 (ref: 09/H0502/106)

Study design

Randomised interventional screening clinical laboratory study

Primary study design

Interventional

Study type(s)

Screening

Health condition(s) or problem(s) studied

Topic: Cardiovascular; Subtopic: Cardiovascular (all Subtopics); Disease: Cardiovascular

Interventions

Patients are not randomised to any treatment in this trial. All enrolled patients are due to stop dual antiplatelet medication as part of their routine care. This study is assessing the effects of stopping clopidogrel at the end of a routine 12 months course of treatment. Patients are followed up in the study for approximately 7 weeks attending the hospital for 7 blood tests over that time.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Clopidogrel is routinely used for 12 months after drugeluting stent implantation

Key secondary outcome(s))

Any differences in inflammatory markers, measured at 12 months

Completion date

01/11/2010

Eligibility

Key inclusion criteria

- 1. Aged greater than 18 years, either sex
- 2. Signed written informed consent
- 3. Prior percutaneous coronary intervention (PCI) with drugeluting stent implantation and due to stop clopidogrel 12 months after PCI
- 4. On maintenance dose aspirin 75 mg and clopidogrel 75 mg daily
- 5. 30 diabetics and 30 nondiabetics

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Aged greater than 85 years
- 2. On regular nonsteroidal antiinflammatory medication or steroids

Date of first enrolment

01/01/2010

Date of final enrolment

01/11/2010

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Southampton General Hospital
Southampton
United Kingdom
SO16 6YD

Sponsor information

Organisation

Southampton University Hospitals NHS Trust (UK)

ROR

https://ror.org/0485axj58

Funder(s)

Funder type

Industry

Funder Name

Haemonetics Limited (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type

Details results

Date created Date added Peer reviewed? Patient-facing?

Results article 01/10/2011 Yes No

Participant information sheet Participant information sheet 11/11/2025 No Yes