

Parkinson's disease, exercise and cognitive training

Submission date 23/03/2016	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 14/04/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 22/01/2018	Condition category Nervous System Diseases	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Parkinson's disease (PD) is a long-term medical condition which is caused by the gradual loss of nerve cells (neurons) in a part of the brain called the substantia nigra. These neurons are normally responsible for producing dopamine, a chemical messenger (neurotransmitter) which carries signals around the brain that help to coordinate movement. In people suffering from PD, these neurons gradually die over time, causing the level of dopamine in the brain to gradually fall. As the levels of dopamine become lower, the brain is unable to coordinate movement as effectively, causing abnormal movements such as stiffness, tremor (uncontrollable shaking) and slowness of movement (bradykinesia). As well as the issues with movement, many patients with PD also experience problems with cognitive function (thinking, reasoning and memory). In the early stages of the disease medications can help to reduce the physical symptoms, but the effects of these drugs on cognitive function are not fully understood. Several studies have suggested that a combination of both physical exercise and cognitive training can help to improve cognitive function in PD patients more efficiently than cognitive training alone, however further research is needed in order to find out how effective this really is. The aim of this study is to find out whether a programme involving both physical exercise and cognitive training is more effective at improving cognitive function than cognitive training alone.

Who can participate?

Adults aged between 40 and 80 who have Parkinson's disease with mild to moderate disability.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the second group take part in cognitive training alone. This involves completing a number of mental tasks on a computer of increasing difficulty. The difficulty levels are adapted for each patient. Those in the second group take part in cognitive training with exercise. This involves completing the cognitive training on a touch screen computer while walking on a treadmill. Two weeks before and after the training and again two months later, participants in both groups complete a number of assessments and questionnaires to test their cognitive and physical performance.

What are the possible benefits and risks of participating?
Participants may benefit from being able to improve their cognitive function. There are no notable risks involved with taking part in this study.

Where is the study run from?
University Hospital Complex of A Coruña (Spain)

When is the study starting and how long is it expected to run for?
December 2014 to February 2018

Who is funding the study?
Ministry of Economy and Competitiveness (Spain)

Who is the main contact?
Professor Miguel Fernández del Olmo
mafo@udc.es

Contact information

Type(s)
Scientific

Contact name
Prof Miguel Fernández del Olmo

ORCID ID
<https://orcid.org/0000-0001-5325-503X>

Contact details
Departamento de Educación Física e Deportiva
Facultade de Ciencias do Deporte e a Educación Física (INEF Galicia)
Avd. Ernesto Che Guevara 121, Pazos-Liáns
A Coruña
Spain
15179
+34 981167000
mafo@udc.es

Additional identifiers

Protocol serial number
DEP2014-53896-R

Study information

Scientific Title
Effects of the combination of physical exercise and cognitive training on cognitive functions in patients with Parkinson's disease

Acronym

EFECEP

Study objectives

Multimodal intervention of physical exercise and cognitive training will induce greater cognitive improvements compared with the unimodal intervention of cognitive training in Parkinson's disease patients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics committee of University of A Coruña, 30/12/2015, ref: 20/2014

Study design

Single-centre randomised parallel trial

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Parkinson's disease

Interventions

Patients will be allocated randomly in two groups, a group of multimodal intervention (physical exercise + cognitive training) and a group of unimodal intervention (cognitive training).

Group 1: Participants complete cognitive training alone. This involves completing computer tasks, individualized for the cognitive capacity, using the software SmartBrain (<http://www.smartbrain.net/>). Participants complete three sessions a weeks for eight weeks, lasting for around 64 minutes per session.

Group 2: Participants complete the cognitive training whiles walking on a treadmill, by using touch screen technology. Participants complete three sessions a weeks for eight weeks, lasting for around 64 minutes per session.

Two weeks before and after the training programs, clinical, motor, neuropsychological and neurophysiological evaluations will be conducted. A two month follow up will also be performed.

Intervention Type

Behavioural

Primary outcome(s)

1. Cognitive performance is measured using CDS (Cognitive Difficulties Scale), WAIS-III (Wechsler Adult Intelligence Scale-III), Digital Span Forward and backwards, Trail Making Test A and B, Stroop Test, Corsi Block, Verbal Fluency (FAS), Tower of London and Wilcoxin Card Sorting Test at 2 month before intervention, 2 weeks before intervention (baseline), 2 weeks post-intervention and two months post-intervention (follow-up)
2. Motor performance is measured using walk at comfortable speed, Time Up & Go and walk with

dual-task at 2 month before intervention, 2 weeks before intervention (baseline), 2 weeks post-intervention and two months post-intervention (follow-up)

3. Quality of life is measured using ADLS (Activities of Daily Living Scale), QOLS (Quality Of Life Scale) and IPAQ (International Physical Activity Questionnaire) at 2 month before intervention, 2 weeks before intervention (baseline), 2 weeks post-intervention and two months post-intervention (follow-up)

Key secondary outcome(s)

1. Functional connectivity is measured using evoked related potentials at 2 weeks before intervention (baseline) and 2 weeks post-intervention
2. White matter is measured using diffusion tensor imaging at 2 weeks before intervention (baseline) and 2 weeks post-intervention

Completion date

15/01/2018

Eligibility

Key inclusion criteria

1. Aged 40–80 years
2. Diagnosed with PD according to UK PD Society Brain Bank Criteria.26
3. Disease severity of Hoehn and Yahr (H&Y) stages I–III

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Sex

All

Key exclusion criteria

1. Significant cognitive impairment (Mini-Mental State Examination < 23)
2. Below average premorbid intelligence (vocabulary subtest, Wechsler Adult Intelligence Scale-III [WAIS-III] typical score < 40)
3. Major depression (GDS-15 > 10)
4. Severe auditory or visual deficits
5. Another psychiatric/neurological condition

Date of first enrolment

01/06/2016

Date of final enrolment

01/06/2017

Locations

Countries of recruitment

Spain

Study participating centre

University Hospital Complex of A Coruña (Complejo Hospitalario Universitario de A Coruña)

As Xubias, 84

A Coruña

Spain

15179

Sponsor information

Organisation

University of A Coruña

ROR

<https://ror.org/01qckj285>

Funder(s)

Funder type

Government

Funder Name

Ministerio de Economía y Competitividad

Alternative Name(s)

Ministry of Economy and Competitiveness, MINECO, MEC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Spain

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request