A comparison to two types of group behaviour therapy for obsessive compulsive disorder: A pilot randomised controlled trial of behaviour therapy and mindfulness-based behaviour therapy for OCD

| Submission date 30/01/2014 | Recruitment status No longer recruiting | [X] Prospectively registered[X] Protocol |
|-------------------------------|--|---|
| Registration date 30/01/2014 | Overall study status Completed | Statistical analysis plan[X] Results |
| Last Edited 11/01/2019 | Condition category Mental and Behavioural Disorders | [] Individual participant data |

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Clara Strauss

Contact details

Psychology Department Guildford United Kingdom GU2 7XH

_

C.Strauss@surrey.ac.uk

Additional identifiers

Protocol serial number 15996

Study information

Scientific Title

A comparison to two types of group behaviour therapy for obsessive compulsive disorder: A pilot randomised controlled trial of behaviour therapy and mindfulness-based behaviour therapy for OCD

Acronym

Mindfulness-based behaviour therapy for OCD: a pilot RCT

Study objectives

Obsessive compulsive disorder (OCD) affects up to one million adults in the UK (NICE, 2005). People diagnosed with OCD experience unwanted obsessive thoughts and engage in compulsive behaviours. Exposure and response prevention (ERP) is a behaviour therapy and it is the psychological therapy with the strongest evidence for effectiveness. However, only 60% of people benefit from ERP. Many people do not find ERP acceptable, with about 25% of people starting ERP dropping out.

Mindfulness-based interventions (MBIs) teach people to notice unpleasant thoughts and feelings without reacting unhelpfully. Although effective for depression, psychosis and a range of anxiety disorders, there is very little research on the effectiveness of MBIs for OCD, although the research that is published shows promise.

We conducted a focus group with six people diagnosed with OCD and who had experience of ERP and MBI. We had intended to compare MBI to ERP, however participants advised us to combine ERP with MBI. They suggested that effectiveness might be based by combining the two together. We have followed their advice in this study.

We intend to compare standard ERP group behaviour therapy (BT) to mindfulness-based ERP group behaviour therapy (MBBT) to see if mindfulness practice improves ERP outcomes and reduces drop-out. Before we can answer this question in a fully-sized trial we first need to conduct a pilot study. This pilot study is a small version of the fully-sized trial. It will tell us: (a) the approximate difference in outcome between BT and MBBT, (b) how easily we can recruit to the study, and (c) how many people drop out. This information will allow us to know how many participants we need for the full trial, how long recruitment will take and how to reduce dropout rates. Once we have this information we will apply for funding for the full trial.

Ethics approval required

Old ethics approval format

Ethics approval(s)

13/LO/1768

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Mental Health Research Network; Subtopic: Personality disorder; Disease: Personality disorders

Interventions

Behaviour Therapy: A 10 session exposure and response prevention for OCD group intervention (maximum 10 participants per group). Each session will run for two hours.

Mindfulness Behaviour Therapy: A 10 session mindfulness-enhanced exposure and response prevention for OCD group intervention (maximum 10 participants per group). Each session will run for two hours. Mindfulness practice and discussion will be added to the standard ERP protocol.

Follow Up Length: 6 month(s); Study Entry: Registration only

Intervention Type

Behavioural

Primary outcome(s)

Yale-Brown Obsessive Compulsive Scale - Second Edition (Y-BOCS-II; Goodman et al, 2006).; Timepoint(s): Baseline, post-therapy (10 weeks post-baseline) and 6-months post-therapy

Key secondary outcome(s))

- 1. Beck Depression Inventory second edition (BDI-II; Beck et al, 1996)
- 2. Change Interview (Elliott et al, 2001).; Timepoint(s): Post-therapy only
- 3. Five-Facet Mindfulness Questionnaire Short Form (FFMQ-SF; Bohlmeijer et al, 2011)
- 4. Mini Neuropsychiatric Interview 6.0.0; Timepoint(s): Baseline, post-intervention and six months post-therapy
- 5. Obsessional Beliefs Questionnaire Revised OBQ-44; Timepoint(s): Baseline, post-therapy and six months post-therapy
- 6. Short Warwick-Edinburgh Mental Well-Being Scale (2007); Timepoint(s): Baseline, post-therapy and 6-months post-therapy
- 7. Therapy attrition; Timepoint(s): Percentage of participants who complete their intervention will be recorded as a primary outcome

Completion date

31/12/2014

Eligibility

Key inclusion criteria

- 1. Meet diagnostic criteria for OCD (MINI 6.0.0 interview)
- 2. Have been stable on psychiatric medication for at least three months prior to the consent meeting, or not taking psychiatric medication for at least three months prior to the consent meeting
- 3. Have no plans for changes to psychiatric medication during the course of the study
- 4. Have not received psychological therapy in the past three months or to have any plans for psychological therapy during the course of the study
- 5. Be aged over 18 years of age

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Αll

Key exclusion criteria

Exclusion criteria will be that participants will not have an identified organic cause for their OCD symptoms or a diagnosed learning disability. This will be ascertained through the care team. People will be excluded if they meet diagnostic criteria, based on the MINI interview, for; post-traumatic stress disorder, anorexia nervosa, an autistic spectrum disorder or a psychotic disorder.

To reflect the reality of mental health services and the co-morbidity of OCD with other anxiety disorders and with depressive disorders (NICE, 2005), co-morbidity with these conditions will not be an exclusion criterion. The exception to this is that people meeting diagnostic criteria for PTSD will be excluded given some concerns in the MBI literature about the offering MBIs to this population. People presenting with hoarding only compulsions will be excluded from the study, given the recent move to classify hoarding as a distinct condition from OCD (Pertusa et al, 2008).

Date of first enrolment

01/02/2014

Date of final enrolment

31/12/2014

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
University of Surrey
Guildford
United Kingdom
GU2 7XH

Sponsor information

Organisation

Sussex Partnership NHS Foundation Trust (UK)

ROR

https://ror.org/05fmrjg27

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research: Research for Patient Benefit (RfPB)

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created Date a | added Peer reviewed | d? Patient-facing? |
|-------------------------------|-------------------------------|---------------------|---------------------|--------------------|
| Results article | results | 01/06/2018 | Yes | No |
| <u>Protocol article</u> | protocol | 16/04/2015 | Yes | No |
| HRA research summary | | | /2023 No | No |
| Participant information sheet | Participant information sheet | 11/11/2025 11/11 | /2025 No | Yes |