

Randomised, Double-blinded, Placebo-controlled Clinical Trial of Pandemic Influenza Inactive Vaccine on Healthy Subjects

Submission date
25/07/2006

Recruitment status
No longer recruiting

☐ Prospectively registered

☐ Protocol

Registration date
17/08/2006

Overall study status
Completed

☐ Statistical analysis plan

☒ Results

Last Edited
03/05/2019

Condition category
Infections and Infestations

☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Jiangtao Lin

Contact details

Department of Respiratory Internal
China-Japan Friendship Hospital
Hepinli Yinhua East Road
Chaoyang District
Beijing
China
100029
+86 106 422 1122 2313
jiangtao_L@263.net

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

NCT00356798

Secondary identifying numbers

PRO-PanFlu-1001

Study information

Scientific Title

Randomised, Double-blinded, Placebo-controlled Clinical Trial of Pandemic Influenza Inactive Vaccine on Healthy Subjects

Acronym

RDPCTPIIVHS

Study objectives

An inactivated monovalent A/H5N1 (influenza A virus) whole-virion, aluminum hydroxide adjuvated influenza vaccine will be safe and immunogenic in humans.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval gained from the ethical review committee of China Japan Friendship Hospital dated on 12 December 2005 (reference: 2005NO[37]Total[99]).

Study design

A stratified, randomised, placebo-controlled and double-blind phase I clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Prevention

Participant information sheet

Health condition(s) or problem(s) studied

Pandemic influenza

Interventions

The trial is divided into four dosage groups and processed from low dosage vaccine to high dosage vaccine:

1. Group A: 30 subjects will be randomly injected with vaccines among which 24 subjects will be injected with the trial vaccine (1.25 µg) antigen and the other six subjects will be randomised into a placebo group. All subjects will be observed for 30 minutes for immediate reactions. Local reactions and systemic reactions were then observed at six, 24, 48 and 72 hours. The second dose will be injected 28 days after the first dose. The dosage, injection and observation method are the same as the first one. Subjects will be followed up for 210 days. Serum will be collected before the injection and then at 14, 28, 42, 56, 210 days after injection. The serums will be tested by the Haemagglutination-Inhibition (HI) assay.

2. Group B: 72 hours after the injection of Group A, when the safety of the vaccine is confirmed, 30 subjects will be randomly injected with vaccines, among them, 24 subjects will receive the trial vaccine (2.5 µg) and the other six subjects will be receive the control. The methods of safety observation and blood collection are as same as Group A.

3. Group C: 72 hours after the injection of Group B, when the safety of vaccine is confirmed, 30 subjects will be randomly injected with the vaccine. In the same way, 24 subjects will receive the trial vaccine (5.0 µg) and the other six subjects will receive the control. The methods of safety observation and blood collection are as same as Group A.

4. Group D: 72 hours after injection for the Group C, when the safety of vaccine is confirmed, 30 subjects will be randomly injected with vaccines. Among them, 24 subjects will receive the trial vaccine (10.0 µg) and the other six subjects will receive the control. The methods of safety observation and blood collection are as same as Group A.

Trial vaccine is designed 0.5 ml per dose. The administration regimen is designed as a two dose schedule, which are given at day zero and day 28.

Intervention Type

Drug

Phase

Phase I

Drug/device/biological/vaccine name(s)

Influenza A virus vaccine

Primary outcome measure

To evaluate the safety of pandemic inactivated influenza vaccine by different doses.

Secondary outcome measures

To evaluate the immunogenicity of pandemic inactivated influenza vaccine by different doses.

Overall study start date

20/12/2005

Completion date

05/06/2006

Eligibility

Key inclusion criteria

1. Males and females, aged from 18 to 60 years old
2. Able to provide proof of identity to the satisfaction of the study clinician completing the enrolment process
3. Able and willing to complete the informed consent process

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

60 Years

Sex

Both

Target number of participants

120

Total final enrolment

120

Key exclusion criteria

1. Women who are breast-feeding or planning to become pregnant during the following 210 days of study participation
2. Subjects who engage in the occupations of culturist, slaughter, sale and forwarder of any avian organisms
3. Subject has a medical history of any of the following:
 - a. allergic history, or allergic to any ingredient of vaccine, such as egg, egg protein etc.
 - b. serious adverse reactions to vaccines such as anaphylaxis, hives, respiratory difficulty, angioedema, or abdominal pain
 - c. autoimmune disease or immunodeficiency
 - d. asthma that is unstable or required emergency care, urgent care, hospitalisation or intubation during the past two years or that requires the use of oral or intravenous corticosteroids
 - e. diabetes mellitus (type I or II), with the exception of gestational diabetes
 - f. history of thyroidectomy or thyroid disease that required medication within the past 12 months
 - g. serious angioedema episodes within the previous three years or requiring medication in the previous two years
 - h. bleeding disorder diagnosed by a doctor (e.g. factor deficiency, coagulopathy, or platelet disorder requiring special precautions) or significant bruising or bleeding difficulties with Intramuscular (IM) injections or blood draws
 - i. malignancy that is active or treated malignancy for which there is not reasonable assurance of sustained cure or malignancy that is likely to recur during the period of study
 - j. seizure disorder other than febrile seizures under the age of two, seizures secondary to alcohol withdrawal more than three years ago, or a singular seizure not requiring treatment within the last three years

- k. asplenia, functional asplenia or any condition resulting in the absence or removal of the spleen
- l. Guillain-Barre Syndrome (GBS)
- 4. The abnormal result of laboratory tests as below:
 - a. biochemistry assaying: Alanine Aminotransferase (ALT)/Serum Glutamate Pyruvate Transaminase, Total Bilirubine (TBIL), Direct Bilirubine (DBIL), Blood Urea Nitrogen (BUN) and Creatinine (Cr)
 - b. Routine blood assaying, routine urine assaying
 - c. Hepatitis B surface Antigen (HBsAg) positive
 - d. pregnancy test positive
- 5. Subject has received any of the following substances:
 - a. immunosuppressive medications or cytotoxic medications or inhaled corticosteroids within the past six months (with the exception of corticosteroid nasal spray for allergic rhinitis or topical corticosteroids for an acute uncomplicated dermatitis)
 - b. blood products within three months prior to initial study vaccine administration
 - c. other study drug within 30 days prior to initial study vaccine administration
 - d. live attenuated vaccines within 30 days prior to initial study vaccine administration
 - e. medically indicated subunit or killed vaccines, e.g. pneumococcal, or allergy treatment with antigen injections, within 14 days of study vaccine administration
 - f. current anti-tuberculosis prophylaxis or therapy
- 6. Fever before vaccination, axillary temperature 37.0°C
- 7. Psychiatric condition that precludes compliance with the protocol, past or present psychoses, past or present bipolar disorder requiring therapy that has not been well controlled on medication for the past two years, disorder requiring lithium, or suicidal ideation occurring within five years prior to enrolment
- 8. Any medical, psychiatric, social condition, occupational reason or other responsibility that, in the judgment of the investigator, is a contraindication to protocol participation or impairs a volunteer's ability to give informed consent

Date of first enrolment

20/12/2005

Date of final enrolment

05/06/2006

Locations

Countries of recruitment

China

Study participating centre

Department of Respiratory Internal

Beijing

China

100029

Sponsor information

Organisation

Sinovac Biotech Co. Ltd (China)

Sponsor details

No. 39 Shangdi Xi Road

Haidian District

Beijing

China

100085

+86 108 289 0088

sinovac@sinovac.com

Sponsor type

Industry

Website

<http://www.sinovac.com>

ROR

<https://ror.org/057f25d66>

Funder(s)**Funder type**

Government

Funder Name

The study was funded by a grant (2005BA723B02) from Ministry of Science and Technology of the People's Republic of China.

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	16/09/2006	03/05/2019	Yes	No