

Comparison of the effects of spontaneous breathing trial with T-piece versus pressure-support ventilation in patients recovering from cardiac surgery: a randomised controlled pilot study

Submission date	Recruitment status	<input type="checkbox"/> Prospectively registered
28/09/2007	No longer recruiting	<input type="checkbox"/> Protocol
Registration date	Overall study status	<input type="checkbox"/> Statistical analysis plan
28/09/2007	Completed	<input type="checkbox"/> Results
Last Edited	Condition category	<input type="checkbox"/> Individual participant data
10/11/2011	Surgery	<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Ms Lee Moi Hew

Contact details

ITU

London Chest Hospital

Bonner Road

London

United Kingdom

E2 9JX

+44 (0)20 8983 2209

leemoi.hew@bartsandthelondon.nhs.uk

Additional identifiers

Protocol serial number

N0205182184

Study information

Scientific Title

Study objectives

To compare the mean levels of anxiety reported by two groups of patients using a Visual Analogue Scale (VAS) at specific time points during spontaneous breathing trials with:

1. The T-piece method, or
2. Pressure-support ventilation (PSV) of 10 cm H₂O with positive end expiratory pressure (PEEP) 5 cm H₂O

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled pilot study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Spontaneous breathing after ventilation

Interventions

Patients are randomised to:

1. The T-piece method
2. Pressure-support ventilation (PSV) of 10 cm H₂O with positive end expiratory pressure (PEEP) 5 cm H₂O

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome(s)

Anxiety scores before operation, at PSV of 10 cm H₂O with PEEP of 5 cm H₂O, 30-minute after PSV of 10 cm H₂O with PEEP of 5 cm H₂O, before extubation, one hour after extubation and before discharge from ICU.

Key secondary outcome(s))

No secondary outcome measures

Completion date

30/09/2006

Eligibility

Key inclusion criteria

Not provided at time of registration

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

22/03/2006

Date of final enrolment

30/09/2006

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

ITU

London

United Kingdom

E2 9JX

Sponsor information

Organisation

Funder(s)

Funder type

Government

Funder Name

Barts and The London NHS Trust (UK)

Funder Name

NHS R & D Support Funding (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration