

# To determine the effectiveness of amitriptyline compared to placebo in the management of moderate to severe HIV related peripheral neuropathy.

<b>Submission date</b> 06/11/2014	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 17/11/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 15/05/2015	<b>Condition category</b> Nervous System Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

The peripheral nervous system is part of the nervous system that is made up of the nerves and ganglia (group of nerve cells) outside of the brain and spinal cord (the central nervous system). It connects the central nervous system to the limbs and organs of the body. Damage to peripheral nerves, especially the long nerves of the feet and hands, is common in people infected with HIV. This damage may be caused by the virus itself, some of the antiretroviral drugs used to treat HIV/AIDS, or as a result of opportunistic infections, for example thrush, herpes and tuberculosis. The nerve damage is often painful, can have a detrimental effect on a person's quality of life and has significant socioeconomic effects. Unlike other types of peripheral neuropathy (e.g., the neuropathy that develops in people with diabetes mellitus), there are, as yet, no effective treatments for painful HIV neuropathy. Here, we want to see if amitriptyline, an antidepressant drug that has been shown to relieve pain in other types of neuropathy, can provide pain relief better than that of a placebo (a dummy pill) in patients with painful HIV neuropathy.

### Who can participate?

Adult HIV-infected individuals that have been diagnosed with painful HIV-associated sensory neuropathy. Participants must either be on stable antiretroviral therapy or have never been exposed to antiretroviral therapy for HIV infection.

### What does the study involve?

HIV-positive patients attending clinics at Chris Hani Baragwanath Hospital, Soweto, South Africa and that can walk are randomly allocated to receive amitriptyline (25 mg) or a placebo for six weeks. The drug/placebo dose is increased every three days over the first two weeks of the treatment until participants achieve pain relief, intolerable side effects, or a maximum dose of 150 mg per day. The trial drug and the placebo are identical in appearance. Participants visit the study centre every three weeks to receive their medication and to record a pain score. At the end of the first six-week period, participants are taken off their assigned medication for a three-week 'washout' period. After this washout period, those participants who receive amitriptyline

before are now given the placebo and vice versa for another 6 weeks. Dose titration and pain assessments will follow the same protocol used in the first six-week period.

What are the possible benefits and risks of participating?

Some participants may achieve meaningful pain relief when receiving amitriptyline, a drug recommended as first-line therapy for other types of painful peripheral neuropathy. However, any positive results of this study will mostly benefit the large population of HIV-infected patients with painful peripheral neuropathy, for which there are no proven treatments. Risks of participating in the trial include side-effects of the medication, typically sedation, dry mouth and difficulty urinating.

Where is the study run from?

Nthabiseng HIV Clinic and the Centre for Palliative Care, Chris Hani Baragwanath Hospital, Soweto, (South Africa)

When is the study starting and how long is it expected to run for?

April 2009 to April 2010

Who is funding the study?

The Diana Princess of Wales Memorial Fund (UK)

Who is the main contact?

Dr Peter Kamerman

peter.kamerman@wits.ac.za

Dr Natalya Dinat

natalyadinat5@gmail.com

## Contact information

**Type(s)**

Scientific

**Contact name**

Dr Natalya Dinat

**Contact details**

86, 14th Street  
Parkhurst  
Johannesburg  
South Africa  
2193

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

N/A

## Study information

### Scientific Title

Double blind randomised controlled trial to assess the effectiveness of amitriptyline compared to placebo in the management of moderate to severe HIV related peripheral neuropathy.

### Study objectives

Amitriptyline is superior to placebo in the management of moderate to severe HIV related peripheral neuropathy.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Human Research Ethics Committee, University of the Witwatersrand, South Africa, 16/09/2014, ref: M080709

### Study design

Placebo-controlled double-blind crossover group randomised controlled trial

### Primary study design

Interventional

### Secondary study design

Randomised cross over trial

### Study setting(s)

Hospital

### Study type(s)

Treatment

### Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

### Health condition(s) or problem(s) studied

Pharmacological management of painful neuropathy

### Interventions

All study participants will receive amitriptyline (titrated to efficacy, side effects or a maximum dose of 150 mg daily) or inert placebo in random order for 6 weeks, with a three-week washout period between interventions.

### Intervention Type

Drug

### Phase

Not Applicable

**Drug/device/biological/vaccine name(s)**

Amitriptyline

**Primary outcome measure**

Self-reported pain intensity, assessed on an 11-point numerical pain rating scale.

**Secondary outcome measures**

1. Pain interference on activities of daily living, assessed using the pain interference sub-scale of the Brief Pain Inventory
2. Physical symptoms, psychological, emotional and spiritual, and information and support needs, assessed using the African Palliative Care Outcomes Scale (APOS)

Primary and secondary outcomes were measured at baseline, three and six weeks for each of the two intervention periods. Across the duration of the study, that represents measurements at 0, 3, 6, 9, 12, and 15 weeks on each participant.

**Overall study start date**

01/04/2009

**Completion date**

30/04/2010

## **Eligibility**

**Key inclusion criteria**

1. Confirmed and documented HIV positive status, either on Arv therapy or not on ARV therapy
2. Diagnosed with moderate to severe peripheral neuropathy using the BPNS and DN4 screening tools and scoring  $\geq 4$  on an 11 point Likert scale visual analogue scale
3. Age above 18 on the date of recruitment
4. Ability to mentally and physically understand to make informed choice to participate in the study

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

124

**Key exclusion criteria**

All conditions which are likely to cause confounding will be excluded from the study, together with

conditions which are likely to cause drug interactions.

1. Severe pain from DSN caused by ARV drugs which warrants a change in regime
2. Already taking amitriptyline drug, or has taken the drugs at least three weeks prior
3. Limb amputation
4. Kaposi sarcoma of the lower limbs
5. Current post-herpetic neuralgia or herpes zoster
6. Pregnancy or intention to fall pregnant
7. TB treatment
8. Malignancy not related to HIV
9. Major psychiatric disorders including mania or epilepsy
10. Clinically significant renal failure
11. Diabetic neuropathy
12. Clinically significant liver failure or past history as defined by encephalopathy, oedema and jaundice
13. Patient in extreme pain or exhausted
14. Participating on another trial or study
15. MAOI, other tricyclic antidepressants or anti-epileptic drugs
16. Recent myocardial infarction, arrhythmias, heart block
17. History of urinary retention, urinary hesitancy or closed angle glaucoma

**Date of first enrolment**

01/04/2009

**Date of final enrolment**

30/04/2010

## **Locations**

**Countries of recruitment**

South Africa

**Study participating centre**

**86, 14th Street**

Johannesburg

South Africa

2193

## **Sponsor information**

**Organisation**

Diana Princess of Wales Memorial Fund (UK)

**Sponsor details**

The Royal Foundation of The Duke and Duchess of Cambridge and Prince Harry  
Kensington Palace  
Palace Green  
London  
United Kingdom  
W8 4PU

**Sponsor type**

Charity

## Funder(s)

**Funder type**

Charity

**Funder Name**

Diana Princess of Wales Memorial Fund (UK)

## Results and Publications

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	14/05/2015		Yes	No