The effect of pre-operative warming on wound infection rates after clean surgery

Submission date	Recruitment status	Prospectively registered
01/03/2001	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
01/03/2001	Completed	[X] Results
Last Edited	Condition category	[] Individual participant data
07/01/2010	Surgery	

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Mr Andrew Melling

Contact details

Professorial Unit of Surgery
North Tees and Hartlepool NHS Trust
North Tees General Hospital
Stockton-on-Tees
United Kingdom
TS19 8PE
+44 (0)1642 624087
AndyMelling@compuserve.com

Additional identifiers

Protocol serial number AP0728

Study information

Scientific Title

Laser doppler flowmetry of skin circulation to measure the effects of using two warming therapies: a randomised clinical trial of surgical patients

Study objectives

To assess whether warming patients before short duration, clean surgery will reduce infection rates.

Ethics approval required

Old ethics approval format

Ethics approval(s)

No ethics approval information required at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Infection during surgery

Interventions

Patients were randomised to one of three groups:

- 1. Standard treatment no warming
- 2. Local warming a minimum of 30 minutes localised warming to the potential incision site using a waming dressing
- 3. Systemic warming a minimum of 30 minutes warming to the whole body using a warm blown air blanket

Both warming devices were applied prior to surgery. A cohort of patients had laser doppler and TcPO2 recordings taken prior to warming, after warming and after surgery.

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

Wound infection; masked outcome assessments made at 2 and 6 weeks

Key secondary outcome(s))

Added as of 07/01/2010:

- 1. Skin preparation; masked outcome assessments made at 2 and 6 weeks
- 2. Length of operation; masked outcome assessments made at 2 and 6 weeks
- 3. Administration of prophylactic antibiotics; masked outcome assessments made at 2 and 6 weeks
- 4. Experience of the operating surgeon; masked outcome assessments made at 2 and 6 weeks
- 5. Core temperature, measured before any treatment, after any warming, and after surgery

Completion date

31/12/2001

Eligibility

Key inclusion criteria

1. Patients having clean breast, varicose vein or hernia surgery

Added as of 07/01/2010:

2. Surgery results in a scar longer than 3 cm in length

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Added as of 07/01/2010:

- 1. Under the age of 18 years
- 2. Pregnant
- 3. Taking long-term oral steroids
- 4. Received radiotherapy (to the incision site) or chemotherapy in the last 4 weeks
- 5. Had an infection at the time of surgery

Date of first enrolment

01/01/2001

Date of final enrolment

31/12/2001

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Professorial Unit of Surgery

Stockton-on-Tees United Kingdom TS19 8PE

Sponsor information

Organisation

Action Medical Research (UK)

ROR

https://ror.org/01wcqa315

Funder(s)

Funder type

Charity

Funder Name

Action Medical Research (UK)

Alternative Name(s)

action medical research for children, actionmedres, The National Fund for Research into Crippling Diseases, AMR

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output typeDetailsDate createdDate addedPeer reviewed?Patient-facing?Results articleresults15/09/2001YesNo