

# Phase I study of oral administration of S 78454 in association with doxorubicin in patients with solid tumour

<b>Submission date</b> 02/09/2013	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 24/09/2013	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 18/04/2018	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

Not provided at time of registration and not expected to be available in the future

## Contact information

### Type(s)

Scientific

### Contact name

Dr Jean-Pierre Delord

### Contact details

Clinical Research Unit and Pharmacology Lab EA 3035  
Institut Claudius Regaud  
20-24, rue du Pont Saint Pierre  
Toulouse  
France  
31052

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

CL1-78454-005

# Study information

## Scientific Title

Phase I dose-escalation study of oral administration of S 78454 given with a fixed dose infusion of doxorubicin administered weekly 3 out of 4 weeks in patients with solid tumour

## Study objectives

To establish the safety profile and the recommended Phase II dose of S 78454 in combination with a fixed dose infusion of doxorubicin.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Ethics approval was obtained before recruitment of the first participants

## Study design

Multicentric non-randomised open dose escalation Phase I study

## Primary study design

Interventional

## Secondary study design

Non randomised controlled trial

## Study setting(s)

Hospital

## Study type(s)

Treatment

## Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

## Health condition(s) or problem(s) studied

Solid tumours

## Interventions

1. Capsules containing 20 mg and 100 mg of S 78454 / Oral use / Treatment duration is 6 cycles
2. Fixed dose infusion of 25 mg/m<sup>2</sup> of doxorubicin, weekly 3 out of 4 weeks / Treatment duration is 6 cycles

No control group is involved

## Intervention Type

Drug

## Phase

Phase I

**Drug/device/biological/vaccine name(s)**

S 78454, doxorubicin

**Primary outcome measure**

1. Dose-limiting toxicities (DLTs) and maximum tolerated doses (MTDs) at each visit - Methods used: blood samples, physical examination, electrocardiogram (ECG)
2. Safety profile of the combination at each visit

**Secondary outcome measures**

1. Tumour response at baseline evaluation every 2 cycles using the Response Evaluation Criteria In Solid Tumors (RECIST) guideline
2. Pharmacokinetic parameters during cycle 1 and 2 by blood samples
3. Pharmacodynamic parameters during cycle 1 by blood samples and tumour biopsies

**Overall study start date**

15/11/2010

**Completion date**

15/11/2013

**Eligibility****Key inclusion criteria**

1. Solid tumour, with measurable or evaluable disease, that has relapsed or is refractory to conventional, standard forms of therapy
2. Ability to swallow oral capsule(s) without difficulty
3. Estimated life expectancy > 12 weeks
4. Eastern Cooperative Oncology Group (ECOG) performance status < or equal to 1
5. Adequate haematological, renal and hepatic functions

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

36 patients

**Key exclusion criteria**

1. Major surgery within previous 4 weeks
2. Chemotherapy within previous 3 weeks (6 weeks in case of nitrosoureas)
3. Radiotherapy within previous 4 weeks (except for palliative radiotherapy at localised lesions)
4. Immunotherapy or hormonotherapy within previous 2 weeks, except stable oral glucocorticoid and mineralocorticoid replacement for adrenal insufficiency, or oral contraceptives or hormonal replacement therapy
5. Risk factors for, or use of drugs known to prolong QTc interval and that may be associated

with Torsades de Pointes

6. Patients treated by valproic acid within previous 5 days

7. Phenytoin (and by extension fosphenytoin) within previous 3 weeks

**Date of first enrolment**

15/11/2010

**Date of final enrolment**

15/11/2013

## **Locations**

**Countries of recruitment**

Belgium

France

**Study participating centre**

Clinical Research Unit and Pharmacology Lab EA 3035

Toulouse

France

31052

## **Sponsor information**

**Organisation**

Institut de Recherches Internationales Servier (France)

**Sponsor details**

50 rue Carnot

Suresnes

France

92284

**Sponsor type**

Industry

**Website**

<http://www.servier.com/>

**ROR**

<https://ror.org/034e7c066>

# Funder(s)

## Funder type

Industry

## Funder Name

Institut de Recherches Internationales Servier (France)

# Results and Publications

## Publication and dissemination plan

Publication plan:

Summary results are published in <https://clinicaltrials.servier.com>.

## Intention to publish date

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from <https://clinicaltrials.servier.com> if a Marketing Authorisation has been granted after 1st January 2014.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Basic results</a>				No	No