

# Scleroderma heart study

<b>Submission date</b> 09/11/2016	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 09/03/2017	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 31/01/2022	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Scleroderma is a rare disease affecting a few thousand patients in the UK. The cause is not known. Patients develop fibrosis (a kind of scar tissue) in the skin, making it tight and hard. Internal organs are often affected too, including the lungs, kidney and bowel. Scleroderma can affect the heart, which can be mild or severe and devastating. Because this complication is uncommon, little is known about how best to diagnose and treat this condition. Current practice is to use MRI scans and sometimes to take samples (biopsies) of the heart to make a diagnosis of scleroderma heart involvement. Treatment is often with drugs to suppress the immune system. However, it is not known how much inflammation (swelling) there is in the heart in patients with scleroderma or whether the treatment might work to suppress that inflammation. In this study, a type of scan called a PET-CT scan will be performed on patients who we suspect have heart involvement from their scleroderma. This scan is already widely used in patients with cancer and in some other conditions. It involves the injection into a regular vein of sugar with a tracer attached to it, so the scan can track where the sugar is taken up. The aim of this study is to find out if a PET-CT scan is an effective assessment to use in patients thought to have heart involvement from scleroderma.

### Who can participate?

Adults who have scleroderma with suspected heart involvement.

### What does the study involve?

All patients are recruited at the Scleroderma centre and National Pulmonary Hypertension Service at the Royal Free Hospital. After signing an informed consent form, participants undergo a heart assessment to check that they are suitable to take part. Following this, health information is taken and patients undergo a PET-CT scan and a pregnancy test (if female of child bearing potential). A follow up visit, involving similar assessments is carried out 6-9 months after screening if required.

### What are the possible benefits and risks of participating?

There are no direct benefits involved with participating. The main risk involved in this study will be the radiation dose received by patients. All patients included in this study will undergo a single 18FDG PET-CT scan. Patients with a positive scan may be offered one further scan after a course of treatment. The study protocol has been written in consultation with medical physics expert at the Royal Free who has reviewed and approved radiation doses. Additionally in order

to avoid harm to unborn children, patients of child bearing potential will have a pregnancy test at every trial visit. If the test is positive patients will be withdrawn from the study immediately.

Where is the study run from?  
Royal Free Hospital (UK)

When is the study starting and how long is it expected to run for?  
March 2015 to December 2020

Who is funding the study?  
Royal Free Charity (UK)

Who is the main contact?  
1. Ms Ivy Wanjiku (public)  
ivy.wanjiku@nhs.net  
2. Dr Benjamin Schreiber (scientific)  
benjamin.schreiber@nhs.net

## Contact information

**Type(s)**  
Public

**Contact name**  
Ms Ivy Wanjiku

**Contact details**  
Clinical Trials  
Rheumatology Department  
Lower Ground Floor  
Royal Free London NHS Foundation Trust  
London  
United Kingdom  
NW3 2QG  
+44 20 7317 7544  
ivy.wanjiku@nhs.net

**Type(s)**  
Scientific

**Contact name**  
Dr Benjamin Schreiber

**Contact details**  
Clinical Trials  
Rheumatology Department  
Lower Ground Floor  
Royal Free London NHS Foundation Trust  
London  
United Kingdom  
NW3 2QG

+44 20 7317 7544  
benjamin.schreiber@nhs.net

## **Additional identifiers**

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

9729

## **Study information**

### **Scientific Title**

A study of using 18FDG-PET to define presence of inflammation in scleroderma myocardial involvement

### **Study objectives**

Use of a combined 18FDG-PET-CT scan may aid in assessment of patients thought to have cardiac involvement from scleroderma.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

East Midlands - Derby Research Ethics Committee, 30/08/2016, ref: 16/EM/0292

### **Study design**

Observational cross-sectional study

### **Primary study design**

Observational

### **Secondary study design**

Cross sectional study

### **Study setting(s)**

Hospital

### **Study type(s)**

Diagnostic

### **Participant information sheet**

### **Health condition(s) or problem(s) studied**

Systemic sclerosis with heart involvement

## **Interventions**

All patients will be recruited at the Scleroderma centre and National Pulmonary Hypertension Service at the Royal Free Hospital.

### **Pre-treatment Evaluation**

Patients with suspected scleroderma cardiac involvement are followed up at the Scleroderma and Pulmonary Hypertension services at the Royal Free Hospital as part of standard practice, where a full 'Cardiac Evaluation' is undertaken; follow up visits are approximately 1-6 monthly depending on the clinical scenario. The Cardiac Evaluation includes rheumatic and cardiac history and examination, blood tests including autoantibodies, CK level, and cardiac biomarkers (troponin, NTproBNP), six minute walking distance, ECG, echocardiogram, 24 hour holter, an implantable loop heart rhythm recorder device in selected patients (those with ectopic beats on holter, presyncope or palpitations), cardiac MRI and cardiac catheterisation with assessment of coronary artery disease if not already excluded and right heart catheterisation as indicated. Patients with scleroderma cardiac involvement will often undergo cardiac biopsy -with samples sent for histology and to exclude viral myocarditis.

This standard rheumatic and cardiac assessment will form the baseline evaluation for patients included in this study. Any UK patient with a diagnosis of scleroderma with cardiac involvement who has undergone a cardiac evaluation at the Royal Free Hospital is potentially eligible for entry into the trial. The patient will be consented for the trial by an appropriately trained member of the Scleroderma and Pulmonary Hypertension services at the Royal Free Hospital. When patients are seen at the Scleroderma and Pulmonary Hypertension services at the Royal Free Hospital, they will be given the patient information sheet. All patients will be allowed as much time as needed to review the information sheet and ask any questions to the Scleroderma and Pulmonary Hypertension services team.

After completion of the informed consent form, a trial number will be assigned to each the patient. Following registration, Scleroderma and Pulmonary Hypertension services at the Royal Free Hospital will contact the nuclear medicine department to arrange a trial slot of imaging. The nuclear medicine department will undertake trial imaging within 6 weeks of the initial evaluation and consent procedures. If there is a  $\geq 6$  week delay before the imaging can be undertaken, patient has to be withdrawn from the trial. Such a patient would be eligible to be screened again and considered for inclusion if he/she fulfils the eligibility criteria. Once a patient has been registered onto the trial, the patient will be provided with a copy of their signed consent form.

## **Intervention Type**

Procedure/Surgery

### **Primary outcome measure**

1. Uptake of 18FDG-PET-CT in the heart in patients with known cardiac scleroderma is assessed by PET-CT scanning at baseline and 6-9 months
2. Degree and pattern of uptake with echocardiography, cardiac markers, cardiac MRI
3. Prognosis in patients with "active" disease on 18FDG-PET-CT to those with "inactive" disease

### **Secondary outcome measures**

1. Degree and pattern of uptake with echocardiography, cardiac markers and cardiac biopsy findings

2. Level of uptake of 18FDG-PET in other organs known to be involved by scleroderma
3. Change in cardiac 18FDG-PET-CT in patients with “active” disease after completion of therapy
4. 18FDG-PET uptake in the lungs with dedicated pulmonary CT imaging (where available)

**Overall study start date**

01/03/2015

**Completion date**

01/12/2020

## Eligibility

**Key inclusion criteria**

1. Patients with Scleroderma
2. Age 18 or older
3. Satisfied the ACR diagnostic criteria for scleroderma (also known as systemic sclerosis)
4. Suspicion of cardiac involvement from scleroderma with one of:
  - 4.1. Left ventricular ejection fraction < 40% on echocardiogram
  - 4.2. Abnormal heart muscle on cardiac MRI suggestive of scleroderma involvement
  - 4.3. Diastolic heart failure in a patient without known risk factors (must satisfy: age <50, no known systemic hypertension, no diabetes mellitus, pulmonary arterial wedge pressure >15 mmHg on right heart catheterisation at rest)
  - 4.4. Persistently raised troponin (> 2 times upper limit of normal on two occasions at least 28 days apart) without coronary artery disease
  - 4.5. Ventricular arrhythmias
5. No coronary artery disease. This needs to be excluded by CT
6. No coronary artery disease (needs to be excluded by CT coronary angiogram, myocardial perfusion scanning, dobutamine stress echocardiogram or conventional coronary angiography)
7. Able to give informed consent

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

20

**Key exclusion criteria**

1. Inability to lie flat
2. Pregnancy or breastfeeding
3. Unwilling to undergo pregnancy test prior to study (in women of child bearing potential)
4. Coronary artery disease. This needs to be excluded by CT coronary angiogram, myocardial

perfusion scanning, dobutamine stress echocardiogram or conventional coronary angiogram  
5. Inclusion in a clinical trial involving an investigational medical product in the last 28 days

**Date of first enrolment**

01/08/2016

**Date of final enrolment**

01/07/2020

## **Locations**

**Countries of recruitment**

England

United Kingdom

**Study participating centre**

**Royal Free Hospital**

Clinical Trials

Rheumatology Department

Lower Ground Floor

London

United Kingdom

NW3 2QG

## **Sponsor information**

**Organisation**

Royal Free London NHS Foundation Trust

**Sponsor details**

Research and Development Office

Medical School Building

Pond Street

London

England

United Kingdom

NW3 2QG

+44 20 7317 7558 ext. 33211

rf.randd@nhs.net

**Sponsor type**

Hospital/treatment centre

**ROR**

## Funder(s)

### Funder type

Charity

### Funder Name

Royal Free Charity

## Results and Publications

### Publication and dissemination plan

The results will be disseminated to high impact rheumatology and or cardiology meetings and a paper will be submitted to a high impact cardiology and/or rheumatology journal.

### Intention to publish date

31/12/2022

### Individual participant data (IPD) sharing plan

Data generated from the study will be analysed and stored by Dr Benjamin Schreiber for up to 10 years as it may be important for future or follow up extension data.

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No