

Neck movement and eye function

Submission date 23/06/2016	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 23/06/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 20/11/2017	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Neck pain is a very common condition, which can be cause great discomfort and restriction of movement. Patients with neck pain often have visual problems however the reason for this is unknown. One possibility is that the amount of neck movement a person undertakes can influence the eye function. Many patients with neck pain may move their necks less to avoid pain, which may be causing the visual problems. The aim of this study is to look at whether a lot of movement or restricted movement of the neck has an effect on eye function in healthy people with no neck pain, to try to find the out if there is link between neck movement and visual problems.

Who can participate?

Adults with no neck pain who are able to move their necks freely and have normal or corrected-to-normal vision

What does the study involve?

Participants are randomly allocated to undertake two tests spaced one week apart in a different order. Before and 20 minutes after each test, participants have their eye function (measuring of the eye stabilization reflexes) measured. One test involves moving the neck around extensively for 20 minutes. The other test involves wearing a stiff neck collar that restricts movement for one hour.

The measurement of eye function will be in a darkened room and lasts around 20 minutes. It involves having to look straight forward while the chair they are sitting on is rotating, while the position of their eyes is measured with an infrared eye tracking device. Two weeks later, 11 participants return to repeat the condition with the neck collar, after wearing it for two hours. The eye function tests are repeated before and 20 minutes after wearing the collar.

What are the possible benefits and risks of participating?

There are no direct benefits for participants taking part in this study.

Where is the study run from?

Erasmus MC (Netherlands)

When is the study starting and how long is it expected to run for?

August 2014 to May 2015

Who is funding the study?
Erasmus MC (Netherlands)

Who is the main contact?
Miss Britta Ischebeck
bischebeck@sjcn.nl

Contact information

Type(s)
Scientific

Contact name
Mrs Britta Ischebeck

ORCID ID
<https://orcid.org/0000-0003-0801-5242>

Contact details
Department of Neuroscience
Erasmus MC
's-Gravendijkwal 230
Rotterdam
Netherlands
3015 CE
+31 (0)10 4642211
bischebeck@sjcn.nl

Additional identifiers

Protocol serial number
2016_6

Study information

Scientific Title
The influence of cervical movement on eye stabilization reflexes: a randomized cross-over trial

Study objectives
The aim of this study is to investigate the influence of the amount of cervical movement on the cervico-ocular reflex (COR) and vestibulo-ocular reflex (VOR) in healthy individuals.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Ethical board of the Erasmus MC, 19/07/2011, ref: MEC-2011-273

Study design

Single-centre randomised cross-over trial

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Neck pain

Interventions

In the main experiment, two types of intervention are applied to all participants in a cross-over design: hypokinesia and hyperkinesia. Directly before and after the intervention, the eye stabilization reflexes and the active range of motion are measured.

In the hypokinesia intervention, the neck is immobilized by using a stiff neck collar (size 4, Laerdal Stifneck® Select™) for one hour.

In the hyperkinesia intervention, active neck movement in all possible directions of movement is evoked by having the participants move their neck excessively in all directions for twenty minutes.

Each participant receives both interventions on two different days separated by 6 or 7 days. The order of the two interventions was pseudo-randomized and balanced across participants.

In the replication experiment, eleven participants wear the neck collar for two hours (prolonged hypokinesia). This experiment takes place two weeks after the end of the main experiment.

Intervention Type

Behavioural

Primary outcome(s)

Amplitude of the cervico-ocular reflex (COR) and vestibulo-ocular reflex (VOR) is measured with a rotational chair with infrared eye tracking device pre and post each intervention.

Key secondary outcome(s)

The amount of the active cervical range of motion (CROM) in both the horizontal and vertical plane was also measured device (Performance Attainments Associates, USA) pre and post each intervention.

Completion date

01/05/2015

Eligibility

Key inclusion criteria

1. 18 years or older
2. No neck pain
3. Able to move their neck

4. Able to sit on a chair for 20 minutes
5. Normal or corrected-to-normal visual acuity

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Neck pain
2. History of neck trauma
3. Use of tranquilizing medication

Date of first enrolment

01/12/2014

Date of final enrolment

01/05/2015

Locations**Countries of recruitment**

Netherlands

Study participating centre

Erasmus MC

Department of Neuroscience

's-Gravendijkwal 230

Rotterdam

Netherlands

3015 CE

Sponsor information**Organisation**

Erasmus MC

ROR

<https://ror.org/018906e22>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Erasmus Medisch Centrum

Alternative Name(s)

Erasmus Medical Center, Erasmus MC, Erasmus Universitair Medisch Centrum, Erasmus University Medical Center, Universitair Medisch Centrum Rotterdam, Erasmus Universitair Medisch Centrum Rotterdam, EMC

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

Netherlands

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2018		Yes	No