

# Ovulation induction in women with newly diagnosed polycystic ovary syndrome: a randomised double blind clinical trial comparing clomiphene citrate plus metformin with clomiphene citrate plus placebo

<b>Submission date</b>	<b>Recruitment status</b>	<input type="checkbox"/> Prospectively registered
27/01/2006	No longer recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
27/01/2006	Completed	<input checked="" type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
04/04/2013	Nutritional, Metabolic, Endocrine	

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr Etelka Moll

### Contact details

Academic Medical Center  
Department of Obstetrics & Gynecology  
P.O. Box 22660  
Amsterdam  
Netherlands  
1100 DD  
E.Moll@amc.uva.nl

## Additional identifiers

### Protocol serial number

NTR485

# Study information

## Scientific Title

### Acronym

The Metformin trial

## Study objectives

To compare the effectiveness of clomiphene citrate with metformin versus clomiphene citrate only in women with newly diagnosed polycystic ovary syndrome with respect to ovulation, pregnancy and spontaneous abortions.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Polycystic Ovary Syndrome (PCOS)

## Interventions

1. Metformin 4 dd 500 mg plus clomiphene citrate 50 - 150 mg
2. Control: placebo 4 dd plus clomiphene citrate 50 - 150 mg

Patients used the medication as long as they were participating in the study. They stopped taking medication when they were clomiphene citrate resistant, pregnant or dropped out for a different reason.

## Intervention Type

Drug

## Phase

Not Specified

## Drug/device/biological/vaccine name(s)

Clomiphene citrate, metformin

## Primary outcome(s)

Ovulation.

**Key secondary outcome(s)**

1. Ongoing pregnancy
2. Spontaneous abortion
3. Clomiphene citrate resistance

**Completion date**

01/06/2004

## Eligibility

**Key inclusion criteria**

All patients with chronic anovulation World Health Organization (WHO) type II, polycystic ovaries diagnosed by transvaginal ultrasonography and child wish.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

Female

**Key exclusion criteria**

1. Other causes of anovulation
2. Age over 40 years and liver, kidney or heart disease/failure (i.e. abnormal results on liver function tests)
3. Serum creatinine concentration greater than 95 umol/l or a history of heart disease/failure
4. Sperm quality indicating male factor subfertility (total motile count less than  $10 \times 10^6$ )

**Date of first enrolment**

01/06/2001

**Date of final enrolment**

01/06/2004

## Locations

**Countries of recruitment**

Netherlands

**Study participating centre**

## Academic Medical Center

Amsterdam  
Netherlands  
1100 DD

## Sponsor information

### Organisation

Academic Medical Center (AMC) (The Netherlands)

### ROR

<https://ror.org/03t4gr691>

## Funder(s)

### Funder type

Industry

### Funder Name

Merck B.V. (The Netherlands)

## Results and Publications

### Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	24/06/2006		Yes	No
<a href="#">Results article</a>	results	01/11/2012		Yes	No