PREVEC Trial: Prevention of Reperfusion damage associated with percutaneous coronary angioplasty following acute myocardial infarction

Submission date 21/01/2013	Recruitment status No longer recruiting	[X] Prospectively registered	
21/01/2013		[X] Protocol	
Registration date	Overall study status Completed	Statistical analysis plan	
28/01/2013		Results	
Last Edited	Condition category	Individual participant data	
03/06/2014	Circulatory System	Record updated in last year	

Plain English summary of protocol

Background and study aims

Acute myocardial infarction (AMI) is the leading cause of mortality in Chile and worldwide. It also has negative effects on quality of life in survivors.

Existing treatments have reduced mortality. But they have the side effect of restoring the blood flow to the heart which is called an ischemia-reperfusion event. Ischemia-reperfusion increases heart damage through a process called oxidative stress. Ischemia reperfusion damage may be preventable. The aim of the study is to assess whether increasing the antioxidant defences will decrease heart damage. This will be done by giving antioxidant vitamin C (intravenously) and vitamin E (orally) to acute myocardial infarction patients during the standard procedure for AMI called percutaneous coronary angioplasty.

Who can participate?

Patients of either sex, over 18 years old, with an indication of primary percutaneous coronary angioplasty and experiencing their first acute myocardial infarction (symptoms must have started during the last 12 hours).

What does the study involve?

Patients will be randomly allocated to one of two groups. One group (Vitamin-treatment group) will receive the standard procedure for an acute myocardial infarction plus a high-dose infusion of vitamin C and an oral dose of vitamin E.

The other group (Control Group) will receive the standard procedure for an acute myocardial infarction plus an infusion of a harmless substance (saline solution) instead of vitamin C and a harmless oral dose of vegetal oil.

Patients in both groups will give blood at entry, during the intervention procedure and at discharge. Cardiac magnetic resonance assessment will be performed 6 and 84 days after the procedure.

What are the possible benefits and risks of participating?

For the Vitamin-treatment group, it is possible that the infarct size in the heart is reduced in comparison to the Control group. Benefits for the Control group are a longer follow-up by the medical team. Possible reported side effects in the Vitamin-treatment group are lethargy or fatigue (0.17% of patients) and/or phlebitis (0.05% of patients).

Where is the study run from?

The University of Chile Clinical Hospital, Cardiovascular Department (Chile).

When is the study starting and how long is it expected to run for? The study will start in February 2013 and will run for 3 years or until the required number of 132 patients have been treated and evaluated.

Who is funding the study?

The Scientific and Technological Development Fund (FONDECYT) - Chilean Government

Who is the main contact? Dr Ramon Rodrigo rrodrigo@med.uchile.cl

Contact information

Type(s)

Scientific

Contact name

Dr Ramón Rodrigo

Contact details

Independencia 1027 Santiago Chile 8380453

Additional identifiers

Protocol serial number

1120594

Study information

Scientific Title

PREVEC Trial: Prevention of Reperfusion damage associated with percutaneous coronary angioplasty following acute myocardial infarction: a randomized double-blind placebocontrolled clinical trial

Acronym

PREVEC

Study objectives

Patients subjected to percutaneous coronary angioplasty to restore the coronary blood flow previously impaired by an acute myocardial infarction, while receiving a short-term infusion of high doses of vitamin C, plus oral doses of the recommended dose of vitamin E, will present a lower infarct size, as well as an attenuation of the functional and biochemical damage occurring during the reperfusion following to the sudden loss of blood supply, as compared with placebotreated patients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Local Ethics Committees (University of Chile Clinical Hospital, Faculty of Medicine of the University of Chile and Health Ministry of the Chilean Government), 07/2011, ref: Project number - 06-2011

Study design

Randomized double-blind placebo-controlled clinical trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Acute Myocardial Infarction

Interventions

Intervention:

Short term massive infusion of intravenous vitamin C (320 mmol/L)

Oral dose of vitamin E (400 IU/day) throughout the protocol

Oral dose of vitamin C (500 mg/12 hours) following angioplasty, throughout the protocol

Control: Placebo

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Infarct size, assessed by cardiac magnetic resonance (CMR) will be measured twice: 6 and 84 days following coronary angioplasty

Key secondary outcome(s))

Biomarkers for oxidative stress, antioxidant status, heart damage and inflammation will be measured at baseline, at the onset of reperfusion, 6-8 hours after revascularization and at hospital discharge.

Completion date

Eligibility

Key inclusion criteria

- 1. Subjects may be of either sex and must be at least 18 years old
- 2. Subjects must have indication of primary percutaneous coronary angioplasty (PCA) as follows:
- 2.1. Angina or equivalent at least 120 min duration
- 2.2. Electrocardiogram (ECG) with STEMI that concerns more than 2 contiguous leads (>2mm)
- 3. Presentation within 12 h of symptoms onset
- 4. First myocardial infarction
- 5. Subject must be able and willing to sign informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. History of renal or hepatic insufficiency
- 2. History of renal lithiasis (oxalates)
- 3. History of heart failure (NYHA III, IV)
- 4. Cardiogenic shock
- 5. Any serious medical co-morbidity that determine life expectancy < 6 month
- 6. Current participation in any other clinical investigation
- 7. Pregnancy
- 8. Glucose 6-phosphate dehydrogenase deficiency

Date of first enrolment

04/02/2013

Date of final enrolment

31/03/2016

Locations

Countries of recruitment

Chile

Study participating centre Independencia 1027 Santiago Chile 8380453

Sponsor information

Organisation

Scientific and Technological Development Fund (FONDECYT Fondo de Desarrollo Científico y Tecnológico) (Chile)

ROR

https://ror.org/02ap3w078

Funder(s)

Funder type

Government

Funder Name

Scientific and Technological Development Fund (FONDECYT) (Chile) grant number 1120594.

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created Date added	l Peer reviewed?	Patient-facing?
Protocol article	protocol	29/05/2014	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	5 No	Yes