

Pre-operative short-course radiotherapy versus neoadjuvant radiochemotherapy in locally advanced rectal cancer (uT2N+, uT3N-/+)

Submission date 25/05/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 02/07/2007	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 04/01/2021	Condition category Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Peter M Schlag

Contact details
Department of Surgery and Surgical Oncology
Charité-Universitätsmedizin Berlin
Campus Berlin-Buch
Lindenberger Weg 80
Berlin
Germany
D-13125
-
pmschlag@charite.de

Additional identifiers

Clinical Trials Information System (CTIS)
2004-001606-27

Protocol serial number
N/A

Study information

Scientific Title

Pre-operative short-course radiotherapy versus neoadjuvant radiochemotherapy in locally advanced rectal cancer (uT2N+, uT3N-/+)

Acronym

BRCT (Berlin Rectal Cancer Trial)

Study objectives

Standard treatment for locally advanced cancer of the rectum is pre-operative short-course radiotherapy or combined neoadjuvant radiochemotherapy with 5-fluorouracil (5-FU) plus post-operative chemotherapy with 5-FU. Similar long-term survival, local control and late morbidity have been reported for both these methods in non-comparative studies. In addition to other ongoing comparative trials we include a larger number of patients for adequate power and we avoid the adjuvant treatment bias by mandatory adjuvant chemotherapy in both groups. It is our hypothesis that the rate of local recurrence after five years is 12% in pre-operative short-course radiotherapy and 7% in combined radiochemotherapy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved by the Ethics Committee of the Charité and the responsible authorities. Certified and recommended by the German Cancer Society ("Gütesiegel A") on the 29th September 2003 (ref: AA3/03/38; EudraCT-number: 2004-001606-27).

Study design

Randomised controlled multicentre trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Rectal cancer

Interventions

Group one: receiving pre-operative short-course radiotherapy (five times 5 Gy) followed by total mesorectal excision (TME) and adjuvant continuous 5-FU infusion therapy for 12 weeks.

Group two: receiving neoadjuvant combined radiochemotherapy (50.4 Gy and continuous 5-FU) followed by TME and adjuvant continuous 5-FU infusion therapy for 12 weeks.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

5-fluorouracil (5-FU)

Primary outcome(s)

Local recurrence, median follow up five years

Key secondary outcome(s)

1. Overall survival, median follow up five years
2. Disease-free survival, median follow up five years
3. Complete resection rate (R0 resection): measured at the date of surgery of the last patient entered
4. Rate of sphincter saving resection: measured at the date of surgery of the last patient entered
5. Acute and late toxicity (radiation related side effects), median follow up five years
6. Quality of life including long term bowel function, median follow up five years

Completion date

31/12/2009

Eligibility**Key inclusion criteria**

1. Aged 18 years or over
2. Karnofsky Index 80% or better
3. Histological diagnosis of adeno- or mucinous carcinoma of rectum
4. Primary rectal cancer:
 - 4.1. Maximum 12 cm above dentate line (upper limit)
 - 4.2. Staged T2N+ or T3N0 or T3N+ (by endorectal ultrasound or computed tomography [CT] /magnetic resonance imaging [MRI] scan)
5. No evidence of metastatic disease as determined by chest X-ray and abdominal ultrasound (or CT-scan of chest and abdomen or other investigations such as positron emission tomography [PET] scan or biopsy if required)
6. Adequate bone marrow function with platelets more than $100 \times 10^9/l$ and neutrophils more than $2.0 \times 10^9/l$
8. Creatinine clearance more than 50 ml/min
7. Serum bilirubin less than 2.0 x upper limit of institutional normal range (ULN)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Not Specified

Key exclusion criteria

1. Rectal cancer other than adeno- or mucinous carcinoma
2. Previous or concurrent malignancies, with the exception of adequately treated basal cell carcinoma of the skin or in situ carcinoma of the cervix
3. Patients with locally advanced inoperable disease, such as T4-tumour
4. Presence of metastatic disease or recurrent rectal tumour
5. Any previous chemotherapy or radiotherapy, and any investigational treatment for rectal cancer
6. Concurrent uncontrolled medical conditions
7. Pregnancy or breast feeding
8. Clinically significant (i.e., active) cardiac disease (e.g., congestive heart failure, symptomatic coronary artery) or myocardial infarction within the last six months
9. Stenotic tumour which can not be passed by the colonoscope and pre-operative need of diverting stoma
10. Evidence of hereditary colorectal cancer (hereditary non-polyposis colorectal cancer [HNPCC] and familial adenomatous polyposis [FAP])
11. Medical or psychiatric conditions that compromise the patients ability to give informed consent

Date of first enrolment

15/01/2004

Date of final enrolment

31/12/2009

Locations

Countries of recruitment

Austria

Germany

Study participating centre

Department of Surgery and Surgical Oncology

Berlin

Germany

D-13125

Sponsor information

Organisation

Berlin Cancer Society (Berliner Krebsgesellschaft e.V.) (Germany)

ROR

<https://ror.org/020yxp837>

Funder(s)

Funder type

Research organisation

Funder Name

Berlin Cancer Society (Berliner Krebsgesellschaft e.V.) (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	06/02/2009	04/01/2021	Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes