# The affect of diode laser on healing after dental extraction in patients diabetes mellitus

Submission date	Recruitment status	Prospectively registered
14/02/2025	No longer recruiting	<pre>Protocol</pre>
Registration date	Overall study status	Statistical analysis plan
20/02/2025	Completed	Results
Last Edited	Condition category	Individual participant data
20/02/2025	Oral Health	[X] Record updated in last year

### Plain English summary of protocol

Background and study aims

Patients visiting dental clinics and those suffering from type 2 diabetes often experience vascular fragility, subsequent bleeding, delayed healing, and the need for various diluents. Therefore, it is essential to develop special materials for these patients that reduce bleeding, accelerate healing, and minimize infections, making extraction safe, easy, and comfortable for both the doctor and the patient. One of these materials is a diode laser, which has numerous positive effects on the body, including pain relief, reduced bleeding, and the promotion of growth factors, thereby enhancing healing.

### Who can participate?

Patients aged 35 years and older with Type II Diabetes require symmetrical tooth extraction.

### What does the study involve?

The control group will receive a placebo laser after the tooth while the experimental group will receive a diode laser after tooth extraction. The participants will be followed up for 7 days.

### What are the possible benefits and risks of participating?

The potential benefits of using lasers for diabetics include pain relief, bleeding, and expedited healing. However, the potential risks involve prolonged working time and Contraindications in cases of local infection.

# Where is the study run from?

Oral and Maxillofacial Surgery Hospital, Faculty of Dentistry, Damascus University (Syria)

When is the study starting and how long is it expected to run for? November 2022 to January 2025

Who is the funding of the study? Investigator initiated and funded

Who is the main contact?
Dr Mhd Obada Shakkour, oshakkour95@gmail.com

# Contact information

### Type(s)

Public, Scientific, Principal investigator

#### Contact name

Dr Mhd Obada Shakkour

#### **ORCID ID**

https://orcid.org/0009-0004-9692-2276

### Contact details

Mezzeh Highway Damascus Syria 0000 +963 999041252 obada95.shakkour@damascusuniversity.edu.sy

# Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

The affect of diode laser on healing after dental extraction in patients type 2 diabetes mellitus

# Study objectives

The null hypothesis posited that there is a significant difference between a diode laser and a placebo laser in enhancing healing after tooth extractions in individuals with type 2 diabetes mellitus.

### Ethics approval required

Ethics approval required

### Ethics approval(s)

approved 26/11/2024, Damascus University (Oral and Maxillofacial Surgery Department - Faculty of Dental Medicine) (Mezzeh Highway, Damascus, -, Syria; +963 1133923192; ap. srd@damascusuniversity.edu.sy), ref: DN-261124-351

# Study design

Double-blind randomized controlled clinical trial

### Primary study design

Interventional

### Study type(s)

Quality of life, Treatment, Efficacy

### Health condition(s) or problem(s) studied

Improving healing after dental extraction in individuals with diabetes

#### **Interventions**

This study is a randomized controlled trial, involving the application of a diode laser and a placebo laser. The control group will receive a placebo laser after the tooth while the experimental group will receive a diode laser after tooth extraction. The diode and placebo lasers will be randomized into two groups using http://www.randomizer.org/. The follow-up period for each patient is one week, with visits:

1) 2 days postoperative and 2) 7 days postoperative.

### Intervention Type

Procedure/Surgery

### Primary outcome(s)

Wound healing will be measured at three endpoints: T0 immediately after laser application, T1 at 2 days, and T2 at 7 days by Landry Gingival healing index

# Key secondary outcome(s))

Pain will be assessed using a visual analog scale and the painkiller pills at T0 immediately after laser application, T1 at 2 days, and T2 at 7 days

### Completion date

15/01/2025

# Eligibility

# Key inclusion criteria

- 1. Patients with Type 2 diabetes and HbA1c levels between 8-10%.
- 2. Individuals over 35 years of age.
- 3. Patients classified as ASA type II III.

# Participant type(s)

**Patient** 

# Healthy volunteers allowed

No

# Age group

Adult

# Lower age limit

### Upper age limit

70 years

### Sex

All

### Total final enrolment

20

### Key exclusion criteria

- 1. HbA1c less than 8 or greater than 10.
- 2. Antibiotic or NSAID use1- days before extraction.
- 3. Smoking more than 10 cigarettes a day.
- 4. Local infection at the site of the tooth to be extracted.

### Date of first enrolment

12/01/2024

### Date of final enrolment

05/01/2025

# Locations

# Countries of recruitment

Syria

# Study participating centre

Oral and Maxillofacial Surgery Hospital, Faculty of Dentistry, Damascus University

Mezzeh Highway Damascus

Syria

4671

# Sponsor information

### Organisation

**Damascus University** 

### **ROR**

https://ror.org/03m098d13

# Funder(s)

### Funder type

Other

### **Funder Name**

Investigator initiated and funded

# **Results and Publications**

### Individual participant data (IPD) sharing plan

the data sharing plans for the current study are unknown and will be made available at a later date

# IPD sharing plan summary

Data sharing statement to be made available at a later date

# **Study outputs**

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 11/11/2025 No Yes