# Promoting mental health for university students in China through mindfulness: A randomized controlled trial across six universities

<b>Submission date</b> 05/09/2025	Recruitment status  No longer recruiting	[X] Prospectively registered
		∐ Protocol
<b>Registration date</b> 09/09/2025	Overall study status Ongoing	Statistical analysis plan
		☐ Results
<b>Last Edited</b> 24/09/2025	Condition category  Mental and Behavioural Disorders	Individual participant data
		[X] Record updated in last year

#### Plain English summary of protocol

Background and study aims

Mindfulness-based programs (MBPs), rooted in contemplative practices, have emerged as promising interventions for promoting mental well-being and resilience. Unfortunately, no large-scale study has assessed the efficacy of MBPs for university students in China, a population with a high prevalence of mental health issues. Building off of a pilot study implemented in spring 2025, the primary goal of this larger-scale study is to evaluate the impact of a mindfulness-based program specifically tailored for university students. We will investigate the effects on mental illness symptoms, stress (perceived and physiological), state and trait mindfulness, emotion regulation skills, and other indicators of well-being, such as sleep quality and somatic symptoms. We will also conduct ecological momentary assessments for one week (both pre- and post-intervention), four times per day, to explore the effects on direct, immediate experience.

#### Who can participate?

Undergraduate students at participating universities with mild to severe symptoms of anxiety and/or depression (see inclusion and exclusion criteria above).

#### What does the study involve?

Students in the intervention group will participate in an eight-week mindfulness program (in-person).

What are the possible benefits and risk of participating?

Potential benefits are improved mental and physical health. Students will also receive small financial rewards for participating in the study.

Potential risks are unlikely, but may include increased awareness of mental or physical discomfort, as well as time stress due to extra time dedicated to participation.

#### Where is the study run from?

Overall, students from six universities will participate in the study. The two main China-based universities (Fudan University and Beijing Normal University) will oversee the study implementation, facilitating participant recruitment and data collection at five other universities in China. Researchers at Stanford University will work with these partners to implement the project.

When is the study starting and how long is it expected to run for? The study will begin recruitment for the mid-September 2025. The intervention is planned to begin in mid-October and last for nine weeks.

#### Who is funding the study?

Study funders include the Cyrus Tang Foundation, the Enlight Foundation, and Vincent Woo Foundation.

#### Who is the main contact?

Researchers may contact Dr. Huan Wang (huanw@stanford.edu), Dr. Hui-Qi Tong (htong@stanford.edu), or Cody Abbey (cjabbey@stanford.edu) for more information about this study.

#### Contact information

#### Type(s)

Public, Scientific

#### Contact name

Mr Cody Abbey

#### **ORCID ID**

https://orcid.org/0000-0002-1281-5472

#### Contact details

616 Jane Stanford Way, Stanford University Stanford United States of America 94305 +1 8282803225 cjabbey@stanford.edu

#### Type(s)

Principal investigator

#### Contact name

Dr Huan Wang

#### Contact details

616 Jane Stanford Way, Stanford University Stanford United States of America 94305 +1 650-724-9254 huanw@stanford.edu

#### Type(s)

Principal investigator

#### Contact name

Dr Hui Qi Tong

#### Contact details

91 Campus Drive Li Ka Shing Building Stanford United States of America 94305 +1 650-724-9254 htong@stanford.edu

#### Type(s)

Principal investigator

#### Contact name

Dr Xinshu She

#### Contact details

91 Campus Drive Li Ka Shing Building Stanford United States of America 94305 +1 650-724-9254 xinshe@stanford.edu

#### Type(s)

Principal investigator

#### Contact name

Prof Scott Rozelle

#### Contact details

616 Jane Stanford Way, Stanford University Stanford United States of America 94305 +1 650-724-9254 rozelle@stanford.edu

#### Type(s)

Principal investigator

#### Contact name

**Prof Lian Tong** 

#### Contact details

220 Handan Rd, Yangpu District Shanghai China 200437 +86 21 5423 7499 ltong@fudan.edu.cn

#### Type(s)

Principal investigator

#### Contact name

Prof Danhua Lin

#### Contact details

19 Xinwai Ave, Beitaipingzhuang, Hai Dian Qu Beijing China 100875 +86 010 58804075 danhualin@bnu.edu.cn

# Additional identifiers

#### Clinical Trials Information System (CTIS)

Nil known

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

AEARCTR-0016685

# Study information

#### Scientific Title

Promoting mental health for university students in China through mindfulness: A randomized controlled trial across six universities

#### **Study objectives**

Participation in the mindfulness-based program will reduce anxiety and depression symptoms more than an experimental waitlist control

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

approved 18/12/2024, Stanford University Institutional Review Board (1705 El Camino Real Palo Alto, Stanford, 94305, United States of America; +1 (650) 724-7141; kateri.noble@stanford.edu), ref: 75117

#### Study design

Interventional randomized controlled trial

#### Primary study design

Interventional

#### Study type(s)

Prevention, Treatment, Efficacy

#### Health condition(s) or problem(s) studied

Prevention of chronic mental disorders in students with depression and/or anxiety symptoms

#### **Interventions**

We will conduct a randomized study with 360 undergraduate students recruited from six universities in China. Using the RCT method of impact evaluation, we will be able to ensure that the intervention and control groups have similar characteristics at baseline, which is vital for drawing a causal connection between the intervention and any changes in student outcomes. By assuring the similarity of characteristics (such as sex, baseline mental health, etc.) between the treatment and waitlist control groups at baseline, we can confidently attribute any significant differences in outcomes between the control and intervention group to the program.

For this study, the 360 sample students will be randomly allocated into two experimental arms (in-person mindfulness-based program and a control). Using STATA 16 software (https://www.stata.com/), with 80% power, a significance level of 0.05, we determined that a sample size of 139 students per arm was required to detect a difference of 0.30 standard deviations in outcome measures. Our recruitment of 180 students per arm (360 in total) allows for 20% attrition.

#### In-person mindfulness instruction:

- The in-person mindfulness instruction will consist of weekly 90-minute group sessions, held in a university classroom over eight consecutive weeks, not including a introductory session prior to the formal start of classes.
- Participants will learn foundational principles related to the role of mindfulness in specific domains such as responding to stress, mindful communication in relationships, and gratitude.
- Participants will also be taught formal mindfulness practices (e.g., sitting meditation, body scan, qigong, loving-kindness meditation) during the weekly sessions and informal mindfulness practices (e.g., mindful eating, mindful walking, and mindful daily routine activities) during the weekly sessions.

They will be asked to commit to completing assigned mindfulness practice independently throughout the intervention period.

Participants assigned to the control group will receive no other intervention during the program (pure control).

All study participants will receive adequate financial compensation for participating in the study.

#### Intervention Type

Behavioural

#### Primary outcome(s)

1. Anxiety symptoms will be measured using the General Anxiety Disorder Scale (GAD-7) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention (post-graduation from undergraduate studies) 2. Depression symptoms will be measured using the Patient Health Questionnaire Scale (PHQ-9) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention (post-graduation from undergraduate studies)

#### Key secondary outcome(s))

Current secondary outcome measures as of 24/09/2025:

- 1. Physiological stress will be measured using cortisol levels. The cortisol levels will be measured via saliva samples starting one week before the intervention, three times on the same day at consistent times. At the end of the intervention, the cortisol levels will be measured one week after the course ends, three times on the same day (at the same times as before the intervention).
- 2. Perceived stress is measured using the Perceived Stress Scale (PSS), measured at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 3. Emotion regulation will be measured using the Emotion Regulation Questionnaire (ERQ) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 4. Coping strategies will be measured using the Coping Strategies Scale at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 5. Trait mindfulness will be measured using the short-form of the Five Facet Mindfulness Questionnaire (FFMQ-SF) at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 6. State mindfulness will be measured using the Multidimensional State Mindfulness Questionnaire (MSMQ) baseline, endline, measured weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 7. Loneliness will be measured using the short-form of the 6-item UCLA Loneliness Scale at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 8. Recent stressful events will be measured using the Adolescent Self-Rating Life Events Checklist (ASLEC) at baseline and endline, and in several follow-ups at three months, six months, and four years after the intervention
- 9. Sleep quality will be measured using the Pittsburgh Sleep Quality Index (PSQI) at baseline, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 10. Flourishing will be measured using the Flourishing Scale (FS) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 11. Life satisfaction will be measured using the Satisfaction with Life Scale (SWLS) at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 12. Academic performance will be measured using the semester GPA and national English test results at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention

- 13. Lifestyle behaviors will be measured using items asking about the amount and frequency of behaviors such as screen use and exercise at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 14. Face will be measured using the CPAI-2 at baseline and endline, and in several follow-ups at three months, six months, and four years after the intervention
- 15. Somatic symptoms will be measured using the brief version of the Somatic Symptom Scale (SSS), at baseline and endline, and in several follow-ups at three months, six months, and four years after the intervention
- 16. Perceived benefits and adverse effects and helpfulness of the program will be measured with Likert-type items measured weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 17. Compliance with mindfulness practice will be measured based on self-report length and frequency of practice, measured weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 18. Acceptability of the intervention will be measured via qualitative interviews and focus groups with students and instructors
- 19. Self-compassion will be measured using the Self-Compassion Scale, measured at baseline and endline, as well as weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 20. Sense of purpose will be measured using the Meaning of Life Questionnaire, measured at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 21. Experiential avoidance will be measured using the Brief Experiential Avoidance Questionnaire at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 22. Emotion Reactivity and Instability will be measured using the RIPOST-Y at baseline, endline, weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 23. Career outcomes will be measured using items asking about post-graduation trajectories four years following the intervention
- 24. Interpersonal tolerance will be measured using the CPAI-2 at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 25. Perceived ostracism will be measured using the Ostracism Experience Scale (OES-A), measured at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 26. Fear of negative evaluation will be measured using the Brief Fear of Negative Evaluation Scale (BFNE-II), measured at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 27. Neuroticism will be measured using the Big Five Personality Inventory (brief version), at baseline and endline
- 28. Openness will be measured using the Big Five Personality Inventory (brief version), at baseline and endline
- 29. Emotion reactivity will be measured using the Perth Emotion Reactivity Scale, measured at baseline and endline
- 30. Gratitude will be measured using the five-item Gratitude Questionnaire, measured at baseline, endline, weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 31. Stress mindset will be measured using the Stress Mindset Scale, measured at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention
- 32. Positive and negative emotions will be measured using the PANAS, measured at baseline, endline, weekly during the intervention, and in several follow-ups at three months, six months,

and four years after the intervention

33. Interpersonal tolerance will be measured using the tolerance scale of the CPAI-2, measured at baseline, endline, and in several follow-ups at three months, six months, and four years after the intervention

Ecological momentary assessment scales are conducted for one week (both pre- and post-intervention), four times per day, to explore the effects on direct, immediate experience:

- 1. State mindfulness (MSMQ-9)
- 2. Depression (PHQ-2)
- 3. Anxiety (GAD-2)
- 4. Life satisfaction (SWLS-2)
- 5. Positive and negative emotions (14 emotion words, drawn from the I-PANAS-SF, PANAS-X, and parts of the LAPA)
- 6. Sleep duration and quality (one item each from the PSQI)
- 7. Perceived stress (one self-developed item)
- 8. Emotion regulation efficacy (two self-developed items)
- 9. Negative and positive events (seven self-developed items each)
- 10. Overall daily emotional stability (adapted from the TIPI)

#### Previous secondary outcome measures:

- 1. Stress (physiological) will be measured using cortisol levels. The cortisol levels will be measured via saliva samples starting one week before the intervention, three times on the same day at consistent times. At the end of the intervention, the cortisol levels will be measured one week after the course ends, three times on the same day (at the same times as before the intervention).
- 2. Perceived stress is measured using the Perceived Stress Scale (PSS), measured at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 3. Emotion regulation will be measured using the Emotion Regulation Questionnaire (ERQ) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 4. Coping strategies will be measured using the Coping Strategies Scale at baseline and endline
- 5. Adverse Childhood Experiences will be measured using the Adverse Childhood Experiences International Questionnaire (ACES-IQ) and Childhood Trauma Questionnaire short form (CTQ-SF) at baseline
- 6. Trait mindfulness will be measured using the short-form of the Five Facet Mindfulness Questionnaire (FFMQ-SF) at baseline and endline
- 7. State mindfulness will be measured using the Multidimensional State Mindfulness Questionnaire (MSMQ) at baseline and endline, and measured weekly during the intervention 8. Loneliness will be measured using the short-form of the 6-item UCLA Loneliness Scale at baseline and endline
- 9. Attachment style will be measured using the Revised Adult Attachment Scale at baseline and endline
- 10. Recent stress will be measured using the Adolescent Self-Rating Life Events Checklist (ASLEC) at baseline and endline, and in several follow-ups at three months, six months, and four years after the intervention
- 11. Sleep quality will be measured using the Pittsburgh Sleep Quality Index (PSQI) at baseline, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 12. Flourishing will be measured using the Flourishing Scale (FS) at baseline, weekly during the intervention, at endline, and in several follow-ups at three months, six months, and four years after the intervention

- 13. Life satisfaction will be measured using the Satisfaction with Life Scale (SWLS)
- 14. Academic performance will be measured using the semester GPA and national English test results at baseline and endline, at endline, and in several follow-ups at three months, six months, and four years after the intervention
- 15. Lifestyle behaviors will be measured using items asking about the amount and frequency of behaviors such as screen use and exercise at baseline and endline
- 16. Neuroticism will be measured using the Big Five Personality Inventory (brief version), at baseline and endline
- 17. Openness will be measured using the Big Five Personality Inventory (brief version), at baseline and endline
- 18. Face will be measured using the CPAI-2 at baseline and endline, and in several follow-ups at three months, six months, and four years after the intervention
- 19. Somatic symptoms will be measured using the brief version of the Somatic Symptom Scale (SSS), at baseline and endline
- 20. Perceived benefits and adverse effects and helpfulness of the program will be measured with Likert-type items measured weekly during the intervention
- 21. Compliance with mindfulness practice will be measured based on self-report length and frequency of practice, measured weekly during the intervention.
- 22. Acceptability of the intervention will be measured via qualitative interviews and focus groups with students and instructors
- 23. Self-compassion will be measured using the Self-Compassion Scale, measured at baseline and endline, as well as weekly during the intervention
- 24. Sense of purpose will be measured using the Meaning of Life Questionnaire, measured at baseline and endline
- 25. Experiential avoidance will be measured using the Brief Experiential Avoidance Questionnaire at baseline and endline
- 26. Emotion Reactivity and Instability will be measured using the RIPOST-Y at baseline, endline, weekly during the intervention, and in several follow-ups at three months, six months, and four years after the intervention
- 27. Career outcomes will be measured using items asking about post-graduation trajectories four years following the intervention
- 28. Interpersonal tolerance will be measured using the CPAI-2 at baseline and endline
- 29. Harmony will be measured using the CPAI-2 at baseline and endline
- 30. Self vs. social orientation will be measured using the CPAI-2 at baseline and endline

#### Completion date

30/01/2030

# **Eligibility**

#### Key inclusion criteria

- 1. Undergraduate student enrolled at participating universities in China
- 2. Reporting mild, moderate, or severe symptoms of depression and/or anxiety (PHQ-9>=5; GAD-7 >= 5)
- 3. Willing to attend weekly classes and engage in mindfulness practice at home
- 4. 18 years or older

#### Participant type(s)

Learner/student

#### Healthy volunteers allowed

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

- 1. In graduating year of undergraduate program
- 2. Has moderate/severe risk of self-harm or harm to others (thoughts of self-harm and/or harm to others occurring several or more days per week)
- 3. Has received formal, teacher-led mindfulness meditation training before and/or has a regular meditation practice (once or more per week over the past two months)
- 4. Is currently receiving psychotherapy and/or taking psychiatric medication

#### Date of first enrolment

15/09/2025

#### Date of final enrolment

07/10/2025

## Locations

#### Countries of recruitment

China

#### Study participating centre Beijing Normal University

19 Xinwai Ave, Beitaipingzhuang, Haidian District Beijing China 100875

# Study participating centre Fudan University

220 Handan Rd, Yang Pu Qu Shanghai China 200437

# Sponsor information

#### Organisation

Stanford University

#### **ROR**

https://ror.org/00f54p054

# Funder(s)

#### Funder type

Charity

#### **Funder Name**

Cyrus Tang Foundation

#### Alternative Name(s)

**CTF** 

#### **Funding Body Type**

Private sector organisation

#### **Funding Body Subtype**

Trusts, charities, foundations (both public and private)

#### Location

United States of America

#### **Funder Name**

**Enlight Foundation** 

#### Alternative Name(s)

#### **Funding Body Type**

Government organisation

#### Funding Body Subtype

Trusts, charities, foundations (both public and private)

#### Location

**United Kingdom** 

# **Results and Publications**

# Individual participant data (IPD) sharing plan

Databases will be shared for replication studies upon request and reasonable conditions. huanw@stanford.edu

## IPD sharing plan summary

Available on request