

A multicenter, randomized, double-blind, placebo-controlled investigation of long-term safety and efficacy of LCAP (leukocytapheresis using "Cellsorba FX") in patients with refractory, chronic active ulcerative colitis

Submission date 28/10/2004	Recruitment status Stopped	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 17/01/2005	Overall study status Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 13/12/2007	Condition category Digestive System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Reinhard Klingel

Contact details
Apheresis Research Institute
Stadtwaldgürtel 77
Cologne
Germany
50935
+49-221-406 317 0
afi@apheresis-research.de

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

1.6 - 07/2004

Study information

Scientific Title

Acronym

MICELL-UC

Study objectives

Study hypothesis added as of 8 June 2007: Assessment of long-term safety and efficacy of leukocytapheresis with Cellsorba FX in comparison to sham-leukocytapheresis for patients with refractory, chronic active ulcerative colitis (CAI 6-10).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval information added as of 8 June 2007:

Approvals of the following ethics committees were obtained on 19 August 2004:

1. Hannover Medical School (MHH)
2. University (LMU) of Munich
3. University of Erlangen
4. Medical association of Mecklenburg-Western Pomerania (Rostock)
5. University of Munster
6. Charité, University of Berlin
7. Medical association of Rhineland-Palatinate (Mainz)

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Refractory, chronic active ulcerative colitis

Interventions

Please note that this study was terminated on 25 April 2007 due to unsatisfactory patient enrolment.

Interventions provided at registration:

Extracorporeal leukocytapheresis (LCAP; verum group) versus sham-leukocytapheresis (placebo group)

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Primary outcome measures added as of 8 June 2007:

The primary efficacy parameter is the 7-item Clinical Activity Index (CAI). A sum score will be calculated summing up all items that are differently weighted. The range of the sum score is 0 to 29 points. A sum score of less than four (or less than or equal to 4) points at the end of the therapy will be assessed as remission or success.

Secondary outcome measures

Secondary outcome measures added as of 8 June 2007:

1. Inflammatory Bowel Disease Questionnaire (IBDQ), German translation. This questionnaire consists of four domains and 32 items. Domains are simply the sum of specific items, i.e. bowel symptoms, systemic symptoms, emotional functions and social functions.
2. Endoscopic Index (EI), which includes 4 items with different weights. The range of the sum score is 0 to 12 points.
3. Cumulative steroid dose over intensive and maintenance phase.
4. CAI, item-wise analysis.

Overall study start date

01/11/2004

Completion date

31/12/2006

Reason abandoned (if study stopped)

The trial will be terminated due to unsatisfactory patient enrolment: during the scheduled period of 24 months, less than 50% of patient enrolment has been completed.

Eligibility**Key inclusion criteria**

1. Ulcerative colitis with chronic active disease state
2. Colitis endoscopically covering at least 15 cm
3. Clinical Activity Index: 6-10
4. No long term remission using combined standard therapy including 5-Aminosalicylate (ASA), prednisolone and/or azathioprine
5. Cumulative steroid dosage within the last 2 months in total at least 600 mg
6. Steroid dosage constant 10 mg per day during 2 weeks before start of treatment (pre-treatment phase)
7. Negative test of pregnancy for female patients
8. Patient is able to understand and sign informed consent

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

Planned number of recruitment added as of 8 June 2007: 95 (less than 50% achieved).

Key exclusion criteria

Exclusion criteria added as of 8 June 2007:

1. Age < 18 or ≥ 80 years
2. Effective response to conventional Ulcerative Colitis (UC) therapy
3. Diagnosis of proctitis or mild UC (usually controlled by aminosalicylates and suppository steroid therapies)
4. Active symptoms which would exclude the patient from undergoing routine diagnostic colonoscopy i.e. evidence of active bowel obstruction, intestinal perforation, significant GI hemorrhage or known high-grade stricture
5. Body weight is less than 40 kg
6. Any malignant disease currently or in history
7. Renal failure and/ or hepatic failure (Glutamate Oxalate Transferase [GOT], Glutamic-Pyruvic Transaminase [GPT], total bilirubin, creatinine > twice the normal value)
8. Chronic hypotension (80 mmHg or lower systolic)
9. Therapeutic anticoagulation (Cumarine) or coagulation disorder
10. Active bacterial or viral infection, especially acute or chronic Hepatitis B or C virus infection, or HIV infection
11. Severe cardiovascular disease (New York Heart Association [NYHA] III-IV or Canadian Cardiovascular Society [CCS] III-IV), which would not permit any extracorporeal treatment
12. Breast feeding, pregnancy, drug abuse or dementia
13. Participation in another clinical study in the last 3 months

Date of first enrolment

01/11/2004

Date of final enrolment

31/12/2006

Locations

Countries of recruitment

Germany

Study participating centre

Apheresis Research Institute

Cologne

Germany

50935

Sponsor information

Organisation

Asahi Kasei Medical Europe GmbH (Germany)

Sponsor details

Lyoner strasse 44-48

Frankfurt

Germany

60528

Sponsor type

Industry

ROR

<https://ror.org/040cmp171>

Funder(s)

Funder type

Industry

Funder Name

Source of funding added as of 8 June 2007:

Funder Name

Asahi Kasei Medical Europe GmbH, Lyoner Strasse 44-48, 60528 Frankfurt (Germany)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration