The United Kingdom Transcatheter Aortic Valve Implantation (UK TAVI) Trial. A multi-centre randomised controlled trial to assess the clinical effectiveness and cost utility of TAVI, compared with conventional surgical aortic valve replacement (AVR), in patients with severe symptomatic aortic stenosis at intermediate or high operative risk

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
12/06/2013		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
12/06/2013	Completed	[X] Results		
Last Edited 18/05/2022	Condition category Circulatory System	[] Individual participant data		

Plain English summary of protocol

Background and study aims

Aortic stenosis is a narrowing of the valve through which blood flows as it leaves the heart, and affects 13% of people over the age of 70 years. It is usually due to age-related degeneration. If the narrowing becomes severe, chest pain, breathlessness, fluid retention and fainting are common, and the outlook is poor with a high risk of death within 2-3 years. The only effective conventional treatment is surgical replacement of the valve, which involves open chest surgery and temporarily stopping the heart, with use of a heart-lung machine. Surgical results are generally excellent but the elderly often have other medical problems that may result in an increased risk of death and complications. Transcatheter Aortic Valve Implantation (TAVI) is a recently developed technique to implant an artificial aortic valve without major surgery, using a catheter to deliver the valve to the heart through the arteries, which are usually accessed by puncturing the skin in the groin. In some cases, the valve is delivered directly through a small incision in the chest wall. The proposed study will compare TAVI with conventional surgery.

Who can participate?

Patients aged 70 years or over for whom either treatment is an option but in whom the risks of surgery are considered by an expert local medical team to be intermediate or high.

What does the study involve?

Each patient who consents to take part will be randomly assigned to receive either TAVI or

surgery. Outcomes in each group will be compared to see whether TAVI offers advantages in the short term and whether the long-term results, over 5 years, are as good as surgery. The cost implications for the NHS will be assessed to see if TAVI offers value for money.

What are the possible benefits and risks of participating? The results of the study will inform NHS policy for the use of TAVI and guide treatment decisions for future patients.

Where is the study run from? University of Leicester (UK)

When is the study starting and how long is it expected to run for? August 2013 to October 2022

Who is funding the study?
NIHR Health Technology Assessment (UK)

Who is the main contact? Dr William D. Toff w.toff@le.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr William D Toff

Contact details

Department of Cardiovascular Sciences University of Leicester Clinical Sciences Wing Glenfield Hospital Groby Road Leicester United Kingdom LE3 9QP

w.toff@le.ac.uk

Additional identifiers

Protocol serial number 14550; HTA 09/55/63

Study information

Scientific Title

The United Kingdom Transcatheter Aortic Valve Implantation (UK TAVI) Trial. A multi-centre randomised controlled trial to assess the clinical effectiveness and cost utility of TAVI, compared with conventional surgical aortic valve replacement (AVR), in patients with severe symptomatic aortic stenosis at intermediate or high operative risk

Acronym

UK TAVI

Study objectives

Aortic stenosis, narrowing of the valve through which blood flows as it leaves the heart, affects 1-3% of people over the age of 70 years. It is usually due to age-related degeneration. If the narrowing becomes severe, chest pain, breathlessness, fluid retention and fainting are common, and the outlook is poor with a high risk of death within 2-3 years.

The only effective conventional treatment is surgical replacement of the valve, which involves open chest surgery and temporarily stopping the heart, with use of a heart-lung machine. Surgical results are generally excellent but the elderly often have other medical problems that may result in an increased risk of death and complications.

Transcatheter Aortic Valve Implantation (TAVI) is a recently developed technique to implant an artificial aortic valve without major surgery, using a catheter to deliver the valve to the heart through the arteries, which are usually accessed by puncturing the skin in the groin. In some cases, the valve is delivered directly through a small incision in the chest wall.

The proposed study will compare TAVI with conventional surgery in a group of patients, aged 70 years or over, for whom either treatment is an option but in whom the risks of surgery are considered by an expert local medical team to be intermediate or high. Each patient who consents to take part will be randomly assigned to receive either TAVI or surgery. Outcomes in each group will be compared to see whether TAVI offers advantages in the short-term and whether the long-term results, over 5 years, are as good as surgery.

The cost implications for the NHS will be assessed to see if TAVI offers value for money. The results of the study will inform NHS policy for the use of TAVI and guide treatment decisions for individual patients.

The primary study hypothesis is that in patients with severe symptomatic aortic stenosis who are at intermediate or high operative risk, Transcatheter Aortic Valve Implantation (TAVI) is non-inferior to surgical aortic valve replacement in respect of death from any cause at one year.

Ethics approval required

Old ethics approval format

Ethics approval(s)

First Medical Research Ethics Committee (MREC), 10/04/2013, 13/LO/0451

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Cardiovascular; Subtopic: Cardiovascular (all Subtopics); Disease: Cardiac Surgery

Interventions

- 1. Surgical AVR, Surgical Aortic Valve Implantation
- 2. TAVI, Transcatheter Aortic Valve Implantation

Follow Up Length: 60 month(s); Study Entry: Single Randomisation only

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

All-cause mortality; Timepoint(s): One year

Key secondary outcome(s))

- 1. All-cause mortality at 30 days, 2, 3, 4 and 5 years
- 2. Cardiovascular mortality at 30 days and annually to 5 years
- 3. Quality-adjusted survival at 3 months and annually to 5 years
- 4. Stroke at 30 days and annually to 5 years
- 5. Re-intervention at 30 days and annually to 5 years
- 6. Death from any cause or stroke at 30 days and annually to 5 years
- 7. Death from any cause or disabling stroke at 30 days and annually to 5 years
- 8. Death from any cause, stroke or re-intervention at 30 days and annually to 5 years
- 9. Quality of life (Minnesota Living With Heart Failure Questionnaire at 6 weeks and 1 year; EQ-
- 5D-5L at 2 weeks, 6 weeks, 3 months, 6 months and annually to 5 years)
- 10. Symptoms and functional capacity (CCS scale and NYHA class at 6 weeks, 3 months, 6 months and annually to 5 years; Nottingham EADL and 6-min walk at 6 weeks and 1 year)
- 11. Cognitive function (Mini-Mental-State-Index at 6 weeks and 1 year)
- 12. Procedural success and in-hospital complications
- 13. Duration of post-procedural hospital stay
- 14. Vascular complications at 30 days and 1 year
- 15. Major bleeding at 30 days and 1 year
- 16. Infective endocarditis at 30 days and annually to 5 years
- 17. Myocardial infarction at 30 days and annually to 5 years
- 18. Conduction disturbance requiring permanent cardiac pacing pre-discharge and at 1 year
- 19. Renal replacement therapy at 30 days and 1 year
- 20. Echocardiographic measures (left ventricular ejection fraction, mass, dimensions and volumes; aortic regurgitation; aortic valve gradient and area, at 6 weeks and 1 year)
- 21. Costs, cost-utility and incremental cost-effectiveness ratio at 1 year, 5 years and estimated over lifetime using an extrapolation model

Completion date

31/10/2022

Eligibility

Key inclusion criteria

- 1. Severe symptomatic aortic stenosis referred for intervention
- 2. Age \geq 80 years or age \geq 70 years with intermediate or high operative risk from conventional AVR, as determined by the MDT
- 3. Both conventional AVR and TAVI deemed to be acceptable treatment options;
- 4. Participant able and willing to give written informed consent
- 5. Participant able (in the investigators opinion) and willing to comply with all study requirements Target gender: male and female; lower age limit: 70 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

ΔII

Total final enrolment

913

Key exclusion criteria

- 1. Intervention deemed inappropriate due to co-morbidity or frailty
- 2. Life expectancy less than one year due to co-morbidity
- 3. Previous AVR or TAVI
- 4. Technically unsuitable for either AVR or TAVI
- 5. Concomitant coronary artery disease requiring revascularisation for which only surgery is considered appropriate
- 6. Predominant aortic regurgitation
- 7. Severe mitral regurgitation or likely need for concomitant surgery or cardiac intervention other than planned coronary artery surgery or percutaneous coronary intervention as part of the treatment strategy

Date of first enrolment

01/08/2013

Date of final enrolment

31/03/2018

Locations

Countries of recruitment

United Kingdom

England

Study participating centre University of Leicester Leicester United Kingdom LE3 9QP

Sponsor information

Organisation

University of Leicester (UK)

ROR

https://ror.org/04h699437

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme; Grant Codes: 09/55/63

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		17/05/2022	18/05/2022	Yes	No
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes