

# Asthma Reduction with Inhaled corticoSteroids in Children with a high risk for the development of asthma (At RISC)

<b>Submission date</b> 27/01/2006	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 27/01/2006	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 25/08/2009	<b>Condition category</b> Respiratory	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr A.J. Nijholt

**Contact details**  
Maastricht University  
Department of General Practice  
P.O. Box 616  
Maastricht  
Netherlands  
6200 MD  
+31 (0)43 3884186  
annemiek.nijholt@hag.unimaas.nl

## Additional identifiers

**Protocol serial number**  
NTR397

## Study information

**Scientific Title**

**Acronym**

At RISC

**Study objectives**

Asthma reduction at age 6 is possible with inhaled corticosteroids in children of 1-4.5 years old with a familial predisposition for asthma in the first degree, who develop wheeze symptoms.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Received from local medical ethics committee

**Study design**

Multicentre randomised double blind placebo controlled parallel group trial

**Primary study design**

Interventional

**Study type(s)**

Prevention

**Health condition(s) or problem(s) studied**

Asthma

**Interventions**

After written parental informed consent has been obtained the children are randomly assigned to receive one of the following drug treatments for 12 months:

First 4 weeks:

Budesonide 2 puffs 200 ug/puff MDI via Nebunette spacer with appropriate facemask

Next 11 months:

Budesonide 1 puff 200 ug MDI via Nebunette spacer with appropriate facemask

OR

Placebo-to-match-budesonide MDI via Nebunette spacer with appropriate facemask

**Intervention Type**

Other

**Phase**

Not Specified

**Primary outcome(s)**

The determination of the effect of a course of 1 month budesonide 400 ug MDI via Nebunette and 11 months budesonide 200 ug versus placebo MDI via Nebunette on the development of asthma at the age of 6 years.

**Key secondary outcome(s))**

1. Differences between the group who received treatment with budesonide versus the group who was treated with placebo medication occur in lung function characteristics, the body height,

the combined asthma score, the presented and reported symptoms and exacerbations, and the adverse events

2. The cost-effectiveness of the treatment and the quality of life with the course of budesonide will be assessed
3. At age 6 changes in control group versus intervention group are described in the total IgE and the specific IgE for cat, dog, and house dust mite

**Completion date**

15/01/2008

## Eligibility

**Key inclusion criteria**

1. Male and female patients aged 1-4.5 years with a familial predisposition for asthma in the first degree, who have experienced at least 2 separate periods with wheeze lasting at least 2 days each documented by their general practitioner
2. For each patient who enters the study a written informed consent by the parent or guardian of the patient should be obtained

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Child

**Lower age limit**

1 years

**Upper age limit**

4.5 years

**Sex**

All

**Key exclusion criteria**

1. Patients who have been treated with pulmonary anti-inflammatory inhaled drugs for more than 2 weeks or anti-inflammatory oral drugs for more than 1 week preceding the study
2. Patients who have been hospitalised for asthma in the 2 weeks prior to the study
3. Patients who have serious respiratory morbidity (e.g. broncho-pulmonary dysplasia, cystic fibrosis, tuberculosis)
4. Patients, who have laboratory or clinical evidence of serious uncontrolled systemic disease (as judged by the investigator)
5. Patients with anatomical abnormalities of the upper airways or lungs
6. Patients currently participating in another drug intervention study
7. When the general practitioner considers it detrimental to the patient to participate in the study

**Date of first enrolment**

10/09/2001

**Date of final enrolment**

15/01/2008

## Locations

**Countries of recruitment**

Netherlands

**Study participating centre**

**Maastricht University**

Maastricht

Netherlands

6200 MD

## Sponsor information

**Organisation**

Care and Public Health Research Institute (CAPHRI), University Maastricht (Netherlands)

**ROR**

<https://ror.org/02jz4aj89>

## Funder(s)

**Funder type**

Charity

**Funder Name**

Netherlands Asthma Foundation (Netherlands)

**Funder Name**

Astra Zeneca BV (Netherlands)

## Results and Publications

## **Individual participant data (IPD) sharing plan**

### **IPD sharing plan summary**

Not provided at time of registration