Randomised phase III study of the local treatment of liver metastases by radiofrequency combined with chemotherapy versus chemotherapy alone in patients with unresectable colorectal liver metastases

Submission date	Recruitment status	Prospectively registered		
04/06/2003	Stopped	☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
23/07/2003	Stopped Condition category	[X] Results		
Last Edited		Individual participant data		
23/04/2021	Cancer	Record updated in last year		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Wendy Wood

Contact details

CR UK and UCL Cancer Trials Centre 90 Tottenham Court Road, London London United Kingdom W1T 4TJ +44 (0)20 7679 9858 W.Wood@CTC.UCL.AC.UK

Additional identifiers

ClinicalTrials.gov (NCT)

NCT00043004

Protocol serial number

EORTC 40004

Study information

Scientific Title

Randomised phase III study of the local treatment of liver metastases by radiofrequency combined with chemotherapy versus chemotherapy alone in patients with unresectable colorectal liver metastases

Acronym

CLOCC

Study objectives

The liver is the common site of relapse in patients with colorectal cancer. Chemotherapy is the treatment of choice for patients where surgical removal of lesions is not possible. The outcome for patients with unresectable liver metastases treated with chemotherapy is poor; five year survival of less than 1% and median survival less than one year.

Tumour destruction using radio frequency may offer an improvement in outcome. The purpose of this trial is to treat patients who have inoperable metastases with chemotherapy or RadioFrequency Ablation (RFA) and chemotherapy. The chemotherapy schedule used will be a combination of 5-Fluorouracil (5FU), leucovorin and oxaliplatin.

Ethics approval required

Old ethics approval format

Ethics approval(s)

LREC approval for UK sites

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Liver metastases in colorectal cancer

Interventions

Arm one: Radiofrequency ablation and chemotherapy

Arm two: Chemotherapy alone.

Radiofrequency:

- 1. Tumour ablative technique which can be combined with surgery
- 2. Imaging of procedure by ultrasound
- 3. During radiofrequency thermal heat injury leads to tissue coagulation

- 4. Effective in the local destruction of liver metastases
- 5. Suitable for lesions which are not totally resectable due to number or location and close to large blood vessels

Chemotherapy:

- 1. Oxaliplatin 85mg/m^2 day one only
- 2. Leucovorin 175mg/m² day one only
- 3. 5-Fluorouracil 400mg/m² bolus
- 4. 5FU 2400mg/m^2 at 46 hours intravenous infusion
- 5. Fortnightly for six months i.e. 12 courses

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

5-Fluorouracil (5FU), LeucoVorin (LV) and oxaliplatin.

Primary outcome(s)

To determine whether radiofrequency in combination with chemotherapy leads to superior overall survival compared to chemotherapy alone.

Key secondary outcome(s))

- 1. Progression free survival
- 2. Quality of Life
- 3. Health economics

Completion date

30/09/2007

Reason abandoned (if study stopped)

Participant recruitment issue

Eligibility

Key inclusion criteria

Patients with unresectable colorectal liver metastases:

- 1. Resection of primary tumour
- 2. Unresectable liver metastases
- 3. No extrahepatic disease
- 4. Total number of metastatic deposits less than ten
- 5. Maximum diameter of lesions 4cm
- 6. Patient consent
- 7. Aged 18 to 80 years
- 8. Normal haematology and biochemistry

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

119

Key exclusion criteria

- 1. World Health Organisation (WHO) status of more than or equal to one
- 2. Hepatic sufficiency (Bilirubin, Alkaline Phosphatase more than three times the Upper Limit of Normal [ULN])
- 3. Peripheral neuropathy Common Toxicity Criteria (CTC) grade more than or equal to one
- 4. Uncontrolled congestive heart failure, angina pectoris, hypertension or arrythmia
- 5. Any contraindication to the use of 5FU/LV/Oxaliplatin
- 6. Active infection
- 7. Pregnant or lactating women

Date of first enrolment

01/04/2003

Date of final enrolment

30/09/2007

Locations

Countries of recruitment

United Kingdom

England

Study participating centre CR UK and UCL Cancer Trials Centre

London United Kingdom W1T 4TJ

Sponsor information

Organisation

Cancer Research UK

ROR

https://ror.org/054225q67

Funder(s)

Funder type

Charity

Funder Name

Cancer Research UK (ref: C8374/A3488)

Alternative Name(s)

CR_UK, Cancer Research UK - London, Cancer Research UK (CRUK), CRUK

Funding Body Type

Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/10/2012	23/04/2021	Yes	No
Abstract results				No	No
Abstract results		20/05/2008	23/04/2021	No	No
Plain English results				No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes