# What is the effect of alternative graphical displays used to present the benefits of antibiotics for sore throat on decisions about whether to use them?

Submission date	Recruitment status No longer recruiting	Prospectively registered		
09/03/2007		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
04/05/2007	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
02/09/2009	Ear, Nose and Throat			

#### Plain English summary of protocol

Not provided at time of registration

#### Contact information

#### Type(s)

Scientific

#### Contact name

Ms Cheryl Carling

#### Contact details

PO box 7004
St Olavs Plass
Oslo
Norway
N-0130
+47 98627327
cheryl.carling@kunnskapssenteret.no

### Additional identifiers

**EudraCT/CTIS** number

**IRAS** number

ClinicalTrials.gov number

#### Secondary identifying numbers

N/A

# Study information

#### Scientific Title

#### Acronym

HIPPO (Health Information Project, Presentation Online)

#### Study objectives

The null hypotheses for this study are that in terms of congruence between peoples' values and treatment decisions there is no difference between the following graphical displays used to present the benefits of antibiotics for sore throat:

- 1. Faces at 3 days and Bar graphs at 3 days
- 2. Bar graphs of Duration of symptoms and Bar graphs at 3 days, and
- 3. Bar graphs at 0, 3 and 7 days and Bar graphs at 3 days.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

The University at Buffalo (New York, USA), Health Sciences Institutional Review Board, approved on 15 May 2002.

#### Study design

Randomised controlled trial

#### Primary study design

Interventional

#### Secondary study design

Randomised controlled trial

#### Study setting(s)

Not specified

#### Study type(s)

**Not Specified** 

#### Participant information sheet

#### Health condition(s) or problem(s) studied

Sore throat (hypothetical scenario)

#### **Interventions**

Presentations of graphical displays showing evidence of penicillin's effect on sore throat.

#### Intervention Type

#### Other

#### **Phase**

**Not Specified** 

#### Primary outcome measure

Hypothetical treatment decision to take or not to take penicillin.

#### Secondary outcome measures

Understanding of and satisfaction with information, and preferred graphical display.

#### Overall study start date

01/09/2004

#### Completion date

31/10/2004

# **Eligibility**

#### Key inclusion criteria

Fluency in Norwegian and at least 18 years of age.

#### Participant type(s)

**Patient** 

#### Age group

Adult

#### Lower age limit

18 Years

#### Sex

**Not Specified** 

#### Target number of participants

3000

#### Key exclusion criteria

Previous participation in this trial.

#### Date of first enrolment

01/09/2004

#### Date of final enrolment

31/10/2004

# **Locations**

Countries of recruitment

# Study participating centre PO box 7004

Oslo Norway N-0130

# Sponsor information

#### Organisation

Norwegian Knowledge Centre for the Health Services (Norway)

#### Sponsor details

PO box 7004 St Olavs Plass Oslo Norway N-0130 +47 (0)23 25 50 00 post@kunnskapssenteret.no

#### Sponsor type

Other

#### Website

http://www.kunnskapssenteret.no/

#### **ROR**

https://ror.org/01thff661

# Funder(s)

#### Funder type

Research council

#### **Funder Name**

Norwegian Research Council (Norway)

## **Results and Publications**

#### Publication and dissemination plan

Not provided at time of registration

#### Intention to publish date

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2009		Yes	No