

# ACES primary care: Alternative cervical screening in primary care

<b>Submission date</b> 08/03/2022	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 09/03/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 11/04/2023	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Cervical screening can save lives from cervical cancer, yet only 7 in 10 women in the UK attend screening, the lowest rate in 20 years. Reasons include embarrassment, fear of examination and inconvenience. Cervical screening is carried out by collecting cells from the cervix (neck of the womb) with a soft brush. These cells are tested for a virus known to cause cancer called human papillomavirus (HPV). If HPV is detected, the cells are examined under the microscope. If they look abnormal, the woman is referred to colposcopy clinic, where cells that are found to be 'pre-cancerous' (cells with the potential to become cancer cells) are identified and treated. To increase screening rates, vaginal 'self-sampling' has been tried, where a woman collects cells from her vagina at home and returns the sample by post, however only 1 in 10 women return the sample. There is therefore an urgent need for new ways to reverse declining rates of cervical screening.

We have developed a urine test that can detect HPV. This test has the potential to remove many of the current barriers to screening and could substantially increase the number of women attending. This study will see if a urine test can accurately identify women with cervical pre-cancer by comparing HPV detection rates in urine and cervical samples.

### Who can participate?

Women and people with a cervix attending routine cervical screening appointments at participating GP practices or NHS clinics.

### What does the study involve?

Individuals will be asked to complete some brief questions about their health and provide a first void urine sample. They will also be asked to complete a short questionnaire to understand views and preferences of current cervical screening attendees. Samples will be tested for high-risk HPV and HPV-positive samples will undergo methylation testing.

This study will help establish whether the clinical performance of urine testing is sufficient to recommend its use as an NHS cervical screening test.

### What are the possible benefits and risks of participating?

There are no immediate benefits to the individual taking part in this study. We will use the

results to help us know whether urine HPV testing could be a reasonable alternative to routine cervical screening. This could encourage more women to participate in cervical screening in the future.

We do not expect there to be any side effects of taking part.

Where is the study run from?

The University of Manchester (UK)

When is the study starting and how long is it expected to run for?

September 2020 to December 2024

Who is funding the study?

National Institute for Health Research (NIHR) (UK)

Who is the main contact?

Suzanne Carter (public), [suzanne.carter@manchester.ac.uk](mailto:suzanne.carter@manchester.ac.uk)

Prof. Emma Crosbie (scientific), [emma.crosbie@manchester.ac.uk](mailto:emma.crosbie@manchester.ac.uk)

### **Study website**

<https://sites.manchester.ac.uk/aces/>

## **Contact information**

### **Type(s)**

Public

### **Contact name**

Miss Suzanne Carter

### **Contact details**

The University of Manchester

Division of Cancer Sciences

School of Medical Sciences

Faculty of Biology Medicine and Health

St Mary's Hospital

Manchester

United Kingdom

M13 9WL

+44 161 701 6941

[suzanne.carter@manchester.ac.uk](mailto:suzanne.carter@manchester.ac.uk)

### **Type(s)**

Principal Investigator

### **Contact name**

Prof Emma Crosbie

### **ORCID ID**

<http://orcid.org/0000-0003-0284-8630>

### **Contact details**

The University of Manchester  
Division of Cancer Sciences  
School of Medical Sciences  
Faculty of Biology Medicine and Health  
St Mary's Hospital  
Manchester  
United Kingdom  
M13 9WL  
+44 161 701 6941  
emma.crosbie@manchester.ac.uk

## Additional identifiers

### EudraCT/CTIS number

Nil Known

### IRAS number

309113

### ClinicalTrials.gov number

Nil known

### Secondary identifying numbers

IRAS 309113, CPMS 51674

## Study information

### Scientific Title

Urine HPV testing for cervical screening in primary care

### Acronym

ACES Primary Care

### Study objectives

Urine HPV testing is an accurate alternative to routine cervical screening in general screening population

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Approved 14/02/2022, South West - Cornwall & Plymouth Research Ethics Committee (Ground Floor, Temple Quay House, 2 The Square, Bristol, BS1 6PN, UK; +44 (0)207 1048071; cornwallandplymouth.rec@hra.nhs.uk), ref 22/SW/0007

### Study design

Multicentre observational cross sectional study

### Primary study design

Observational

## **Secondary study design**

Cross sectional study

## **Study setting(s)**

GP practice

## **Study type(s)**

Screening

## **Participant information sheet**

<https://sites.manchester.ac.uk/aces/>

## **Health condition(s) or problem(s) studied**

Identification of cervical pre cancer in the general screening population using urine HPV testing

## **Interventions**

Current intervention as of 11/04/2023:

This study will see if a urine test can accurately identify individuals with cervical pre-cancer and those who continue to be HPV positive after treatment by comparing HPV detection rates in matched urine and cervical samples. Samples will be tested for high-risk HPV. HPV-positive samples will undergo methylation testing.

Prior to routine clinical procedures, we will collect a voided urine sample. Urine samples will be self-collected at the GP practice or other NHS clinic, in the privacy of the clinic bathroom. Urine will be collected with a Colli-Pee device, which reliably collects a standardised volume of first void urine. Urine collection must be done before routine procedures to mirror what would happen in 'real life' if a urine test were to replace routine screening. It must also be done on the same day as the cervical sample, to preclude changes in viral status between sampling time points affecting the validity of the results. A routine cervical screening ('Pap' smear) will then be taken as part of the participant's routine clinical cervical screening care.

We will directly compare HPV detection rates, concordance between urine and matched cervical samples and CIN2+ detection rates in urine samples.

If the participant attends for a follow-up visit (e.g. for treatment after initial assessment), we may ask them to provide a second or third set of samples, if they consent. This will help us understand more about how well the urine test could work during the natural history of HPV infection and the management of abnormal smears.

Participants will answer a short acceptability questionnaire to gauge their views on urine testing for cervical screening. Those who decline participation will be asked to record their reasons on a short questionnaire. This is entirely optional.

---

## **Previous intervention:**

This study will see if a urine test can accurately identify women with cervical pre-cancer and those who continue to be HPV positive after treatment by comparing HPV detection rates in matched urine and cervical samples. Samples will be tested for high risk HPV. HPV positive samples will undergo methylation testing.

Prior to routine clinical procedures, we will collect a voided urine sample. Urine samples will be self collected at the GP practice, in the privacy of the clinic bathroom. Urine will be collected with a Colli-Pee device, which reliably collects a standardised volume of first void urine. Urine collection must be done before routine procedures to mirror what would happen in 'real life' if a urine test were to replace routine screening. It must also be done on the same day as the cervical sample, to preclude changes in viral status between sampling time points affecting the validity of the results. A routine cervical screening ('Pap' smear) will then be taken as part of the participant's routine clinical cervical screening care.

We will directly compare HPV detection rates, concordance between urine and matched cervical samples and CIN2+ detection rates in urine samples.

If the participant attends for a follow up visit (e.g. for treatment after initial assessment), we may ask them to provide a second or third set of samples, if they consent. This will help us understand more about how well the urine test could work during the natural history of HPV infection and the management of abnormal smears.

Participants will answer a short acceptability questionnaire to gauge their views on urine testing for cervical screening. Those who decline participation will be asked to record their reasons on a short questionnaire. This is entirely optional.

## **Intervention Type**

Other

## **Primary outcome measure**

1. High risk HPV detection rate in matched urine and cervical samples at baseline
2. CIN2+ detection rates according to routine cervical screening outcomes

The above measures will be used to calculate sensitivity, specificity, negative predictive value (NPV) and positive predictive value (PPV) of the urine HPV test for CIN2+ detection compared to routine cervical screening

## **Secondary outcome measures**

1. Concordance of HR-HPV+ test results in matched urine and cervical samples measured as in the primary outcome measure at baseline
2. Presence of CIN2+ measured using urine HPV+ test combined with a methylation+ test at baseline (used to calculate diagnostic test accuracy (sensitivity, specificity, NPV and PPV)
3. Preference for urine compared to routine sampling for cervical screening assessed by participant questionnaire at baseline
4. Reasons for declining to take part in the study assessed by participant questionnaire at the time of declining

## **Overall study start date**

01/09/2020

## **Completion date**

31/12/2024

# **Eligibility**

## **Key inclusion criteria**

1. Age 24-70 years
2. Written, informed consent to participate
3. Undergoing routine NHS cervical screening or management of abnormal cervical screening

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

24 Years

**Upper age limit**

70 Years

**Sex**

Female

**Target number of participants**

1,500 - 2,000

**Key exclusion criteria**

1. Pregnant
2. Previous hysterectomy
3. Unable to provide a urine sample
4. Unable to understand the Participant Information Sheet and consent form
5. Any condition that would compromise participant safety or data integrity

**Date of first enrolment**

14/03/2022

**Date of final enrolment**

01/09/2024

**Locations****Countries of recruitment**

England

United Kingdom

**Study participating centre**

**Bowland Medical Practice**

52 Bowland Road

Baguley

Manchester  
United Kingdom  
M23 1JX

**Study participating centre**  
**Didsbury Medical Centre - Dr Whitaker**  
645 Wilmslow Road  
Didsbury  
Manchester  
United Kingdom  
M20 6BA

**Study participating centre**  
**Surrey Lodge Practice**  
11 Anson Road  
Victoria Park  
Manchester  
United Kingdom  
M14 5BY

**Study participating centre**  
**The Maples Medical Centre**  
2 Scout Drive  
Newall Green  
Manchester  
United Kingdom  
M23 2SY

**Study participating centre**  
**Ancoats Urban Village Medical Practice**  
Old Mill Street  
Manchester  
United Kingdom  
M4 6EE

**Study participating centre**  
**Hawthorn Medical Centre**  
Unit K, Fallowfield Shopping Centre  
Birchfields Road  
Fallowfield  
Manchester

United Kingdom  
M14 6FS

**Study participating centre**

**Central Manchester University Hospitals NHS Foundation Trust**  
Trust Headquarters, Cobbett House  
Manchester Royal Infirmary  
Oxford Road  
Manchester  
United Kingdom  
M13 9WL

## **Sponsor information**

**Organisation**

University of Manchester

**Sponsor details**

Faculty of Biology  
Medicine and Health  
Carys Bannister Building  
Dover Street  
Manchester  
England  
United Kingdom  
M13 9PL  
+44 (0)161 275 5436  
FBMHethics@manchester.ac.uk

**Sponsor type**

University/education

**Website**

[www.manchester.ac.uk](http://www.manchester.ac.uk)

**ROR**

<https://ror.org/027m9bs27>

## **Funder(s)**

**Funder type**

Government



**Funder Name**

National Institute for Health Research

**Alternative Name(s)**

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## Results and Publications

**Publication and dissemination plan**

Planned publication in a high impact peer-reviewed journal.

**Intention to publish date**

01/09/2025

**Individual participant data (IPD) sharing plan**

At the end of the project we will deposit a fully anonymised dataset in an open data repository where it will be permanently stored. Researchers at other institutions can access the anonymised data directly from the repository and use it for further research or to check our analysis and results

The data will be completely anonymised and stored on Figshare (<https://figshare.manchester.ac.uk/>). Researchers at other institutions can access the anonymised data directly from the repository. We are seeking consent from participants for this.

**IPD sharing plan summary**

Stored in publicly available repository

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Plain English results</a>			08/08/2022	No	Yes
<a href="#">HRA research summary</a>			28/06/2023	No	No