

Dyslipidemia in Japanese Patients with Rheumatoid Arthritis

Submission date 02/06/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 29/06/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 29/06/2010	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Taro Mawatari

Contact details
Maidashi 3-1-1
Higashi-ku
Fukuoka
Japan
812-8582

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title
Dyslipidemia in Japanese Patients with Rheumatoid Arthritis: A Retrospective Multicentre Observational Study

Study objectives

By investigating the prevalence of dyslipidemia and its relationships with disease activity, or medical interventions, it is possible to reduce subsequent mortality and morbidity of cardiovascular disease in patients with rheumatoid arthritis (RA).

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Institutional Review Board of the Kyushu University School of Medicine approved on the 10th of May 2010

Study design

Retrospective multicentre observational study

Primary study design

Observational

Study type(s)

Screening

Health condition(s) or problem(s) studied

Rheumatoid arthritis

Interventions

Non-interventional, observational study.

The blood samples from a cohort of patients being treated with lipid-lowering drugs will be taken at baseline, 2, 4, 6, 8 months after administration.

Information will be collected on the patient's history of cardiovascular events.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. To investigate the lipid profile and estimate the prevalence of dyslipidemia
2. To determine the relationships between the lipid profiles and the RA disease activity, or the treatments such as Disease Modifying Antirheumatic Drugs (DMARDs), biologics, glucocorticoids.

Key secondary outcome(s)

To investigate the effect of administered statins in the treatment of dyslipidemia in Japanese patients with RA.

Completion date

31/03/2013

Eligibility

Key inclusion criteria

1. Age > 18 yrs
2. Fulfilled the 1987 revised criteria for RA of the American College of Rheumatology
3. Access to medical record information

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

History of medical conditions other than RA which may affect lipid profile

Date of first enrolment

01/06/2009

Date of final enrolment

31/03/2013

Locations**Countries of recruitment**

Japan

Study participating centre

Maidashi 3-1-1

Fukuoka

Japan

812-8582

Sponsor information**Organisation**

Kyushu University (Japan)

ROR

<https://ror.org/00p4k0j84>

Funder(s)

Funder type

University/education

Funder Name

Graduate School of Medical Sciences, Kyushu University (Japan) - Department of Orthopaedic Surgery

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes