A prospective study to assess the effectiveness of lignocaine versus normal saline in the reduction of pain associated with dressing removal in fingertip injuries

Submission date	Recruitment status	Prospectively registered
28/09/2007	Stopped	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
28/09/2007	Stopped	Results
Last Edited	Condition category	Individual participant data
27/08/2008	Signs and Symptoms	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Mr D N Quinton

Contact details

Derby Hospitals NHS Foundation Trust
Derbyshire Royal Infirmary
London Road
Derby
United Kingdom
DE1 2QY
+44 01332 347141 ext: 2249
david.quinton@derbyhospitals.nhs.uk

Additional identifiers

Protocol serial number N0077183569

Study information

Scientific Title

Study objectives

Does local anaesthetic (lignocaine) reduce pain associated with dressing removal of fingertip injuries when compared to normal saline?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Signs and Symptoms: Pain

Interventions

Patients will be recruited from the Pulvertaft Hand Center upon clinic appointment. This is a stop interview /evaluation that takes place in the Hand Clinic. Certain data will be collected - age, sex, nature of injury, etc. This will be kept confidential. Patients will be asked whether they would like to be involved in the study. Those patients who agree will be asked to sign a consent form. The patients who do not wish to participate in the study will receive the normal standard of care for such injuries, ie. normal saline.

The following protocol will be set up:

- Affected finger will be immersed in 10 milliliters of local anaesthetic (lignocaine) or normal saline.
- The dressing is then removed by the sister in the clinic and the patient asked to log their level of pain. This will be achieved by a simple (linear analogue) scoring system.

The whole process will be randomised and blind to those involved in the study to avoid or prevent bias.

Added 27 August 2008: trial was stopped.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

lignocaine versus normal saline

Primary outcome(s)

The reduction in the level of pain on dressing removal of patients with fingertip wounds as shown on the linear visual analogue pain scale, 0-10, 0 being the lowest level of pain and 10 the highest level of pain.

Key secondary outcome(s))

No secondary outcome measures

Completion date

31/01/2008

Reason abandoned (if study stopped)

Too much bureaucracy

Eligibility

Key inclusion criteria

- 1. Patients sustaining a nail bed injury
- 2. Patients who are old enough to give consent (over 16 years old)
- 3. Patients willing to participate

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Not Specified

Key exclusion criteria

- 1. Patients who are unable to give consent
- 2. Previous injury or surgery to the presenting digit
- 3. Patients within any other concomitant trial involving analgesics, patient allergy to lignocaine

Date of first enrolment

21/08/2006

Date of final enrolment

31/01/2008

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Derby Hospitals NHS Foundation Trust
Derby
United Kingdom
DE1 2QY

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

Derby Hospitals NHS Foundation Trust (UK), NHS R&D Support Funding

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration