Evaluation of the efficacy of buprenorphine injection around the stellate ganglion in pain syndromes of the upper body-half

Submission date	Recruitment status No longer recruiting	Prospectively registered		
12/09/2003		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
12/09/2003	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
15/04/2008	Signs and Symptoms			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Andreas Goebel

Contact details

c/o Pain Relief Unit
The Churchill
Headington
Oxford
United Kingdom
OX3 7LJ
+44 (0)1865 226161
andreasgoebel@rocketmail.com

Additional identifiers

Protocol serial number N0176115639

Study information

Scientific Title

Study objectives

- 1. To determine if 0.025 mg buprenorphine, a highly lipohilic morphine derivate, is more efficient than saline in relieving chronic, non-nociceptive pain when injected in proximity to the stellate ganglion. The stellate ganglia are collections of autonomic nerve cells and synapses located on either side of the neck.
- 2. A minimally traumatising method for injection of the stellate ganglion, not previously been published in the English literature will be used; this method is applied in German pain clinics. Our objective is to ascertain the usefulness of this method in conjunction with stellate opioid injections.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pain

Interventions

Two interventions:

- 1. Normal injection into buttocks
- 2. Injection into neck

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Buprenorphine

Primary outcome(s)

- 1. Pain reduction over the first 8 hours following stellate injection, as calculated from the median of four, two-hourly measurements of present pain reduction using a visual analogue scale (VAS) and compared between buprenorphine and placebo stellate injections
- 2. Absolute pain levels as expressed using a VAS scale and calculated from the median of four, two-hourly measurements of present pain intensity (PPI), each divided by the pre-injection PPI and compared between buprenorphine and placebo stellate injections

Key secondary outcome(s))

Not provided at time of registration

Completion date

15/11/2004

Eligibility

Key inclusion criteria

Not provided at time of registration

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/10/2002

Date of final enrolment

15/11/2004

Locations

Countries of recruitment

United Kingdom

England

Study participating centre c/o Pain Relief Unit

Oxford United Kingdom OX3 7LJ

Sponsor information

Organisation

Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

Oxford Radcliffe Hospitals NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/04/2008		Yes	No