

The role of 3D laparoscopic rectal surgery: a randomised controlled trial

Submission date 09/12/2015	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 16/12/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 10/01/2023	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-trial-to-improve-keyhole-surgery-for-rectal-cancer>

Contact information

Type(s)

Public

Contact name

Mr Nader Francis

Contact details

Yeovil District Hospital
Yeovil
United Kingdom
BA21 4AT

Type(s)

Scientific

Contact name

Mr Nathan Curtis

Contact details

Yeovil District Hospital
Yeovil
United Kingdom
BA21 4AT

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

2D vs. 3D laparoscopic anterior resection: a randomised controlled trial

Study objectives

Using 3D imaging reduces intra-operative technical errors enacted during laparoscopic anterior resection.

Ethics approval required

Old ethics approval format

Ethics approval(s)

South Central Berkshire B Research Ethics Committee, 24/02/2016, ref: 16/SC/0118

Study design

Multicentre randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Rectal adenocarcinoma

Interventions

Patients will be identified at local colorectal Multi-Disciplinary Team meetings and outpatient clinics. Potentially eligible patients will be approached by GCP (Good Clinical Practice) trained staff with delegated responsibility at each centre, who will discuss the study with the patient and answer any questions. Patients will be given a written site specific participant information sheet to take away. If willing and eligibility is confirmed, written consent will be obtained and the patient randomised. All trial participating centres and trial surgeons are experienced in using 3D imaging systems.

Centralised randomisation will take place at Yeovil Hospital Clinical Research Unit. Allocation into one of two groups will be determined using pre-prepared computerised random number generation (zero or one) to allocate in a 1:1 ratio:

1. Group 1 patients undergo laparoscopic anterior resection using a 2D imaging system.
2. Group 2 patients undergo laparoscopic anterior resection using a 3D imaging system.

Intervention Type

Device

Primary outcome(s)

The total number of technical errors during each operation. Each operation will be digitally recorded and analysed unedited by a blinded assessor using Objective Clinical Human Reliability Analysis (OCHRA)

Key secondary outcome(s)

1. Total operating time
2. Operative blood loss
3. Conversion to open surgery
4. Postoperative length of stay
5. 30 day readmission
6. Histological outcomes, including mesorectal specimen quality, which will be assessed by a pathologist using standardised criteria
7. Postoperative complications using the Clavien-Dindo classification
8. 30 day mortality
9. Surgeon-reported feedback on the adverse effects of the imaging system will also be recorded such as blurred vision, altered vision, light headedness and or dizziness
10. Surgeon-reported cognitive load assessment

All complications occurring in hospital (index stay or re-admission) will be recorded up to 30 days.

Completion date

30/04/2018

Eligibility

Key inclusion criteria

1. Patients diagnosed with adenocarcinoma of the rectum (<15cm from anal verge as measured by staging MRI)
2. Scheduled to undergo elective laparoscopic TME/ partial mesorectal excision surgery with curative intent, with or without neo-adjuvant treatment and with or without defunctioning stoma formation
3. Written informed patient consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

88

Key exclusion criteria

1. History of inflammatory bowel disease
2. Patients requiring abdomino-perineal excision or surgery without anastomosis
3. Advanced tumours involving adjacent organs (TNM5 T4aNxMx)
4. Surgery performed with palliative intent or under unplanned/emergency settings
5. Previous treatment (radiotherapy/abdominopelvic surgery) for endometrial, ovarian, prostate, bladder (TURBT ok), anal or vaginal cancer
6. Patient or surgeon refusal to enter study or accept randomisation result

Date of first enrolment

02/01/2016

Date of final enrolment

30/03/2018

Locations

Countries of recruitment

United Kingdom

England

Wales

Study participating centre

Yeovil District Hospital

Higher Kingston

Yeovil

United Kingdom

BA21 4AT

Study participating centre

Royal Surrey County Hospital

Egerton Road

Guildford

United Kingdom

GU2 7XX

Study participating centre

Frimley Park Hospital

Portsmouth Road

Frimley

United Kingdom

GU16 7UJ

Study participating centre
University Hospital of Wales
Heath Park
Cardiff
United Kingdom
CF14 4XW

Study participating centre
Queen Alexandra Hospital
Cosham
Portsmouth
United Kingdom
PO6 3LY

Sponsor information

Organisation
Yeovil District Hospital NHS Foundation Trust

ROR
<https://ror.org/00v5nyn36>

Funder(s)

Funder type
Industry

Funder Name
European Association of Endoscopic Surgeons

Funder Name
Karl Storz GmbH & Co

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2019	13/03/2019	Yes	No
Results article	Factors predicting operative difficulty	01/12/2019	10/01/2023	Yes	No
HRA research summary			28/06/2023	No	No
Plain English results			24/03/2021	No	Yes