

# Neural correlates of cognitive motor interference (CMI) while walking

<b>Submission date</b> 17/11/2008	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 27/11/2008	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 06/06/2016	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Helen Dawes

**Contact details**  
School of Life Sciences  
Oxford Brookes University  
Oxford  
United Kingdom  
OX3 0BP

## Additional identifiers

**Protocol serial number**  
TSA 2007/09

## Study information

**Scientific Title**  
Cortical activation of cognitive motor interference while walking: a near-infrared spectroscopic study

**Acronym**  
CMI

## **Study objectives**

1. There may be a relationship between the degree of regional brain activation and/or the temporal activation pattern and performance in cognitive and motor tasks under dual task conditions
2. This relationship may be different following a stroke than in healthy controls

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Oxfordshire Research Ethics Committee C, 24/07/2008, ref: 08/H0606/55

## **Study design**

Observational cross-sectional study

## **Primary study design**

Observational

## **Study type(s)**

Diagnostic

## **Health condition(s) or problem(s) studied**

Stroke

## **Interventions**

Baseline assessments include:

1. The Mini Mental State Examination: the MMSE is a brief screening tool to provide quantitative assessment of cognitive function. It consists of 11 simple questions or tasks. Typically, they are grouped into seven domains; orientation to time, orientation to place, registration of three words, attention and calculation.
2. Edinburgh Handedness Inventory: this is a brief questionnaire of 10 short items, which determines subject's hand dominance.
3. Barthel Index: a simple index of independence in activities of daily living (ADL). It consists of 10 common ADL activities, which they are assessed for independence/dependence.
4. Modified Physical Activity Scale for the elderly: a self reported physical activity questionnaire that covers one week.
5. Berg Balance Scale (BBS): the BBS provides a quantitative assessment of balance. It consists of 14 items requiring subjects to maintain or complete movement tasks of varying levels of difficulty. All items are common to everyday life.

The results of the assessments above will be reported together with the primary/secondary outcomes, as they might correlate with differences in CMI.

Brain activation in both stroke and control subjects will be examined by near-infrared spectroscopy (NIRS) alongside behavioural performance of walking and cognitive tasks (calculation). The participants will be asked to perform the concurrent tasks when seated and walking. Gait parameters will be measured using a well-validated gait analysis system (Xsens, The Netherlands).

## **Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome(s)**

Task-related changes in oxygenated and deoxygenated haemoglobins in the frontal lobe as measured by NIRS.

**Key secondary outcome(s)**

1. Rate of backward calculation (number per minute)
2. Number of errors in backward calculation (errors per trial)
3. Spatiotemporal gait measures (speed, cadence, step time, stride time, step length, stride length, step width)

**Completion date**

01/07/2010

**Eligibility****Key inclusion criteria**

Patients will be selected from those treated in the Oxford Radcliffe Hospitals who are:

1. More than 6 months following a first stroke
2. Both males and females, aged 45 to 80 years
3. With an ischaemic infarct upon computed tomography (CT) or magnetic resonance imaging (MRI) scan
3. Able to perform a simple reciprocal bilateral foot tapping task, walk safely on a treadmill with or without mobility aids
4. Be able to give informed consent

Sex and age-matched controls will be chosen who do not have a known neurological disease, a history of hypertension, cardiac disease or diabetes.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

1. Dementia
2. Aphasia significantly limiting communication
3. History of previous symptomatic strokes or neurological disease
4. Known psychiatric disease or claustrophobia, or other conditions precluding safe MRI (e.g., pacemaker or other metal implant)

**Date of first enrolment**

01/07/2008

**Date of final enrolment**

01/07/2010

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre**

**Oxford Brookes University**

Oxford

United Kingdom

OX3 0BP

## Sponsor information

**Organisation**

Oxford Brookes University (UK)

**ROR**

<https://ror.org/04v2twj65>

## Funder(s)

**Funder type**

Charity

**Funder Name**

Stroke Association (ref: TSA 2007/09)

**Alternative Name(s)**

TheStrokeAssociation, TheStrokeAssoc

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

Associations and societies (private and public)

**Location**

United Kingdom

## **Results and Publications**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**