

Efficacy of a web and text messaging based intervention to reduce problem drinking in young people

| | | |
|--|---|--|
| Submission date 24/06/2014 | Recruitment status No longer recruiting | <input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol |
| Registration date 10/07/2014 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results |
| Last Edited 04/08/2017 | Condition category Mental and Behavioural Disorders | <input type="checkbox"/> Individual participant data |

Plain English summary of protocol

Background and study aims

Binge drinking is drinking a significant number of alcoholic beverages with the intention of having enough to become intoxicated, or drunk, over a short period of time. It can lead to antisocial behavior and violence, and also to a number of health problems; this not only affects the individual, it also results in increased costs to healthcare services. There have been studies involving the use of tailored feedback generated by a computer programme that have been successful in reducing binge drinking among university and college students in the United States. It works by providing the student with information on their own drinking habits compared with what is considered the social norm in order to change their perception of what they would consider normal drinking behavior. It is hoped that a similar approach will reduce alcohol use in young people with a lower level of education as well, but little research has been done on this as yet. It is thought, however, that shorter and more frequent feedback messages may be needed for individuals with a poorer educational background and that they might benefit from getting support for a longer time period in order to change their drinking behavior long-term. Text messaging (SMS) is a proactive and low cost way of sending short messages at any time and place and it is a way of delivering tailored information to young people at times when they are most likely to drink alcohol. The main aim of this study is to assess how successful a combined web and text messaging based programme (the intervention) is reducing problem drinking in young people with different educational backgrounds.

Who can participate?

Young people over the age of 15 who have a mobile phone.

What does the study involve?

Participants are randomly allocated to one of two groups. Group 1 receive the intervention and have their alcohol use monitored. Group 2 just have their alcohol use monitored. Group 1 get a individually tailored web-based feedback on the number of drinks they have each week compared to a reference group that represents what would be the social norm for their age and gender. The number of times they have indulged in risky drinking (5 drinks on any one occasion for males, 4 times for females) is recorded. They are told how much they spend on alcohol and

how many calories their drinks contain. They also get individually tailored mobile phone messages over a 3 month period that encourage the participant to keep their drinking to a safe limit and resist alcohol altogether. The drinking habits of the participants are then followed up 6 months later.

What are the possible benefits and risks of participating?

We expect that the intervention will reduce binge drinking. There are considered to be no risks.

Where is the study run from?

Swiss Research Institute for Public Health and Addiction (Switzerland)

When is the study starting and how long is it expected to run for?

September 2014 to August 2016

Who is funding the study?

Swiss National Science Foundation (Switzerland)

Who is the main contact?

Dr. Severin Haug

severin.haug@isgf.uzh.ch

Contact information

Type(s)

Scientific

Contact name

Dr Severin Haug

Contact details

Konradstrasse 32

Zurich

Switzerland

8031

+41 (0) 44 448 11 74

severin.haug@isgf.uzh.ch

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

100014_149209/1

Study information

Scientific Title

Efficacy of a web and text messaging based intervention to reduce problem drinking in young people: a cluster-randomized controlled trial

Study objectives

The main objective of the planned study is to test the efficacy of a combined web and text messaging based intervention to reduce problem drinking in young people. Our main hypothesis is that the intervention will be more effective than assessment only in reducing indicators of problem drinking assessed at six month follow-up.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics committee of the Philosophical Faculty of the University of Zurich, Switzerland, 24/06/2014

Study design

Single center, two-arm, cluster randomized, single blind controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Other

Study type(s)

Prevention

Participant information sheet

Not available in web format, please use the contact details below to request a participant information sheet.

Health condition(s) or problem(s) studied

Alcohol consumption, Problem drinking

Interventions

Intervention Group:

Web-based feedback including individually tailored graphical and textual information concerning:

1. The number of drinks consumed per week in relation to the age and gender-specific reference group
2. Financial costs of drinking
3. Calories consumed with alcoholic drinks
4. Number of RSOD occasions in relation to the age and gender-specific reference group

The participants in the intervention group will receive a single web-based feedback immediately after the baseline assessment.

Individually tailored mobile phone text messages to stimulate

1. Positive outcome expectations to drink within low-risk limits
2. Self-efficacy to resist alcohol
3. Planning processes to translate intentions to resist alcohol into action

Participants will receive up to two weekly text messages over a time period of 3 months.

Control Group: No intervention

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Frequency of risky single occasion drinking (RSOD) occasions in the last 30 days (How often did you have (male: 5 female: 4) or more drinks on one occasion in the last 30 days?).

Secondary outcome measures

1. Quantity of alcohol consumption, assessed by a 7 day drinking calendar for which participants are asked to think about a typical week in the past month and, for each day, to record the number of standard drinks they typically consume on that day.
2. Peak blood alcohol concentration assessed by asking participants to report the number of standard drinks consumed and the duration of their heaviest drinking episode in the previous 30 days. This information will be used along with the sex and weight to calculate an estimated peak blood alcohol concentration based on the Widmark Formula.
3. Normative misperceptions of alcohol consumption.

Overall study start date

01/09/2014

Completion date

31/08/2016

Eligibility

Key inclusion criteria

1. Age > 15 years
2. Possession of a mobile phone

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

n=467 per study group (intervention and control) resulting in a total of n=934 study participants

Key exclusion criteria

None

Date of first enrolment

01/09/2014

Date of final enrolment

31/08/2016

Locations

Countries of recruitment

Switzerland

Study participating centre

Konradstrasse 32

Zurich

Switzerland

8031

Sponsor information

Organisation

Swiss Research Institute for Public Health and Addiction (Switzerland)

Sponsor details

c/o Severin Haug

Konradstrasse 32

Zurich

Switzerland

8031

Sponsor type

Research organisation

Website

<http://www.isgf.ch>

ROR

<https://ror.org/02crff812>

Funder(s)

Funder type

Research organisation

Funder Name

Funding for this study is provided by the Swiss National Science Foundation (No. 100014_149209 /1) (Switzerland)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|----------------------------------|----------|--------------|------------|----------------|-----------------|
| Protocol article | protocol | 07/08/2014 | | Yes | No |
| Results article | results | 01/02/2017 | | Yes | No |