

# Intravenous glutamine-dipeptide and early nasojejunal feeding in severe acute pancreatitis

<b>Submission date</b> 01/08/2012	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 27/09/2012	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 27/11/2015	<b>Condition category</b> Digestive System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

We are carrying out a study on patients with severe acute pancreatitis, which is a life-threatening disease where the pancreas becomes inflamed. Parenteral nutrition, also known as intravenous feeding, is a method of getting nutrients into the body through the veins. Other studies have shown the benefits of intravenous glutamine with total parenteral nutrition in severe acute pancreatitis. The aim of this study is to determine the effects of intravenous glutamine in severe acute pancreatitis patients who are being fed through a nasojejunal feeding tube (a tube which goes from the nose to the small intestine).

### Who can participate?

Patients aged over 18 with severe acute pancreatitis.

### What does the study involve?

Participants are randomly allocated into two groups. In the glutamine group, participants receive a glutamine drip for 7 days. In the control group, participants get a drip of other amino acids of the same quantity for the same period. Both groups are otherwise treated in the same way, including early nasojejunal feeding. During their hospital stay, we record the frequency of infectious complications, organ failures, the rate of operations and radiological interventions, length of hospital stay and death rate.

### What are the possible benefits and risks of participating?

Glutamine drips are frequently used in intensive care, but are not so widespread in surgery, probably because other combined amino acid solutions are well-known and cheaper. It has no side effects as it is an amino acid normally found in the human body.

### Where is the study run from?

Petz Aladar County Teaching Hospital, Győr, Hungary.

### When is the study starting and how long is it expected to run for?

September 2008 to September 2012.

Who is funding the study?  
Mulartz Henrik Surgical Foundation (Hungary).

Who is the main contact?  
Nora Hajdu  
kajda@freemail.hu

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Nora Hajdu

**Contact details**  
The Mulartz Henrik Surgical Foundation  
Vasvari P. u. 2.  
Gyor  
Hungary  
9024  
-  
kajda@freemail.hu

## Additional identifiers

**Protocol serial number**  
G-2008/12

## Study information

**Scientific Title**  
A combination of intravenous glutamine-dipeptide and early nasojejunal nutrition in severe acute pancreatitis: a prospective randomized double-blind clinical trial

**Study objectives**  
Intravenous glutamine may be beneficial with early nasojejunal feeding in prevention of septic complications in severe acute pancreatitis.

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**  
Regional Science and Research Ethics Committee of Petz Aladar Teaching Hospital, 04/09/2008, ref: 76-1-28/2008

**Study design**  
Single-center prospective randomized double-blind clinical trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Severe acute pancreatitis

**Interventions**

The trial will involve about 60 patients, admitted to the surgical department of Petz Aladar County Teaching Hospital, Győr, Hungary. Participants will receive the same treatment according to local guidelines, including early nasojejunal nutrition and radiological and surgical interventions, if necessary, except for the supplementation of amino acids. Glutamine group will get glutamine, while control group will get a combination of other amino acids usually used in clinical practice as an intravenous supplementation. Participation will start on admission and end at dismissal from hospital.

**Intervention Type**

Supplement

**Primary outcome(s)**

1. Rate of pancreas-specific infectious complications
2. Organ failures

**Key secondary outcome(s)**

1. Comparison of radiological and surgical interventions
2. Length of hospital stay
3. Mortality rate

**Completion date**

04/09/2012

**Eligibility****Key inclusion criteria**

1. Patients admitted with severe acute pancreatitis
2. Glasgow score at least 3 or CRP level over 150 mg/ml

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

## **Key exclusion criteria**

1. Age under 18 years
2. Pregnancy
3. Patients with complaints starting before the last 48 hours of admission
4. Death within 48 hours of admission

## **Date of first enrolment**

04/09/2008

## **Date of final enrolment**

04/09/2012

## **Locations**

### **Countries of recruitment**

Hungary

### **Study participating centre**

**The Mulartz Henrik Surgical Foundation**

Gyor

Hungary

9024

## **Sponsor information**

### **Organisation**

The Mulartz Henrik Surgical Foundation (Hungary)

## **Funder(s)**

### **Funder type**

Research organisation

### **Funder Name**

The Mulartz Henrik Surgical Foundation (Hungary)

## **Results and Publications**

Individual participant data (IPD) sharing plan

**IPD sharing plan summary**

Not provided at time of registration