Procalcitonin for discontinuation of antibiotic therapy in clinically diagnosed Ventilator Associated Pneumonia

Submission date 11/08/2006	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date 01/09/2006	Overall study status Completed	 [] Statistical analysis plan [X] Results
Last Edited 05/10/2011	Condition category Respiratory	[_] Individual participant dat

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s) Scientific

Contact name Prof Michael Tamm

Contact details

University Hospital Basel Clinic of Pneumology Petersgraben 4 Basel Switzerland 4031 + 41 61 265 5184 mtamm@uhbs.ch

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

Acronym

The ProVAP pilot study

Study objectives

We hypothesise that a Procalcitonin (ProCT) guided approach will increase the number of antibiotic-free days (for Ventilator Associated Pneumonia [VAP]) alive at 28 days by one third without compromising clinical outcomes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Committee for the Protection of Human Subjects in Research (FWA #00004009, docket #H-11990), EKBB Switzerland

Study design

Prospective multicentre randomised controlled trial.

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Ventilator associated pneumonia

Interventions

In this study, all patients, irrespective of the randomisation group, will be treated with antibiotics for 72 hours.

1. In patients randomly assigned to the standard therapy group, antibiotics will be prescribed and discontinued based on the clinical stability, radiologic and laboratory findings as routinely performed in the treating facility. Serum and/or plasma samples for ProCT will be collected daily, and the treating physician will be blinded to the results of the ProCT level. 2. In patients randomly assigned to the ProCT group, the decision to discontinue antibiotic therapy will also be based on the clinical stability, radiologic and laboratory findings. However, in this group, a further assessment of the probability of bacterial infection using ProCT levels will be available. Antibiotic discontinuation will be recommended according to serum ProCT concentrations as follows:

2.1. strongly encouraged if less than 0.25 ug/L

2.2. encouraged if less than 0.5 ng/ml or a decrease more than or equal to 80% as compared to day zero values (or previous values)

2.3. discouraged if more than or equal to 0.5 ng/ml or a decrease less than or equal to 80% as compared to day zero values (or previous values)

2.4. strongly discouraged if more than or equal to 1 ug/L

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Number of antibiotic free-days (for VAP) alive within 28 days of clinically suspicion of VAP.

Secondary outcome measures

1. Clinical deterioration (defined as an increase in Clinical Pulmonary Infection Score [CPIS] of more than two points)

2. Microbiologically documented pulmonary infection recurrence

3. The evolution of signs and symptoms potentially linked to pulmonary infection (fever,

leukocyte counts, partial pressure of oxygen in arterial blood [PaO2]/fraction of inspired oxygen [FiO2], and radiological infiltrates)

4. Number of mechanical ventilation-free days at 28 days

5. The length of stay in the ICU within 30 days

6. In-hospital mortality up to 30 days

7. Mortality at 30 days

8. Percentage of patients in the ProCT group for whom treatment recommendations are followed

9. Correlation of other biomarkers and clinical course

Overall study start date

01/07/2006

Completion date

31/12/2008

Eligibility

Key inclusion criteria

1. Intensive Care Unit (ICU) patients who are intubated and have been mechanically ventilated for at least 48 hours

2. 18 years of age and older

3. Clinical suspicion of VAP based on clinical and radiological criteria (new or progressive radiographic infiltrate) plus at least two of three clinical features:

3.1. fever greater than 38°C3.2. leukocytosis or leucopenia3.3. purulent tracheal secretions

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex Both

Target number of participants 100

Key exclusion criteria

1. Pregnancy

2. Patients with coexisting, documented extrapulmonary infection diagnosed between days one and three that requires antibiotic therapy longer than three days

3. Previous long-term corticosteroid therapy (more than or equal to 0.5 mg/kg per day of prednisolone or equivalent for more than one month)

4. Severe immunosuppression (solid organ transplantation or stem cell transplant recipients, known Human Immunodeficiency Virus [HIV] infection, neutropenic patients and patients after chemotherapy)

Date of first enrolment

01/07/2006

Date of final enrolment 31/12/2008

Locations

Countries of recruitment Switzerland

Study participating centre University Hospital Basel Basel Switzerland 4031

Sponsor information

Organisation University Hospital Basel (Switzerland)

Sponsor details

c/o Professor Michael Tamm University Hospital Basel Clinic of Pneumology Petersgraben 4 Basel Switzerland 4031 + 41 61 261 5184 mtamm@uhbs.ch

Sponsor type University/education

ROR https://ror.org/04k51q396

Funder(s)

Funder type University/education

Funder Name University Hospital Basel (Switzerland)

Results and Publications

Publication and dissemination plan Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary Not provided at time of registration

Study outputs

Output type

Details Date created

Date added

Peer reviewed?

Patient-facing?

Results article	results	01/12/2009	Yes	No
Results article	results	01/03/2011	Yes	No
Results article	results	01/10/2011	Yes	No