

Sevoflurane inhalational introduction in the elderly: a comparison with intravenous induction

Submission date 12/09/2003	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 12/09/2003	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 11/10/2016	Condition category Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr MH Nathanson

Contact details
Department of Anaesthesia
C Floor, University Hospital
Nottingham
United Kingdom
NG7 2UH

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0192080350

Study information

Scientific Title

Sevoflurane inhalational introduction in the elderly: a comparison with intravenous induction

Study objectives

The aim of the study is to compare the effects of an inhalational induction of anaesthesia with sevoflurane with an intravenous induction in elderly patients. In particular, we will assess the effects on cardiovascular parameters (heart rate, blood pressure), and cerebral blood flow velocity. We hypothesise that compared with an intravenous induction an inhalational induction will be more stable.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Anaesthesia

Interventions

Randomised controlled trial:

1. Inhalational induction of anaesthesia with sevoflurane
2. Intravenous induction of anaesthesia with sevoflurane

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Sevoflurane

Primary outcome measure

Change in blood pressure and cerebral flow velocity during induction of anaesthesia compared to baseline values.

Secondary outcome measures

Not provided at time of registration

Overall study start date

20/08/2001

Completion date

14/11/2002

Eligibility**Key inclusion criteria**

Total number of subjects = 40

Participant type(s)

Patient

Age group

Senior

Sex

Both

Target number of participants

40

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

20/08/2001

Date of final enrolment

14/11/2002

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre

University Hospital
Nottingham
United Kingdom
NG7 2UH

Sponsor information

Organisation

Department of Health (UK)

Sponsor details

Richmond House
79 Whitehall
London
United Kingdom
SW1A 2NL

Sponsor type

Government

Website

<http://www.doh.gov.uk>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Queens Medical Centre, Nottingham University Hospital NHS Trust (UK) - NHS R&D Support Funding

Funder Name

Abbott Laboratories (UK)

Alternative Name(s)

Abbott, Abbott U.S., Abbott Alkaloidal Company

Funding Body Type

Government organisation

Funding Body Subtype

For-profit companies (industry)

Location

United States of America

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration