

# The ketogenic diet in bipolar disorder

<b>Submission date</b> 28/02/2022	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 02/03/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 27/01/2025	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Bipolar disorder (BD) is a mental health condition that affects your moods, which can swing from one extreme to another. It is a major lifelong mental health condition that significantly impacts the quality of life of those who live with it. Our current understanding of how it develops is poor, and as such treatments are not always effective. There is some evidence that the ketogenic diet (KD), which is low in carbohydrate and high in fat, may be an effective treatment for BD, and some patients already follow this diet. To date there has been no clinical trial of the KD in BD. This study will pilot the introduction of a KD for 8 weeks in patients with BD. The aim is to find out how easy it is to do this, the main challenges, and how to address these. The acceptability of the intervention will be assessed, looking at how many people remain on the diet, what side effects they experience, and what support they require in sticking to the diet. This will provide information that will help the team design a future clinical trial.

### Who can participate?

Patients aged 18-70 years with bipolar disorder who have not have had any episodes of major depression, mania or hypomania in the preceding 3 months

### What does the study involve?

Participants will follow the diet for 8 weeks, with an additional 2 weeks to stop it gradually (unless they chose to continue). They will have regular support from a dietitian and psychiatrist. They will have tests performed before starting the diet and after 8 weeks at a clinical research facility. These will include questionnaires relating to their mental health, general health, finances and education and employment. They will have physical measurements taken to include blood pressure, weight and height. Blood tests and brain imaging will be performed to look at specific markers which may change on the KD. This will help with understanding how the diet might work. Participants will also monitor their ketone and glucose levels and mood state daily whilst on the KD, and wear a wrist device to record sleep and activity throughout the 10-week study period.

### What are the possible benefits and risks of participating?

Potential benefits to participants include learning more about their mental state over a period of time, as recorded in an app. They will also be able to better understand their sleep and activity pattern, as a summary of data from the actigraph device will be made available to them. For participants who are overweight, a potential benefit is weight loss, a well-known effect of

the ketogenic diet.

Possible risks to participants include discomfort from blood tests, or experiencing anxiety or claustrophobia when having an MRI brain scan. Following the ketogenic diet may be intrusive and impact on participants' quality of life as it will require a completely new set of dietary rules which may impact social situations. To help with this they will have regular support from a dietitian. There are potential side effects from commencing the ketogenic diet which commonly include fatigue, irritability, hunger and altered bowel habit. These are typically mild and can be managed with diet adjustments under the guidance of a dietitian.

Where is the study run from?

This study will be run from the clinical research facility at the Royal Infirmary of Edinburgh. It will be run by the University of Edinburgh with collaboration from NHS Lothian and the University of Glasgow (UK)

When is the study starting and how long is it expected to run for?

October 2021 to April 2023

Who is funding the study?

Baszucki Brain Research Fund (USA)

Who is the main contact?

Prof. Daniel Smith

d.smith@ed.ac.uk

## Contact information

**Type(s)**

Scientific

**Contact name**

Dr Nicole Needham

**ORCID ID**

<https://orcid.org/0000-0002-3610-1669>

**Contact details**

Centre for Clinical Brain Sciences, University of Edinburgh

Kennedy Tower, Royal Edinburgh Hospital

Morningside Park

Edinburgh

United Kingdom

EH10 5HF

+44 (0)131 5376531

nneedham@ed.ac.uk

**Type(s)**

Principal investigator

**Contact name**

Prof Daniel Smith

**ORCID ID**

<https://orcid.org/0000-0002-2267-1951>

**Contact details**

Centre for Clinical Brain Sciences,  
University of Edinburgh  
Kennedy Tower,  
Royal Edinburgh Hospital  
Morningside Park  
Edinburgh  
United Kingdom  
EH10 5HF  
+44 (0)131 537 6509  
d.smith@ed.ac.uk

**Type(s)**

Principal investigator

**Contact name**

Prof Harry Campbell

**ORCID ID**

<https://orcid.org/0000-0002-6169-6262>

**Contact details**

Centre for Global Health  
Usher Institute  
University of Edinburgh  
Teviot Place  
Edinburgh  
United Kingdom  
EH8 9AG  
+44 (0)7876314010  
harry.campbell@ed.ac.uk

**Additional identifiers****Clinical Trials Information System (CTIS)**

Nil known

**Integrated Research Application System (IRAS)**

306939

**ClinicalTrials.gov (NCT)**

Nil known

**Protocol serial number**

Sponsor AC21156, IRAS 306939

**Study information**

**Scientific Title**

A pilot study of the ketogenic diet in bipolar disorder

**Acronym**

KETOBD

**Study objectives**

The researchers have described in detail a novel hypothesis on the mechanism by which hyperinsulinaemia disrupts the production of energy by mitochondria in the brain, resulting in increased levels of specific biomarkers, including lactate and pyruvate. Higher levels of these metabolites have been described in bipolar disorder (BD) compared to the general population and may be linked to mood instability. The ketogenic diet (KD) can provide an alternative energy source to glucose (ketones) that can bypass this abnormal processing of energy and allow for more stable levels of energy production.

This study aims to pilot the use of the KD in people with BD for 8 weeks. This will help us to understand the feasibility and acceptability of the different aspects of compliance and monitoring that will be required in a future trial. There is no specific hypothesis as this is a pilot study without a primary outcome. In a future trial, however, the researchers will hypothesise that the use of the KD in BD will result in an improvement in mental state and a reduction in the levels of certain metabolites measured through metabolomics and magnetic resonance (MR) spectroscopy.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Approved 24/01/2022, South East Scotland Research Ethics Committee 02 (2nd Floor, Waverley Gate, 2-4 Waterloo Place, Edinburgh, EH1 3EG, UK; +44 (0)131 536 9000; sandra.wyllie@nhslothian.scot.nhs.uk), REC ref 22/SS/0007

**Study design**

Single-centre non-randomized interventional pilot study with no control arm

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Bipolar disorder

**Interventions**

All participants will be trialled on a low-carbohydrate, high-fat ketogenic diet for a period of 8 weeks, the first two of which will be an adaptation period. Support will be via dietary supervision to check adherence to prescribed diets, problem solve and identify any adverse effects, which will mainly occur over the phone or via email. Dietitians will provide guidance and practical support around establishing and characterising the diet, using recipe cards and planning, dietary booklets and providing advice on common problems.

Prior to commencing the ketogenic diet, participants will be required to attend initial baseline assessments, including mental health and health economics questionnaires, physical parameters (height, weight, blood pressure), blood tests (for biochemistry and metabolomics) and an MRI brain scan. They will also be taught how to measure their ketone and glucose measurements and monitor their mood on a daily basis and be provided with an actigraph device to wear throughout the study.

## **Intervention Type**

Behavioural

## **Primary outcome(s)**

The feasibility and acceptability of the ketogenic diet as an intervention for bipolar disorder for a future trial. Data will be collected on a range of parameters to understand how to make all aspects of the study workable for a subsequent trial:

1. Timeline estimates for future trial recruitment by measuring the number of interested participants and fully recruited participants from each avenue of recruitment during the estimated 4-month recruitment period
2. Participant level of compliance with dietary interventions measured using fidelity checklists at weeks 0, 2, 4, 6, and 8, and daily ketone measurements during the 8-week intervention period
3. Participant level of compliance with study assessments as recorded on the case report form at baseline and week 8
4. Participant level of compliance with continuous study assessments measured using data from the study devices and apps (actigraph device, ketomojo device and app, and ilumivu app) during the 10-week study period
5. Quality of life measured using the EQ5D-5L at baseline and week 8
6. Productivity and activity impairment measured using the Work Productivity and Activity Impairment Questionnaire at baseline and week 8
7. Participant resource use measured using a Within Trial Resource Use Questionnaire at baseline and week 8
8. Staff time and cost in delivering dietary interventions, measured by keeping contemporaneous records throughout the 10-week study period
9. Attrition rate measured using the number of participants who consent to participate who remain in the study until the end of follow-up at 10 weeks, to help inform estimation of sample size for a future trial
10. Acceptability and experience of the study participants evaluated via telephone interviews after the main study has ended as part of a process evaluation

## **Key secondary outcome(s)**

1. Mood stability measured by the Affective Lability Scale (ALS18) and Beck Depression Inventory (BDI) at baseline and at 8-week follow-up
2. Hypomania/mania symptoms measured using the Young Mania Rating Scale (YMRS) at baseline and 8-week follow up
3. Glucose/ketone ratio measured using ketomojo devices daily throughout the 10 week study period
4. Identification of specific metabolic changes in glucose, ketones and tricarboxylic acid (TCA) metabolites associated with the KD that are predicted by the study hypothesis, measured using serum and brain MRI measures at baseline and week 8
5. Mood, energy, speed of thought, impulsivity and anxiety measured using visual analogue scales from 1-100 at daily ecological momentary assessments during the 10-week study period
6. Episodes of bipolar depression and hypomania/mania occurring during the study period assessed by the MINI Neuropsychiatric assessment at baseline and week 8

7. Sleep duration and circadian activity/rest rhythmicity parameters assessed by actigraphs throughout the 10-week period (to include 2 weeks of stepping down the diet)

**Completion date**

08/04/2023

## Eligibility

**Key inclusion criteria**

Current participant inclusion criteria as of 29/04/2022:

1. Meet the Diagnostic and Statistical Manual of Mental Disorders (DSM-4) diagnostic criteria for bipolar disorder for at least 1 year, assessed using the MINI Neuropsychiatric Interview
2. Clinically stable and currently euthymic, defined as 3 months with no major mood episodes (major depression lasting at least 2 weeks or hypomania/mania lasting at least 1 week), assessed using the MINI Neuropsychiatric Interview
3. Aged 18 to 70 years
4. Able to provide informed consent to take part in the study
5. Able to speak, read and understand English to a level whereby they can understand the participant information sheet (PIS)
6. Currently living in Scotland

Previous participant inclusion criteria:

1. Meet the Diagnostic and Statistical Manual of Mental Disorders (DSM-4) diagnostic criteria for bipolar disorder for at least 1 year, assessed using the MINI Neuropsychiatric Interview
2. Clinically stable and currently euthymic, defined as 3 months with no major mood episodes (major depression lasting at least 2 weeks or hypomania/mania lasting at least 1 week), assessed using the MINI Neuropsychiatric Interview
3. Aged 18 to 70 years
4. Able to provide informed consent to take part in the study
5. Able to speak, read and understand English to a level whereby they can understand the participant information sheet (PIS)

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Mixed

**Sex**

All

**Total final enrolment**

27

**Key exclusion criteria**

Current participant exclusion criteria as of 29/04/2022:

1. Pregnancy or breastfeeding (or those planning to become pregnant within 3 months)

2. Active substance misuse with alcohol or illicit drugs
3. Use of the ketogenic diet in the previous 2 months
4. Currently following a vegan diet
5. Admission to hospital within the past 3 months
6. Current involvement in another research study
7. Inability to complete baseline assessments
8. Liver or kidney disease
9. Cardiovascular disease
10. Severe hyperlipidaemia
11. Type 1 diabetes
12. Insulin dependent type 2 diabetes
13. Currently training for or undertaking very high energy requirement activities (as judged by the study dietitian)

Previous participant exclusion criteria:

1. Pregnancy or breastfeeding (or those planning to become pregnant within 3 months)
2. Active substance misuse with alcohol or illicit drugs
3. Use of the ketogenic diet in the previous 2 months
4. Currently following a vegan diet
5. Admission to hospital within the past 3 months
6. Current involvement in another research study
7. Inability to complete baseline assessments
8. Liver or kidney disease
9. Cardiovascular disease
10. Severe hyperlipidaemia
11. Type 1 diabetes

**Date of first enrolment**

27/04/2022

**Date of final enrolment**

28/10/2022

## **Locations**

**Countries of recruitment**

United Kingdom

Scotland

**Study participating centre**

**NHS Lothian**

Waverley Gate

2-4 Waterloo Place

Edinburgh

United Kingdom

EH1 3EG

**Study participating centre**  
**University of Edinburgh**  
Old College  
South Bridge  
Edinburgh  
United Kingdom  
EH8 9YL

**Study participating centre**  
**University of Glasgow**  
University Avenue  
Glasgow  
United Kingdom  
G12 8QQ

## **Sponsor information**

**Organisation**  
University of Edinburgh

**ROR**  
<https://ror.org/01nrxf90>

**Organisation**  
NHS Lothian

**ROR**  
<https://ror.org/03q82t418>

## **Funder(s)**

**Funder type**  
Research organisation

**Funder Name**  
Baszucki Brain Research Fund

## **Results and Publications**

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to the small sample size and this being a pilot study with no primary outcome.

## IPD sharing plan summary

Not expected to be made available

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		10/10/2023	11/10/2023	Yes	No
<a href="#">HRA research summary</a>			20/09/2023	No	No
<a href="#">Other publications</a>	Process evaluation	21/01/2025	27/01/2025	Yes	No
<a href="#">Participant information sheet</a>	version 3	18/02/2022	29/04/2022	No	Yes
<a href="#">Participant information sheet</a>	version 1	14/04/2022	29/04/2022	No	Yes
<a href="#">Preprint results</a>		03/06/2023	05/06/2023	No	No
<a href="#">Protocol file</a>	version 4	14/04/2022	29/04/2022	No	No
<a href="#">Protocol file</a>	version 7	10/11/2022	26/05/2023	No	No
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes