

Shoulder Injection Trial - A comparison of injection of Tenoxicam with Depo-Medrone in shoulders with subacromial impingement syndrome

Submission date 30/09/2004	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 30/09/2004	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 13/04/2011	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Mr Howard Kwong

Contact details

Orthopaedic Department
Hospital of St Cross
Barby Road
Rugby
United Kingdom
CV22 5PX
+44 (0)1788 572831/545240
htkwong@hotmail.com

Additional identifiers

Protocol serial number

N0295132803

Study information

Scientific Title

Study objectives

Compare the effects of subacromial injection of Tenoxicam (NSAID) with Depo-medrone (steroid) in patients with impingement syndrome.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Added as of 27/03/2008:

1. Warwick Research Ethics Committee (REC) (ref: RE553)
2. Coventry REC (ref: 020/09/03)

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Musculoskeletal Diseases: Shoulder disorders

Interventions

Please note that, as of 27/03/2008, the end date of this trial was updated from 30/09/2004 to 17/07/2006 (date on which the last participant was recruited).

Interventions added as of 27/03/2008:

The participants were randomly allocated to the following two arms in equal numbers:

Arm 1: Subacromial injection of tenoxicam (NSAID) 20 mg single dose

Arm 2: Subacromial injection of depo-medrone (steroid) 40 mg single dose

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Primary outcome measures updated as of 27/03/2008:

Constant-Murley shoulder assessment score at baseline and 6 weeks

Primary outcome measures provided at time of registration:

Constant-Murley Shoulder assessment score and Disabilities of the Arm Shoulder and Hand (DASH) Questionnaire.

Key secondary outcome(s)

Added as of 27/03/2008:

1. Disabilities of the Arm, Shoulder and Hand (DASH) Questionnaire at baseline and 2, 4 and 6 weeks
2. Oxford shoulder score at baseline and 2, 4 and 6 weeks

Completion date

17/07/2006

Eligibility**Key inclusion criteria**

Added as of 27/03/2008:

1. Patients over the age of 18 years
2. Clinical diagnosis of subacromial impingement syndrome
3. Symptoms lasting longer than 3 months

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

Added as of 27/03/2008:

1. Evidence of other pathology causing shoulder pain
2. Injection in the same shoulder within the previous 6 months
3. Patients taking regular systemic NSAIDs or steroids or in whom those drugs were contraindicated
4. If their present shoulder condition was the subject of any legal proceedings or insurance claims
5. Pregnant and breast-feeding mothers

Date of first enrolment

15/02/2004

Date of final enrolment

17/07/2006

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre
Orthopaedic Department
Rugby
United Kingdom
CV22 5PX

Sponsor information

Organisation
Department of Health

Funder(s)

Funder type
Government

Funder Name
University Hospitals Coventry and Warwickshire NHS Trust

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2010		Yes	No