

Napping behaviour in sleep apnoea: a transcultural study in older people

Submission date 12/01/2012	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 28/02/2012	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 07/04/2016	Condition category Nervous System Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Obstructive sleep apnoea syndrome (OSAS) is a condition where the walls of the throat relax and narrow during sleep, interrupting normal breathing. These repeated sleep interruptions can make patients with OSAS feel very tired during the day, causing them to take frequent naps. OSAS can be treated with continuous positive airway pressure (CPAP). This involves wearing a mask that delivers a continuous supply of compressed air that prevents the throat from closing. The aim of this study is to examine the effect of CPAP on napping behaviour in OSAS patients in two different ethnic populations.

Who can participate?

Scottish and Taiwanese patients with OSAS, aged 60 and over, who nap at least 3 times per week with average nap duration of at least 30 minutes.

What does the study involve?

Patients are randomly allocated to one of two groups. One group is treated with CPAP and best supportive care for 12 weeks. The other group is treated with best supportive care for 6 weeks, then CPAP and best supportive care for another 6 weeks. Best supportive care includes normal medical treatment and information about weight loss, stopping smoking, drinking less alcohol, changing sleeping posture and sleep hygiene. Both groups undergo tests at the start of the study and at follow-up. The tests evaluate sleep behaviour, mood and anxiety, and quality of life. All participants undergo 2 weeks of actigraphy (sleep monitoring) before hospital visits. Sleep diaries are recorded over the study period.

What are the possible benefits and risks of participating?

All participants receive detailed advice on managing their OSAS with improved diet, exercise and regular sleep patterns. We may find during the tests that something needs further investigation or treatment that would not have otherwise been found. CPAP is generally very well tolerated. Sometimes it can cause nasal dryness/discomfort, nasal discharge, a dry mouth (and dribbling), facial discomfort and, rarely, ulceration due to poor mask fitting. These effects can be minimised by the use of humidification and careful mask fitting.

Where is the study run from?

1. Royal Infirmary of Edinburgh (UK)
2. Kaohsiung Medical University Hospital (Taiwan)

When is the study starting and how long is it expected to run for?

February 2012 and October 2012

Who is funding the study?

National Science Council (Taiwan)

Who is the main contact?

Dr Chung-Yao Hsu

Contact information

Type(s)

Scientific

Contact name

Dr Chung-Yao Hsu

Contact details

No.100, Tzyou 1st Road

Kaohsiung

Taiwan

80754

Type(s)

Scientific

Contact name

Dr Renata Riha

Contact details

51 Little France Crescent

Old Dalkeith Road

Edinburgh

United Kingdom

EH16 4SA

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Napping Behaviour in Sleep Apnoea (NBSA): a randomised transcultural study in older people

Acronym

NBSA

Study objectives

Napping behaviour in obstructive sleep apnoea (OSAS) patients may reflect a compensatory reaction to sleep deprivation or sleep interruption. If sleep deprivation or sleep interruption due to disturbed breathing improve after continuous positive airway pressure (CPAP) treatment, napping behaviour may decrease.

Ethics approval required

Old ethics approval format

Ethics approval(s)

South East Scotland Research Ethics Committee 01, 21/02/2012, ref: 12/SS/0017

Study design

Randomised single-blind parallel study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Obstructive sleep apnoea syndrome (OSAS)

Interventions

1. An early CPAP group: CPAP + best supportive care (BSC) for 12 weeks
2. A delayed CPAP group: BSC for 6 weeks initially then CPAP + BSC will be combined for another 6 weeks.

BSC includes normal medical treatment and information about weight reduction, smoking cessation, reducing alcohol consumption, change in sleeping posture and sleep hygiene.

Intervention Type

Device

Primary outcome(s)

Change in weekly nap frequency and duration every 6 weeks in both groups

Key secondary outcome(s)

1. CPAP compliance during CPAP intervention in both groups
 2. Change in Epworth Sleepiness Scale (ESS) and Pittsburgh Sleep Quality Index (PSQI) scores
 3. Change in RAND-36 scores
 4. Change in Hospital Anxiety and Depression Scale (HADS) scores
- Measured every 6 weeks in both groups
5. Change in reaction time of Psychomotor Vigilance Task (PVT) assessed post intervention

Completion date

31/10/2012

Eligibility

Key inclusion criteria

1. Aged 60 years and over
2. Have apnoea-hypopnoea index (AHI) ≥ 15
3. Nap at least 3 times per week with average nap duration of at least 30 minutes

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Key exclusion criteria

1. Have previous exposure to continuous positive airway pressure (CPAP) therapy
2. Are unable to give written informed consent or comply with the protocol
3. Are on central nervous system (CNS)-active drug treatment (e.g., hypnotic and antiepileptic drugs) that causes significant daytime sleepiness or cognitive impairment
4. Have an unstable medical or psychiatric illness
5. Have moderate to severe respiratory disease that affects activities of daily living or awake SpO₂ less than 92%

Date of first enrolment

15/02/2012

Date of final enrolment

31/10/2012

Locations

Countries of recruitment

United Kingdom

Taiwan

Study participating centre

No.100, Tzyou 1st Road

Kaohsiung

Taiwan

80754

Sponsor information

Organisation

Kaohsiung Medical University Hospital (Taiwan)

Organisation

Department of International Cooperation (Taiwan)

Organisation

Kaohsiung Medical University Chung-Ho Memorial Hospital

ROR

<https://ror.org/02xmkec90>

Funder(s)

Funder type

Government

Funder Name

National Science Council (NSC 99-2911-I-037-004)

Alternative Name(s)

National Science Council, Taiwan, National Science Council of Taiwan, NSC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Taiwan

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration