

# Hypercoagulation effect evaluation and deep vein thrombosis preventions after laparoscopic surgery.

<b>Submission date</b> 02/11/2014	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 29/12/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 29/12/2014	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Venous thromboembolism (VTE) is a serious, potentially fatal, medical condition where people develop a blood clot in a vein. People who have just had major surgery are at particular risk of VTE; between 20-30% of VTE patients have just had general surgery. About 5.5% of VTE patients have undergone laparoscopic fundoplication, a "keyhole" surgery for treating gastro-oesophageal reflux disease. Here, we want to test whether treating patients with Bemiparin (Zibor) before patients have their laparoscopic fundoplication surgery reduces the risk of deep vein thrombosis (DVT), a blood clot in a leg vein. We will be investigating two different treatment regimens to see which one works best.

### Who can participate?

Adult patients between 18-75 years undergoing laparoscopic fundoplication surgery.

### What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 receive one dose (2500 TV 0,2 ml.) of Bemiparin and intraoperative intermittent pneumatic compression (IPC – a therapy whereby an inflatable device squeezes fluids, such as blood, out of the area where it is applied) one hour before their operation. Those in group 2 receive 3 doses of Bemiparin, one 12 hours before the operation, one 6 hours after the operation and a final dose 30 hours after the operation. IPC is also applied during the surgery. After the operation, the blood is tested for certain components (markers) that would suggest a blood clot is forming (thrombogenesis) and other components that would suggest that the blood is not able to clot (hypocoagulation). These are measured 1 hour before, during, just after and then 2 days after the operation. Deep vein imaging methods are also used to check for DVT before the operation, 2 days afterwards and finally 28 days after the surgery.

### What are the possible benefits and risks of participating?

There are no additional risks to taking part in the study.

Where is the study run from?

The Hospital of Lithuanian University of Health Sciences (LUHS) (Lithuania)

When is the study starting and how long is it expected to run for?

January 2014 to January 2017

Who is funding the study?

The Lithuanian University of Health Sciences (LUHS) (Lithuania)

Who is the main contact?

Dr. Indre Zostautiene

indrajoss@gmail.com

## Contact information

**Type(s)**

Scientific

**Contact name**

Dr Indre Zostautiene

**Contact details**

S. Ciurlionienes-Kymantaites str. 7

Kaunas

Lithuania

LT-46320

-

indrajoss@gmail.com

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

N/A

## Study information

**Scientific Title**

Hypercoagulation effect evaluation and deep vein thrombosis preventions after laparoscopic surgery: a prospective randomized double blinded trial.

**Acronym**

N/A

**Study objectives**

A combination of one dose low-molecular-weight heparin (LMWH) 1h before laparoscopic surgery with intraoperative intermittent pneumatic compression (IPC) causes better hypocoagulation effect than LMWH administered 12h before surgery with intraoperative IPC.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Regional Biomedical Research Ethics Committees in Lithuania: in Kaunas region (based at the Lithuanian University of Health Sciences), 08/10/2014, ref. Nr. BE-2-13.

### **Study design**

Prospective randomized double blinded trial.

### **Primary study design**

Interventional

### **Secondary study design**

Randomised controlled trial

### **Study setting(s)**

Hospital

### **Study type(s)**

Treatment

### **Participant information sheet**

Not available in web format, please use the contact details below to request a patient information sheet

### **Health condition(s) or problem(s) studied**

Patients with hiatal hernia diagnosis and condition after laparoscopic fundoplications. The study domain is to evaluate the hypocoagulation effect and rate of deep vein thrombosis complications of intermittent pneumatic compression (IPC) combination of low molecular weight heparin (LMWH) after laparoscopic fundoplication

### **Interventions**

The participants were randomly allocated into one of two groups:

Group 1 received subcutaneous 2500 TV 0,2 ml LMWH Bemiparin (Zibor) - one dose (1h before the operation)

Group 2 received three doses of 2500 TV 0,2 ml LMWH Bemiparin (Zibor) - (12h before the operation, 6h and 30h after the operation).

### **Blood sampling:**

Venous blood samples were taken (obtained) from the forearm vein four times:

1. 1h. before the operation (before LMWH injection)
2. 1h after the introduction of the laparoscope
3. After extubation
4. 2 days after the operation

## Imaging methods:

1. Duplex examination - before the operation, two days after the operation and 28 days after the operation
2. Spiral CT venography two days after the operation.

## Intervention Type

### Phase

Not Applicable

### Primary outcome measure

1. F1+2 and TAT concentrations in plasma samples were measured by solid phase sandwich ELISA method (Enzygnost® F1+2 and Enzygnost® TAT microenzyme immunoassay, Dade Behring, Germany).
2. FTFPI was performed by an ELISA technique (Diagnostika Stago, France) according to the manufacturers instructions.
3. Thromboelastograma - TEG measures the physical properties of the clot in whole blood via a pin suspended in a cup (heated to 37C) from a torsion wire connected with a mechanical electrical transducer; koalin is used as activator.

### Secondary outcome measures

Ultrasound examination - Toshiba Aplio SSA 770A - compression ultrasound (B-mode imaging only), duplex ultrasound (B-mode imaging and Doppler waveform analysis), and color Doppler imaging alone. Different lower extremity veins are best evaluated with different techniques: compression ultrasound is typically performed on the proximal deep veins - the common femoral, femoral, and popliteal veins, whereas a combination of duplex ultrasound and color Doppler imaging is more often used to interrogate the calf and iliac veins.

Spiral CT venography - GE Light Speed Pro (320 slice) - using 150 mL of IV contrast -

4 mL/sec, with a delay three minutes after the start of injection, 5-mm thick axial CT venograms are acquired from the ankles to the mid abdomen.

### Overall study start date

01/01/2014

### Completion date

01/01/2017

## Eligibility

### Key inclusion criteria

1. Patients undergoing elective laparoscopic funduplications because of gastroesophageal reflux disease, caused by hiatus hernia.
2. The patients in the two groups are similar in terms of age (18 - 75 years), weight, height, gender, duration of surgery and ASA score.

### Participant type(s)

Patient

### Age group

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

100-150

**Key exclusion criteria**

1. Patients with coagulopathies disorders
2. Usage of prescription drug modifying cardiovascular response
3. Participants with systemics nervous system disease
4. Pregnant women
5. Patients, who refused to participate in the study

**Date of first enrolment**

01/01/2014

**Date of final enrolment**

01/01/2017

## **Locations**

**Countries of recruitment**

Lithuania

**Study participating centre**

**S. Ciurlionienes-Kymantaites str. 7**

Kaunas

Lithuania

LT-46320

## **Sponsor information**

**Organisation**

The Lithuanian University of Health Sciences (LUHS) (Lithuania)

**Sponsor details**

Eiveniu str. 2

Kaunas

Lithuania

LT-50009

-  
rastine@kaunoklinikos.lt

**Sponsor type**

University/education

**ROR**

<https://ror.org/0069bkg23>

## **Funder(s)**

**Funder type**

University/education

**Funder Name**

The Lithuanian University of Health Sciences (LUHS) (Lithuania)

## **Results and Publications**

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration