

A study to investigate the feasibility of smartphone-based self-monitoring to characterise cognitive and neurological impairment in participants with multiple sclerosis (Floodlight MS_More Active)

Submission date 02/03/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 11/03/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 20/11/2024	Condition category Nervous System Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Multiple sclerosis (MS) is a chronic (long-lasting), inflammatory, demyelinating (causing damage to the myelin sheath (protective covering) surrounding nerve fibers in the brain), and degenerative disease (characterized by progressive deterioration and loss of function) of the central nervous system (CNS). Symptoms of MS are very different between people but can commonly include tiredness, vision problems, and problems with walking or balance. The main purpose of this study is to analyse how a combination of the information collected during the participant's routine clinic visits (including MS history and magnetic resonance imaging (MRI) scans), along with other information about participants' movement, brain function, hand function, speech and quality of life collected using smartphone applications (App), can be used to improve the way change in function is monitored in people with MS. The study will mainly focus on:

1. The ease of using Floodlight MS (FL MS) cognitive test to identify cognitive (mental processes like thinking, learning, etc.) impairment in participants with MS (pwMS)
2. The ease of using FL MS cognitive test to characterise cognitive trajectories in pwMS
3. Exploring the effect of different frequencies of administration for the FL MS cognitive test on learning and practice effects in pwMS
4. Evaluating the ease of using FL MS gait and hand function tests to characterise neurological impairment and trajectories in pwMS
5. Evaluating the ease of using FL MS digital outcomes to predict future disability progression in pwMS
6. Assessing participant adherence to self-monitoring tasks
7. Assessing participant's user experience with regular digital outcome measures with FL MS
8. Assessing the learning effect for each measure of FL MS in pwMS
9. Assessing the ease of using self-monitoring tasks to characterize clinical relapses in pwMS

Who can participate?

Patients with MS who are over 18 years of age and have a personal smartphone

What does the study involve?

Participants will be asked to be a part of this study for at least 2 years, and information about participants' MS may be accessed for up to 3 years. The participants taking part in this study will also have to agree to participate in the MSBase Registry study. MSBase is a large, ongoing, long-term study that collects information from over 70,000 people with MS in over 34 countries. The participants will have to perform the following tests while they participate in the study:

- FL MS smartphone application tests: This test includes-

1. Completing a survey on how the participant is feeling and assessing their MS symptoms
2. Drawing a shape on the smartphone screen
3. Pinching a shape on the smartphone screen
4. Walking for 2 minutes and as fast as possible but safely (Two-minute walk test)
5. Walking safely and performing at least 5 successive U-turns (U-Turn test)
6. Cognitive test (measuring how fast participants are thinking or processing information)

These tests will take 5-10 minutes to complete

- Expanded Disability Status Scale (EDSS) exams: It is based on a standard neurological examination used for the determination of disability in a participant. The scores of the neurological examination are then used in conjunction with observations and information concerning ambulation (ability to walk) and the use of assistive devices by a participant to determine the EDSS score.

- MSReactor tests: This test is designed to measure changes in the participant's reaction time and memory (cognition), and so may provide slightly different information than the FL MS application tests. At the end of the MSReactor tests, participants will be presented with some questionnaires that measure the quality of life, work productivity, worry, and depression. This will take 10-15 minutes to complete

- Redenlab tests: This measures changes in the participant's speech and consists of five reading aloud and speaking tasks that will take less than 5 minutes to complete.

- Audio Recorded Cognitive Screen (ARCS) test: Participants will be asked to sit in a quiet area with no distractions, unsupervised, and listen to an audio file on headphones and record their responses in the booklet provided. The test will be completed in 35-40 minutes, after which it is scored by a trained person.

The study will consist of 4 periods:

1. Screening Period: Participants will be screened for enrollment in this study by the Investigator.

2. Baseline Visit: The first study visit, during which the investigator/coordinator will assist participants in installing and registering the FL MS, Redenlab, and MSReactor App on their smartphones. This process may be completed through a virtual visit using available video-calling software. After installation of these apps, participants will be asked to complete the following tests:

- FL MS smartphone application tests: Will be performed in-person during the baseline visit
- Redenlab tests: Will be performed in-person during the baseline visit
- Participant's most recent (within 6 months) MRI scans will be collected

3. Familiarisation Period: Following the baseline visit, participants will be asked to perform the following test during the 6-weeks familiarisation period:

- FL MS tests 1 to 5 using the FL MS App: Every other day or at least 3 times per week (18 to 21 times in total). Participants will also need to perform the cognitive test (test 6 in the list above) weekly using the FL MS App
- MSReactor tests: Once per week (or at a minimum, 4 times in total)

- Redenlab tests: Once a week (or at a minimum 4 tests in total)

The purpose of the familiarisation period is for the participants to learn how to use the Apps and to ensure that they are conducting the tests as per the instructions. All the tests will be performed by the participants at their homes.

4. Observational Period: After the familiarisation period, participants will enter the observational period for approximately 2 years. Participants will continue to have the usual routine visits with their neurologist, for approximately 6 months, and will need to attend a study-only visit once during Month 12 and once during Month 24. Participants will have to perform the following tests during the observational period:

- FL MS App Tests: Tests 1-5: For the first 2 weeks, will be performed every other day except the cognitive test (to be performed once weekly) to establish baseline data. After 2 weeks of the observation period, FL MS tests 1-5 will be performed at least twice a month; and test 6 once a week, once in two weeks, or once a month (depending on the participant's study group) for the remainder of the 2-year study period.

- EDSS: A telephonic EDSS (tele-EDS) assessment conducted over the telephone within a ± 2 -week time around the first two weeks of FL MS testing (from Day 29 to Day 70) in the observational period to ensure that a baseline tele-EDSS measurement is available for all participants

- MSReactor tests: At least once a month for the remainder of the 2-year study period. Participants who have done MSReactor tests on a tablet or desktop before may be approached to perform the test twice, once on a tablet computer and once on the smartphone

- Redenlab tests: At least once a month for the remainder of the 2-year study period

- ARCS test: Once a year at a study clinic visit i.e., once during Month 12 and once during Month 24

- Participant's MRI scans performed as part of the routine clinical care will be collected during the 2-year study period

Participants will be divided into three cohorts based on the frequency at which they will perform the FL MS cognitive test. All other tests mentioned above will be performed at the same frequency by all the participants.

- Participants in Cohort 1 will perform the FL MS cognitive test once every week.

- Participants in Cohort 2 will perform the FL MS cognitive test once in two weeks

- Participants in Cohort 3 will perform the FL MS cognitive test once a month

For participants who were a part of the Floodlight Open study (Study MN39878):

1. Participants will complete the final test for the Floodlight Open study at the first visit for this study (Baseline visit).

2. Participants will uninstall the Floodlight Open study application and install the new FL MS software application to be used in this study

What are the possible benefits and risks of participating?

Participants will receive no clear benefit from participating in this research. The results of this study may provide valuable information to the researchers and doctors that may help improve the treatment or care of people with MS as a group. There may be unknown or unforeseen risks, including privacy risks, associated with participating in this study.

Where is the study run from?

F. Hoffman La Roche (Switzerland)

When is the study starting and how long is it expected to run for?

January 2022 to July 2024

Who is funding the study?
F. Hoffman La Roche (Switzerland)

Who is the main contact?
global.trial_information@roche.com

Study website

<https://forpatients.roche.com/en/trials/autoimmune-disorder/multiple-sclerosis/a-prospective-investigation-of-the-feasibility-of-smartphone-bas.html>

Contact information

Type(s)

Public, Scientific

Contact name

Dr Clinical Trials

Contact details

Grenzacherstrasse 124
Basel
Switzerland
4070
+41 616878333
global.trial_information@roche.com

Type(s)

Principal Investigator

Contact name

Prof Anneke van der Walt

Contact details

Monash University
553 St Kilda Rd
Melbourne
Australia
VIC 3004
+61 3 9903 9300
med-clinicaltrials@monash.edu

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

MN42134

Study information

Scientific Title

A prospective investigation of the feasibility of smartphone-based self-monitoring to characterize cognitive and neurological impairment in patients with multiple sclerosis (Floodlight MS_More Active)

Study objectives

The primary purpose of this study is to evaluate the feasibility of using the Floodlight Multiple Sclerosis (FL MS) Cognitive Test to characterize cognitive impairment in participants with MS (pwMS).

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 20/07/2022, The Alfred (PO Box 315, , Prahran, VIC 3181, Australia; +61 (03) 9076 2000; research@alfred.org.au), ref: Project 612/21

Study design

Prospective non-interventional primary data collection study

Primary study design

Observational

Secondary study design

Longitudinal study

Study setting(s)

Other

Study type(s)

Other

Participant information sheet

No participant information sheet available

Health condition(s) or problem(s) studied

Multiple sclerosis

Interventions

- Cohort 1: Participants in Cohort 1 will perform the FL MS Cognitive Test once every week and all other FL MS tests every 2 weeks, MSReactor tests, and Redenlab tests at least once a month and Audio Recorded Cognitive Screen (ARCS) test once a year up to Month 24.
- Cohort 2: Participants in Cohort 2 will perform the FL MS Cognitive Test and all other FL MS

tests every 2 weeks, MSReactor tests, and Redenlab tests at least once a month, and ARCS test once a year up to Month 24.

- Cohort 3: Participants in Cohort 3 will perform the FL MS Cognitive Test every 4 weeks and all other FL MS tests every 2 weeks, MSReactor tests, and Redenlab tests at least once a month, and ARCS test once a year up to Month 24.

Intervention Type

Other

Primary outcome measure

1. Cross-sectional correlation between FL MS Cognitive Test outcome and ARCS total outcome measured using Spearman's Rank-order Correlation at baseline, Month 12, and Month 24

Secondary outcome measures

Current secondary outcome measures as of 21/12/2023:

1. Cross-sectional correlation between composite digital cognition outcome with FL MS and ARCS total score measured using Spearman's Rank-order Correlation at baseline, Month 12, and Month 24
2. Cross-sectional correlation between FL MS digital cognition outcome and Expanded Disability Status Scale (EDSS) measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
3. Cross-sectional correlation between digital outcome measures with FL MS and Redenlab score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
4. Cross-sectional correlation between digital outcome measures with FL MS and MSReactor Cognitive score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
5. Cross-sectional correlation between FL MS digital outcome measures (MS gait and hand function test) and Patient-reported outcome measures (PROMs)- Multiple Sclerosis International Quality of Life (MusiQoL) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24
6. Cross-sectional correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs- Work Productivity and Activity Impairment Questionnaire: General Health (WPAI:GH) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24
7. Cross-sectional correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs- Patient Health Questionnaire-9 (PHQ-9) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24
8. Cross-sectional correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs- Penn State Worry Questionnaire (PSWQ-15) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24
9. Cross-sectional correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs- Patient Global Impression of Severity Questionnaire (PGIS) assessed using Spearman's Rank-order Correlation at Months 12 and 24
10. Cross-sectional correlation between digital outcome measures with FL MS and MRI parameters assessed using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
11. Longitudinal correlation between composite digital cognition outcome with FL MS and ARCS total score measured using Spearman's Rank-order Correlation at baseline, Month 12, and Month 24
12. Longitudinal correlation between digital cognition outcome measures with FL MS and EDSS measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
13. Longitudinal correlation between digital cognition outcome measures with FL MS and Redenlab score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

14. Longitudinal correlation between digital cognition outcome measures with FL MS and MSReactor Cognitive score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
15. Longitudinal correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs (MusIQoL) assessed using Spearman's Rank-order Correlation over 24 months
16. Longitudinal correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs (WPAI:GH) assessed using Spearman's Rank-order Correlation over 24 months
17. Longitudinal correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs (PHQ-9) assessed using Spearman's Rank-order Correlation over 24 months
18. Longitudinal correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs (PSWQ-15) assessed using Spearman's Rank-order Correlation over 24 months
19. Longitudinal correlation between FL MS digital outcome measures (MS gait and hand function test) and PROMs (PGIC) assessed using Spearman's Rank-order Correlation at Months 12 and 24
20. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs (PGIS) assessed using Spearman's Rank-order Correlation over 24 months
21. Longitudinal correlation between digital outcome measures with FL MS and MRI parameters assessed using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
22. Correlation between FL MS Digital Outcomes and subsequent changes in EDSS assessed using Spearman's Rank-order correlation calculated from baseline to Month 12 and Month 12 to 24
23. Cognitive phenotypes identified using a joint latent profile analysis (LPA) on the baseline FL MS Cognitive Test outcome, ARCS Global Score, and MSReactor Cognitive score at baseline (Day 1)
24. Difference in trajectories of FL MS Cognitive Test outcome between participants performing cognitive test every week versus every 2 weeks versus every 4 weeks, measured using a Piecewise Linear Regression Method over the first 6 and 12 months
25. Participant engagement patterns in the first 3 months of usage of FL MS
26. Participant engagement patterns in the first 24 months of usage of FL MS
27. Factors influencing participants engagement (demographics and disease factors) on FL MS, Redenlab, and MSReactor Apps usage recorded from baseline to Month 24
28. Factors influencing participant's adherence (physician-driven factors i.e., phone calls) to self-monitoring recorded from baseline to Month 24
29. In-app FL MS Post-appointment Questionnaire assessed at Month 12 and Month 24
30. FL MS usability score from mHealth Usability Questionnaire assessed at Month 12 and Month 24
31. Number of assessments or familiarisation period duration required to minimize the impact of learning effect assessed using FL MS to ensure stable FL MS baseline performance measured from baseline to Week 6
32. Changes in digital outcome measures with FL MS and the presence of protocol-defined relapse measured from baseline to Month 24
33. Longitudinal correlation between composite digital outcome and ARCS total score measured using Spearman's Rank-order Correlation over 24 months
34. Differences between participants with different learning patterns on FL MS Cognitive Test outcome identified using practice effect trajectories from baseline to Months 6, or 12
35. Differences between participants with different learning patterns on ARCS score identified using practice effect trajectories from baseline to Months 6, or 12
36. Differences between participants with different learning patterns on EDSS scale score

identified using practice effect trajectories from baseline to Month 6, or 12

37. Differences between participants with different learning patterns on MRI outcome measures identified using practice effect trajectories from Months 6 or 12 to Month 24

38. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and EDSS assessed using Spearman's Rank-order Correlation at baseline, Months 6, 12, 18 and 24

39. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and EDSS assessed using Spearman's Rank-order Correlation over 24 months

Previous secondary outcome measures:

1. Cross-sectional correlation between supervised digital outcome measures with FL MS and unsupervised digital outcome measures with FL MS measured using Spearman's Rank-order Correlation at Months 6, 12, 18, and 24

2. Cross-sectional correlation between composite digital cognition outcome with FL MS and ARCS total score measured using Spearman's Rank-order Correlation at baseline, Month 12, and Month 24

3. Cross-sectional correlation between digital cognition outcome with FL MS and Expanded Disability Status Scale (EDSS) measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

4. Cross-sectional correlation between digital cognition outcome with FL MS and Redenlab score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

5. Cross-sectional correlation between composite digital cognition outcome with FL MS and MSReactor Cognitive score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

6. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and Patient-reported outcome measures (PROMs)- Multiple Sclerosis International Quality of Life (MusiQoL) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24

7. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs- Work Productivity and Activity Impairment Questionnaire: General Health (WPAI: GH) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24

8. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs- Patient Health Questionnaire-9 (PHQ-9) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24

9. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs- Penn State Worry Questionnaire (PSWQ-15) assessed using Spearman's Rank-order Correlation at baseline, Months 12 and 24

10. Cross-sectional correlation between digital outcome measures with FL MS and MRI parameters assessed using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

11. Longitudinal correlation between supervised digital outcome measures with FL MS and unsupervised digital outcome measures with FL MS measured using Spearman's Rank-order Correlation at Months 6, 12, 18, and 24

12. Longitudinal correlation between composite digital cognition outcome with FL MS and ARCS total score measured using Spearman's Rank-order Correlation at baseline, Month 12, and Month 24

13. Longitudinal correlation between digital cognition outcome with FL MS and EDSS measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

14. Longitudinal correlation between digital cognition outcome with FL MS and Redenlab score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

15. Longitudinal correlation between digital cognition outcome with FL MS and MSReactor Cognitive score measured using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24

16. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs (MusiQoL) assessed using Spearman's Rank-order Correlation over 24 months

17. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test)

and PROMs (WPAI:GH) assessed using Spearman's Rank-order Correlation over 24 months

18. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs (PHQ-9) assessed using Spearman's Rank-order Correlation over 24 months
19. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and PROMs (PSWQ-15) assessed using Spearman's Rank-order Correlation over 24 months
20. Longitudinal correlation between digital outcome measures with FL MS and MRI parameters assessed using Spearman's Rank-order Correlation at Months 6, 12, 18 and 24
21. Time to unscheduled clinical or paraclinical (e.g., MRI) assessment measured using area under the curve (AUC) from baseline to Month 24
22. Time to first treatment (disease-modifying therapy [DMT]) switch measured using AUC from baseline to Month 24
23. Time to prescription of new MS symptomatic drugs measured using AUC from baseline to Month 24
24. Correlation between FL MS Digital Outcomes and subsequent changes in EDSS assessed using Spearman's Rank-order correlation calculated from baseline to Month 12 and Month 12 to 24
25. Cognitive phenotypes identified using a joint latent profile analysis (LPA) on the baseline FL MS Cognitive Test outcome, ARCS Global Score, and MSReactor Cognitive score at baseline (Day 1)
26. Difference in trajectories of FL MS Cognitive Test outcome between participants performing cognitive test every week versus every 2 weeks versus every 4 weeks, measured using a Piecewise Linear Regression Method over the first 6 and 12 months
27. Participant engagement patterns in the first 3 months of usage of FL MS
28. Participant engagement patterns in the first 24 months of usage of FL MS
29. Factors influencing participants engagement (demographics and disease factors) on FL MS, Redenlab, and MSReactor Apps usage recorded from baseline to Month 24
30. Factors influencing participant's adherence (physician-driven factors i.e., phone calls) to self-monitoring recorded from baseline to Month 24
31. In-app FL MS Post-appointment Questionnaire assessed at Month 12 and Month 24
32. FL MS usability score from mHealth Usability Questionnaire assessed at Month 12 and Month 24
33. Percentage of participants with FL MS-associated adverse events (spontaneously reported) assessed through the PROM questionnaire data from baseline to Month 24
34. Number of assessments or run-in period duration required to minimize the impact of learning effect assessed using FL MS to ensure stable FL MS baseline performance measured from baseline to Week 6
35. Changes in digital outcome measures with FL MS and the presence of protocol-defined relapse measured from baseline to Month 24
36. Longitudinal correlation between FL MS Cognitive Test outcome and ARCS total score measured using Spearman's Rank-order Correlation over 12 and 24 months
37. Differences between participants with different learning patterns on FL MS Cognitive Test outcome identified using practice effect trajectories from baseline to Months 6, or 12
38. Differences between participants with different learning patterns on ARCS score identified using practice effect trajectories from baseline to Months 6, or 12
39. Differences between participants with different learning patterns on EDSS scale score identified using practice effect trajectories from baseline to Month 6, or 12
40. Differences between participants with different learning patterns on MRI outcome measures identified using practice effect trajectories from Months 6 or 12 to Month 24
41. Cross-sectional correlation between FL MS digital outcomes (MS gait and hand function test) and EDSS assessed using Spearman's Rank-order Correlation at baseline, Months 6, 12, 18 and 24
42. Longitudinal correlation between FL MS digital outcomes (MS gait and hand function test) and EDSS assessed using Spearman's Rank-order Correlation over 24 months

Overall study start date

05/01/2022

Completion date

31/07/2024

Eligibility

Key inclusion criteria

Current participant inclusion criteria as of 21/12/2023:

1. Be able to provide informed consent to the study, and to participate in the MSBase registry and the MRI repository
2. Aged 18 years or older
3. EDSS in the range of 1.0 to 7.0 inclusive
4. Diagnosis of MS (new or revised) based on the 2017 revised McDonald criteria
5. Untreated or treated with an approved or off-label DMT, if DMT was initiated >3 months ago
6. Must have a personal smartphone
7. Willing to discontinue participation in any other Floodlight application.

Previous participant inclusion criteria:

1. Be able to provide informed consent to the study, and to participate in the MSBase registry and the MRI repository
2. Aged 18 years or older
3. EDSS in the range of 2.0 to 7.0 inclusive
4. Diagnosis of MS (new or revised) based on the 2017 revised McDonald criteria
5. Untreated or treated with an approved or off-label DMT, if DMT was initiated >3 months ago
6. Must have a personal smartphone
7. Willing to discontinue participation in any other Floodlight application.

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

400

Total final enrolment

279

Key exclusion criteria

1. Unable to provide data to establish a minimum dataset
2. Any contraindication for use of smartphone as per the investigator's discretion
3. Known significant cognitive impairment as per the investigator's discretion
4. Not able to tolerate routine clinical MRI
5. Clinical relapse in the 30 days before enrolment.

Date of first enrolment

26/07/2022

Date of final enrolment

09/04/2024

Locations

Countries of recruitment

Australia

Study participating centre

The Alfred

Dept of Neurosciences Level 6
CCS, Alfred Centre
99 Commercial Road
Melb, Vic
Australia
3004

Study participating centre

John Hunter Hospital

Lookout Road
New Lambton Heights, NSW
Australia
2305

Study participating centre

Royal Brisbane & Women's Hospital

Suite 286, St Andrew's Place
33 North St
Spring Hill, QLD
Australia
4000

Study participating centre

Brain & Mind Research Centre

The University of Sydney
Level 3, F23 Administration Building
Camperdown, NSW
Australia
2006

Study participating centre**Eastern Health**

5 Arnold Street
Box Hill, Vic
Australia
3128

Study participating centre**Menzies Institute**

17 Liverpool Street
Hobart, Tas
Australia
7000

Study participating centre**Perron Institute**

8 Verdun Street
Nedlands, WA
Australia
6009

Study participating centre**Monash Health**

246 Clayton Rd
Clayton VIC
Australia
3168

Sponsor information**Organisation**

F. Hoffmann-La Roche Ltd

Sponsor details

Grenzacherstrasse 124
Basel
Switzerland
4070
+41 616878333
global.trial_information@roche.com

Sponsor type

Industry

Website

http://www.roche.com/about_roche/roche_worldwide.htm

Funder(s)**Funder type**

Industry

Funder Name

F. Hoffmann-La Roche

Alternative Name(s)

Hoffman-La Roche, F. Hoffmann-La Roche Ltd.

Funding Body Type

Private sector organisation

Funding Body Subtype

For-profit companies (industry)

Location

Switzerland

Results and Publications**Publication and dissemination plan**

Planned publication both at scientific congresses and in a peer-reviewed journal

Intention to publish date

31/07/2025

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to participant-level data not being a regulatory requirement

IPD sharing plan summary

Not expected to be made available