Effectiveness of an eleven-valent pneumococcal (type 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F) conjugate vaccine against pneumonia in Philippine children: A doubleblind, placebo-controlled, randomised, multicentre, effectiveness study

Submission date	Recruitment status	Prospectively registered
13/09/2005	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
25/01/2006	Completed	[X] Results
Last Edited	Condition category	Individual participant data
05/11/2012	Respiratory	

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Marilla Lucero

Contact details

Research Institute for Tropical Medicine (RITM)
Research Drive
Filinvest Corporate City
Alabang
Muntinlupa City
Philippines
1781
+63 (0)2 807 2634
mglucero@pldtdsl.net

Additional identifiers

Study information

Scientific Title

Acronym

ARIVAC

Study objectives

Primary objective: The vaccine is efficacious in preventing community-acquired X-ray positive pneumonia. The minimum efficacy, estimated by the lower limit of the 95% confidence interval, is 15%. The relative risk of the X-ray positive pneumonia in the vaccine group is less than 0.85 when compared to the placebo group.

Objective 2a: Hypothesis: The vaccine is efficacious in preventing community-acquired pneumonia requiring hospitalization. The relative risk of pneumonia is lower than one when compared to the placebo group; in other words, the vaccine efficacy is higher than 0%.

Objective 2b: Hypothesis: The vaccine is efficacious in preventing community-acquired pneumonia not requiring hospitalization. The relative risk of pneumonia is lower than one when compared to the placebo group; in other words, the vaccine efficacy is higher than 0%.

Objective 2c: Hypothesis: The vaccine is efficacious in preventing culture proven vaccine type-specific invasive pneumococcal disease. The relative risk of culture proven vaccine type-specific invasive pneumococcal disease is lower than one when compared to the placebo group; in other words, the vaccine efficacy is higher than 0%.

Objective 2d: Hypothesis: The eleven-valent pneumococcal conjugate vaccine is safe when administered concomitantly with the vaccines of Expanded Programmes on Immunization (EPI) and Hib vaccine.

For the nested carriage and immunogenicity study:

Objective 2e: Hypothesis: Children immunized with the eleven-valent pneumococcal vaccine have higher concentrations of antipneumococcal polysaccharide antibodies and higher opsonophagocytic activity in comparison to the placebo recipients.

Objective 2f: Hypothesis: Significantly fewer children immunized with the eleven-valent pneumococcal conjugate vaccine will carry vaccine serotypes of Streptococcus pneumoniae than the placebo recipients.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Technical Review Board and Ethical Review Board (Institutional Review Board [IRB]) of the Research Institute for Tropical Medicine (RITM), Philippines, reviewed and approved the original study protocol in 1999. A counterpart ethical review committee at the National Public Health Institute in Finland likewise reviewed the study protocol, and approved it in the same year. The

RITM IRB evaluated yearly the progress of the study and gave its corresponding approval to proceed with the conduct of the trial. The latest approval was awarded on August 8, 2005.

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Pediatric pneumococcal pneumonia, sepsis, meningitis.

Interventions

Study Vaccine Group: An eleven-valent pneumococcal tetanus-diphtheria toxoid conjugated vaccine, diphtheria-tetanus-pertussis whole cell vaccine, Haemophilus influenzae type b vaccine, hepatitis B vaccine, oral polio vaccine, measles vaccine according to national immunization program schedule.

Control (Placebo) Vaccine Group: Saline (NaCl), diphtheria-tetanus-pertussis whole cell vaccine, Haemophilus influenzae type b vaccine, hepatitis B vaccine, oral polio vaccine, measles vaccine according to national immunization program.

Intervention Type

Biological/Vaccine

Phase

Phase III

Primary outcome(s)

Radiologically confirmed, community-acquired pneumonia at least 14 days after the third dose of the study vaccine in first 2 years of life.

Key secondary outcome(s))

Secondary outcomes: World Health Organisation (WHO) Pneumonia

- 1.1. Any episode of community-acquired pneumonia requiring or not requiring hospitalization
- 1.2. Any episode of community-acquired pneumonia requiring hospitalization
- 2. Any episode of community-acquired pneumonia not requiring hospitalization
- 3. Any episode of culture proven invasive pneumococcal disease, determined as vaccine-specific serotype, vaccine-related serotype, vaccine serogroup, or non-vaccine serotype or group of Streptococcus pneumoniae
- 4. Safety of the 11PCV when administered concomitantly with the EPI and Hib vaccines
- 5. Reactogenicity of the 11PCV after each injection
- 6. Immunogenicity and the opsonophagocytic activity (OPA) of the 11PCV in 11PCV and placebo recipients
- 7. Nasopharyngeal carriage of vaccine and non-vaccine specific serotypes of Streptococcus pneumoniae in 11PCV and placebo recipients

Completion date

Eligibility

Key inclusion criteria

For the effectiveness study and the nested immunogenicity study:

- 1. Any child who comes for routine vaccination to a barangay health station in the 48 barangays in the six municipalities of Bohol, the Philippines (i.e. Baclayon, Balilihan, Cortes, Dauis, Panglao and Tagbilaran)
- 2. Considered to be in good health on the basis of medical history and observation taken at the barangay health station
- 3. At least 6 weeks of age and not older than 6 months of age
- 4. Having at least one parent or other legal representative giving their informed consent attested by signature

For the effectiveness study:

Is a resident of any of the barangays within the catchment of the 45 barangay health stations of the six municipalities for at least the past 3 months with his/her family, or intends to stay permanently

For the nested immunogenicity, safety and carriage study:

Is a resident of any of the barangays within the catchment of the three chosen barangay health stations (i.e. Dampas-Tagbilaran, Danao and Main Health Center-Panglao) for at least the past 3 months with his/her family, or intends to stay permanently

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

6 weeks

Upper age limit

6 months

Sex

All

Key exclusion criteria

For the effectiveness study and the nested immunogenicity study: Any child who:

- 1. Has received the first dose of diphtheria, tetanus, pertussis (DTP) vaccine
- 2. Has acute febrile illness (rectal temperature >/= 38.5°C) at the time of inclusion

- 3. Is suspected to have a neurological disease (a contraindication to the DTP vaccine)
- 4. Has history of hospitalization for and/or treatment for immune suppression
- 5. Is enrolled or scheduled to be enrolled in another clinical trial

Date of first enrolment

05/07/2000

Date of final enrolment

18/12/2004

Locations

Countries of recruitment

Philippines

Study participating centre Research Institute for Tropical Medicine (RITM)

Muntinlupa City Philippines 1781

Sponsor information

Organisation

ARIVAC Consortium (Finland)

Funder(s)

Funder type

Other

Funder Name

National Health and Medical Research Council (NHMRC) (Australia)

Alternative Name(s)

National Health and Medical Research Council, Australian Government, NHMRC National Health and Medical Research Council, NHMRC

Funding Body Type

Government organisation

Funding Body Subtype

Location Australia
Funder Name European Commission (Belgium) - DG Research, INCO Programme
Funder Name Finnish Academy (Finland)
Funder Name Finnish Ministry of Foreign Affairs (Finland)
Funder Name Physicians for Social Responsibility (PSR) (Finland)
Funder Name Program for Appropriate Technology and Health (PATH) (USA)
Funder Name Global Alliance for Vaccines, Accelerated Development and Implementation Plan (GAVI ADIP Pnc) (Switzerland)
Funder Name Provincial Government of Bohol (Philippines)
Funder Name Local government units of Tagbilaran City, Dauis, Panglao, Balilihan, Cortez and Baclayon (Philippines)

National government

Funder Name

Research Institute for Tropical Medicine (RITM) (Philippines)

Funder Name

National Public Health Institute (KTL) (Finland)

Funder Name

University of Colorado (USA)

Funder Name

University of Queensland (Australia)

Alternative Name(s)

University of Queensland in Australia, University of Queensland - Australia, The University of Queensland | Brisbane QLD, uniofqld, The University of Queensland, UQ

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

Australia

Funder Name

Sanofi Pasteur (France)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	21/07/2008		Yes	No
Other publications	evaluation	07/06/2012		Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes