

# Effects of hot pack and ice pack intervention and electrical intervention on sensation and movement ability of the arm and hand in individuals with acute stroke

<b>Submission date</b> 13/05/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 25/05/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 03/07/2025	<b>Condition category</b> Nervous System Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Over 85% of the post-stroke population will experience impaired upper-limb functionality. The functional recovery in the upper limbs for post-stroke is more difficult than that in the lower limbs. Furthermore, the coordination of both sides of upper extremity is needed to accomplish most of the daily activities, such as face washing, tooth brushing, eating, and getting dressed. Therefore, helping patients with stroke recover the use of their affected upper limbs seems to be very essential. Hence, the purpose of this clinical trial is to investigate the effects of electrical stimulation and the hot pack and ice pack stimulation on the sensory and motor recovery in upper extremity for individuals with stroke in acute stage.

### Who can participate?

Acute stroke patients with hemiplegia (paralysis on one side of the body) in Taiwan

### What does the study involve?

Participants were allocated into the electrical stimulation group, thermal stimulation group and conventional physical therapy group. Participants in each group underwent a 5-day intervention. The intervention for the electrical stimulation group includes electrodes attached to the forearm, and, participants might feel itchy during the electrical stimulation protocol. Participants will undergo a total 10 intervention sessions, 2 sessions in a day for 5 days. Participants in the thermal stimulation group will receive a 30-minute heat and cold stimulation, 2 sessions per day for 5 days. Participants will be requested to sit with both hands flat on the table. During the session, the therapist encouraged the participants to move away their limb from the hot pack or ice pack through active movements.

### What are the possible benefits and risks of participating?

The possible benefits are motor function recovery. The possible risks include scald and allergy to the ice pack.

Where is the study run from?  
E-Da Hospital (Taiwan)

When is the study starting and how long is it expected to run for?  
December 2014 to January 2016

Who is funding the study?  
This work was supported by the Chi-Mei KMU Joint Project (106CM-KMU-12), Ministry of Science and Technology in Taiwan (MOST 109-2221-E-037-003-) and the NSYSU-KMU Joint Research Project (#NSYSUKMU 105-P009).

Who is the main contact?  
Hong-Chi Wang  
ohya708@hotmail.com

## Contact information

**Type(s)**  
Principal Investigator

**Contact name**  
Mr Hong-Chi Wang

**Contact details**  
100, Shih-Chuan 1st Road  
Kaohsiung  
Taiwan  
80708  
+886 (7)3121101 ext. 2737 ext. 624  
ohya708@hotmail.com

## Additional identifiers

**EudraCT/CTIS number**  
Nil known

**IRAS number**

**ClinicalTrials.gov number**  
Nil known

**Secondary identifying numbers**  
EMRP-103-113

## Study information

**Scientific Title**  
Comparison of the effects of thermal stimulation and transcutaneous electrical nerve stimulation on upper extremity sensory and motor function in individuals with acute stroke: a randomized controlled pilot study

**Study objectives**

The sensory stimulation might improve the sensory and motor function for individuals with acute stage stroke.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Approved 28/01/2015, Institutional review board of E-Da hospital (No.1, Yida Road, Jiaosu Village, Yanchao District, Kaohsiung City, Taiwan (R.O.C) 82445; +886 7-6151100 ext. 5110; ed107339@edah.org.tw), ref: EMRP-103-113

**Study design**

Single center interventional randomized controlled trial

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Hospital

**Study type(s)**

Treatment

**Participant information sheet**

No participant information sheet available

**Health condition(s) or problem(s) studied**

intervention for individuals with acute stroke

**Interventions**

Participants in this study will be allocated into thermal stimulation group, transcutaneous electrical nerve stimulation group, and the control group by sealed envelope.

For thermal stimulation (TS) group, the following equipment and intervention protocol were used. The hot and cold stimulation devices used were a Firstek heating circulator water bath (B300, Firstek Corp, Taiwan) and a Firstek cooling circulator water bath (B401L, Firstek Corp, Taiwan), respectively. Each was connected to a temperature therapy pad (TP22E, Gaymer Corp, USA). Water temperature was adjusted or maintained at a constant as necessary by using the operation panel, and hot or cold water was then circulated to the therapy pads. The therapy pads (38 × 55 cm) were made of soft materials that allowed the forearm to be covered completely. The participants received hot and cold stimulation in 30-min sessions administered twice daily (once in the morning and afternoon, respectively) over 5 days, totaling 10 sessions. In accordance with the procedure used in one study, participants receiving TS were instructed to sit with both hands flat on the table. Heat stimulation was applied to their healthy arm for no more than 15 s. A thermometer was placed on the stimulated body part to prevent frostbite or burns. For the hot and cold stimulation, the temperatures were set at 51°C and 4°C, respectively. The

therapy pad was wrapped around the palm and wrist of the affected limb. During the session, the therapist encouraged the participants to pull their limb from the therapy pad through active movements. They were instructed to remove their healthy hand when they began feeling discomfort or when a score of seven had been reached on a standard 10-point visual analog scale (administered by the therapist), and the time from the beginning of the session to this point was recorded. The same procedure was repeated with the participants' affected hand. If no adverse skin reactions occurred, heat was applied on their affected arm 10 consecutive times, separated by 3 min of rest. Cold therapy involved the same procedure and was applied alternately with heat therapy. With both heat and cold therapy, stimulation was applied for 15 s, followed by at least 30 min of rest. Heat and cold were applied 20 times in each session (in order: 10 cycles of heat therapy, 10 cycles of cold therapy, 10 cycles of heat therapy, 10 cycles of cold therapy). During each session, the therapist constantly the skin surface temperature on the tested limb to prevent frostbite or burns. Moreover, the participants' blood pressure, heart rates, and breathing before and after the intervention were monitored, and the rest periods were extended as necessary.

For transcutaneous electrical nerve stimulation (TENS) group, the following equipment and intervention protocol were conducted. A portable transcutaneous electrical nerve stimulator (TRIO-310) was used. Patches were adhered to the back of the forearm and the side of the palm. The skin was cleaned with alcohol before and after each session for disinfection and to reduce the possibility of increased electrical resistance. Wounds were avoided during the intervention. The TENS settings were as follows: pulse width 200  $\mu$ s, output frequency 100 Hz, and output time 30 min. The output frequency was selected mainly for the purpose of stimulating the A $\beta$  fibers, which produce sensations of light touch and pressure[19]. The current strength was adjusted as the maximum that the participant could withstand. As with the TS, TENS was applied in 30-min sessions administered twice daily (once in the morning and afternoon, respectively) over 5 days, totaling 10 sessions. Moreover, the therapist monitored the participants during each session and measured their blood pressure, heart rates, and breathing before and after the intervention, adjusting the rest periods as necessary and taking care to prevent electrical burns.

Participants in the control group received regularly scheduled rehabilitation therapy (1 h each of physical and occupational therapy). The physical therapy included therapeutic exercise, facilitation training, and functional training. Occupational therapy involved hand function training and training on activities of daily living.

### **Intervention Type**

Mixed

### **Primary outcome measure**

Measured at baseline (before intervention) and after 5 days (after completing intervention):

1. Motor recovery is measured using Fugl-meyer upper extremity scale assessment
2. Motor function recovery is measured by Brunnstrom stage assessment
3. Sensory is measured by minimal current perception

### **Secondary outcome measures**

Measured at baseline (before intervention) and after 5 days (after completing intervention):

The spasticity of upper limb is measured by Modified ashworth scale assessment

### **Overall study start date**

05/12/2014

**Completion date**

27/01/2016

## **Eligibility**

**Key inclusion criteria**

1. Aged 20 years or older
2. Have a stroke for the first time
3. Have been examined by a physician, and were approved to receive rehabilitation treatment
4. Hospitalized during the acute phase
5. Hemiparesis
6. Do not have obvious cognitive impairments
7. Can independently maintain a sitting posture for at least 30 min

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

30

**Total final enrolment**

27

**Key exclusion criteria**

1. Have skin conditions or injuries (e.g., wounds) on upper limbs or have other contraindications for electrotherapy or TS (e.g., a malignant tumor)
2. Have a language disorder (e.g., aphasia) and is therefore unable to communicate or comply with instructions
3. Have other orthopedic conditions (e.g., severe arthritis) or nerve damage (e.g., peripheral nerve injury) affecting movement in their upper limbs
4. Have diabetes or complete sensory impairment not caused by stroke (e.g., peripheral vascular disease or neuropathy)
5. Have developed neurological disorders during the experiment period or other conditions that may affect the study results
6. Have uncontrolled hypertension, unstable angina, a history of myocardial infarction or epilepsy (excepting febrile seizures) in the past 3 months, or a pacemaker
7. Have participated in other rehabilitation trials or drug trials
8. Unable to cooperate with the researchers because of cognitive or personal reasons or refused to provide written informed consent

**Date of first enrolment**

28/01/2015

**Date of final enrolment**

15/01/2016

## **Locations**

### **Countries of recruitment**

Taiwan

### **Study participating centre**

#### **E-Da hospital**

No. 1, Yida Road

Jiaosu Village

Yanchao District

Kaohsiung City

Taiwan

82445

## **Sponsor information**

### **Organisation**

Kaohsiung Medical University

### **Sponsor details**

No.100, Tzyou 1st Rd.

Sanmin Dist.

Kaohsiung City

Taiwan

80756

+886 (7)3121101 ext. 2646 ext. 614

yuen@kmu.edu.tw

### **Sponsor type**

University/education

### **Website**

<http://english2.kmu.edu.tw/front/bin/home.phtml>

### **ROR**

<https://ror.org/03gk81f96>

## **Funder(s)**

### **Funder type**

Government

**Funder Name**

Ministry of Science and Technology, Taiwan

**Alternative Name(s)**

Ministry of Science and Technology, R.O.C. (Taiwan), Ministry of Science and Technology of Taiwan, MOST

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

Taiwan

**Funder Name**

Chi-Mei Medical Center and Kaohsiung Medical University

**Funder Name**

National Sun Yat-sen University and Kaohsiung Medical University

## Results and Publications

**Publication and dissemination plan**

Planned publication in a high-impact peer-reviewed journal

**Intention to publish date**

31/07/2022

**Individual participant data (IPD) sharing plan**

The datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication.

**IPD sharing plan summary**

Published as a supplement to the results publication

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		28/06/2024	03/07/2025	Yes	No