Concomitant chemoradiotherapy for treatment of non-small cell lung cancer - the Conrad study

Submission date	Recruitment status No longer recruiting	Prospectively registered		
28/04/2009		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
29/05/2009	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
21/01/2019	Cancer			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Roy Bremnes

Contact details

Oncology Department University Hospital of North Norway Tromsø Norway 9038

roy.bremnes@unn.no

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

Norwegian Social Science Data Services no. 15175; UNN (PVO) no. 0147

Study information

Scientific Title

A randomised, multicentre phase III trial of combination chemotherapy +/- thoracic radiotherapy in the treatment of patients with stage III non-small cell lung cancer not eligible for radical therapy

Acronym

Conrad

Study objectives

That a combination of chemotherapy and thoracic radiation is superior to chemotherapy alone in the treatment of patients with stage III non-small cell lung cancer, non-eligible for radical therapy, with overall survival as primary endpoint.

Protocol can be found at: http://www.nlcg.no/uploads/conrad_protokoll.pdf

Ethics approval required

Old ethics approval format

Ethics approval(s)

- 1. Regional Ethical Review Board approved on the 19th July 2006
- 2. Norwegian Social Science Data Services approved on the 7th November 2006

Study design

Open randomised multicentre phase III trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Can be found at: http://www.nlcg.no/uploads/conrad_protokoll.pdf

Health condition(s) or problem(s) studied

Non-small cell lung cancer

Interventions

The participants will be randomised to the following two arms:

Arm 1: Chemotherapy and thoracic radiation

Arm 2: Chemotherapy alone

Chemotherapy:

All patients will receive 4 cycles of identical regimes of chemotherapy: Oral vinorelbine tablets 60 mg/m² orally (per os) day 1 and 8 and intravenous carboplatin AUC = 5 (Calvert's formula) over one hour day 1. However, the doses will be adjusted according to age (patients age greater than 75 year will be given 75% of full dose from cure no.1) and haematological toxicity.

Schedule of radiation therapy:

Simulator planned, two opposing fields with fractionation 2.8 Gy \times 15. The radiotherapy should start at the same time as or just after cycle number two. Cycle number three should be given three weeks after cycle number two.

Intervention Type

Other

Phase

Phase III

Primary outcome measure

Overall survival, followed-up for 52 weeks.

Secondary outcome measures

- 1. Health related quality of life, assessed by the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire for Cancer patients (EORTC QLQ-C30) version 3.0 and the lung cancer specific module EORTC QLQ-LC13. The questionnaires will be sent to the patients every month during the treatment period and every second month after the treatment period for a total of 10 questionnaires completed per patient.
- 2. Time to progression, followed-up for 52 weeks
- 3. In field relapse, followed-up for 52 weeks
- 4. Drug related adverse event frequency and severity, followed-up for 52 weeks
- 5. Health economics, followed-up for 52 weeks

Overall study start date

15/11/2006

Completion date

15/11/2013

Eligibility

Kev inclusion criteria

- 1. Chemo-naive patients with non-small cell lung cancer (NSCLC) locally advanced stage III, not candidates for radical radiotherapy and with no pleural effusion
- 2. Performance status (World Health Organization [WHO]) 0 2
- 3. Ability to understand oral and written study information
- 4. Both males and females, no age limit
- 5. S-creatinine less than 1.5 times upper reference limit, bilirubin and S-transaminase levels less than 2 times upper limits. Normal white blood-cell and platelet count.

Participant type(s)

Patient

Age group

Other

Sex

Both

Target number of participants

352

Key exclusion criteria

- 1. Other active malignancies
- 2. Pregnancy or breast feeding

Date of first enrolment

15/11/2006

Date of final enrolment

15/11/2013

Locations

Countries of recruitment

Norway

Study participating centre Oncology Department

Tromsø Norway 9038

Sponsor information

Organisation

Pierre Fabre Pharma Norden AB (Sweden)

Sponsor details

Turebergs Torg 1 Sollentuna Sweden SE-191 47

_

tore.aalberg@pierre-fabre.com

Sponsor type

Industry

Website

http://www.pierre-fabre.com

ROR

https://ror.org/04hdhz511

Funder(s)

Funder type

Industry

Funder Name

Regional Health Authorities (HELSE NORD) (Norway)

Funder Name

Pierre Fabre Pharma Norden AB (Sweden)

Funder Name

Please note that the Conrad study was initiated by the Norwegian Lung Cancer Group.

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	17/09/2013	21/01/2019	Yes	No
Results article	results	01/06/2014	21/01/2019	Yes	No
Results article	results	01/05/2015	21/01/2019	Yes	No