

Localization of transcranial magnetic stimulation in treatment of depression

Submission date 26/01/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 11/04/2017	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 27/12/2023	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Major depressive disorder (MDD) is one of the most common mental health conditions in the world. The symptoms can vary greatly from person to person, but they generally include low mood, problems with sleeping and/or eating, and a general loss of interest in life. Treatment often relies heavily on antidepressant medications, which work by increasing the activity and levels of a group of chemicals in the brain (neurotransmitters) and talking therapy. Repetitive transcranial magnetic stimulation (rTMS) is a relatively new treatment of depression. It involves using magnetic fields to stimulate brain cells in order to improve the symptoms of depression. Studies have shown that this treatment can produce changes in the brain that have been found to be related to mood regulation. This study is looking at the effects of targeting rTMS on a particular area of the brain.

Who can participate?

Patients with MDD aged between 15 and 60 years old.

What does the study involve?

All participants receive transcranial magnetic stimulation (rTMS) as a part of their normal care. During the first and sixth week of their treatment, participants are interviewed and asked to complete a questionnaire in order to assess the severity of their depression symptoms and their functioning, in order to find out whether the rTMS treatment has made any difference to their condition.

What are the possible benefits and risks of participating?

There are no direct benefits involved with participating in this study. Strong magnetic fields used in MRI and TMS are a risk for patients with magnetic objects or medical devices in the body and so patients with these are unable to take part. There are no other notable risks involved with participating.

Where is the study run from?

Helsinki University Hospital (Finland)

When is the study starting and how long is it expected to run for?
January 2015 to January 2019

Who is funding the study?
Helsinki University Hospital (Finland)

Who is the main contact?
Dr Tuukka Raij

Contact information

Type(s)
Scientific

Contact name
Dr Tuukka Raij

ORCID ID
<https://orcid.org/0000-0002-9834-5570>

Contact details
PL 348
Helsinki
Finland
00029 HUS

Additional identifiers

Protocol serial number
05012017

Study information

Scientific Title
Imaging-based optimization of repetitive transcranial magnetic stimulation for depression

Study objectives
Effectiveness of rTMS treatment of depression depends on E-field location in the left dorsolateral cortical region.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Helsinki University Hospital Ethics Committee for gynaecology and obstetrics, pediatrics and psychiatry, 11/06/2015, ref: 215/13/03/03/2015

Study design
Observational cohort study

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Major depression

Interventions

The brain stimulation treatment and whole-head MRI is part of the normal clinical care of the patients. For the study purposes, location of the stimulating coil is registered to enable computation of stimulation-induced electric fields and to compare the cortical distribution of the fields with treatment response. Patients admitted to the Helsinki University Central Hospital (HUCH) Aurora neuromodulation unit for treatment of MDD with repetitive transcranial magnetic stimulation (rTMS) are first contacted by phone to inquire about their interest in participating in the study. At the first study visit, participants sign informed consent, fill in the patient health questionnaire-9 (phq-9, assessing symptoms of depression), and are evaluated by the study psychiatrist. The evaluation includes the Mini-International Neuropsychiatric Interview (MINI) for diagnostics, Montgomery-Åsberg Depression Rating Scale (MADRS), Social and Occupational Functioning Assessment Scale (SOFAS), and determination of the patient's TMS motor threshold, which is used to customize the rTMS intensity. TMS coil is placed over the EEG electrode location F5 situated over the left dorsolateral prefrontal cortex, and a mark for the coil position is saved in the neuronavigation system. On the subsequent 23 visits, rTMS is applied at the marked position using a TMS neuronavigation system. Exact coil position with respect to the anatomical MRI is recorded during all TMS sessions. TMS is given with a Magstim Rapid stimulator at 110-120% of motor threshold, (4-second trains of 40 pulses at 10 Hz, interleaved with 26-s waiting periods), on 3-5 times days per week at weeks 1-4, on two days at week 5, and once at week 6. At the end of the last rTMS visit at week 6 the participants again fill in the phq-9 and meet the psychiatrist for MADRS and SOFAS assessments. There is no follow-up in the study.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Transcranial magnetic stimulation

Primary outcome(s)

Depression severity is measured using the Montgomery-Åsberg Depression Rating Scale (MADRS) total at week 1 and 6.

Key secondary outcome(s)

1. Depression severity is measured using The Patient Health Questionnaire (PHQ-9) at week 1 and 6
2. Functioning is measured using Social and Occupational Functioning Assessment Scale (SOFAS) at week 1 and 6

Completion date

31/01/2023

Eligibility

Key inclusion criteria

1. Major depressive disorder in MINI interview
2. Aged 15-60 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

15 years

Upper age limit

60 years

Sex

All

Key exclusion criteria

1. Serious psychiatric or somatic illness
2. MRI/TMS contraindications

Date of first enrolment

01/05/2016

Date of final enrolment

31/10/2022

Locations

Countries of recruitment

Finland

Study participating centre

Helsinki University Hospital

Aurora ECT unit

Helsinki

Finland

00029 HUS

Sponsor information

Organisation

Helsinki University Hospital

ROR

<https://ror.org/02e8hzhf44>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Helsinki University Hospital

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available to protect patient identity.

IPD sharing plan summary

Not expected to be made available