

Protocol for a Randomised Phase II Study of the Stanford V regimen compared with ABVD for the treatment of Advanced Hodgkin's Disease

Submission date	Recruitment status	<input type="checkbox"/> Prospectively registered
19/08/2002	No longer recruiting	<input type="checkbox"/> Protocol
Registration date	Overall study status	<input type="checkbox"/> Statistical analysis plan
19/08/2002	Completed	<input checked="" type="checkbox"/> Results
Last Edited	Condition category	<input type="checkbox"/> Individual participant data
18/10/2018	Cancer	

Plain English summary of protocol

<http://cancerhelp.cancerresearchuk.org/trials/a-trial-comparing-stanford-v-chemotherapy-with-abvd-chemotherapy-for-advanced-hodgkins-lymphoma>

Contact information

Type(s)

Scientific

Contact name

Mr Paul Mouncey

Contact details

Lymphoma Trials Office
222 Euston Road
London
United Kingdom
NW1 2DA
+44 (0) 20 7679 8038
pm@ctc.ucl.ac.uk

Additional identifiers

Protocol serial number

STANFORD V

Study information

Scientific Title

Protocol for a Randomised Phase II Study of the Stanford V regimen compared with ABVD for the treatment of Advanced Hodgkin's Disease

Study objectives

To compare the Stanford V regimen with Adriamycin, Bleomycin, Vinblastine, Dacarbazine (ABVD) for the treatment of Advanced Hodgkin's Disease.

Ethics approval required

Old ethics approval format

Ethics approval(s)

This trial is currently awaiting approval for an amendment, and sites will be able to recruit once this has been granted.

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Lymphoma (Hodgkin's)

Interventions

STANFORD V regimen: Chemotherapy with mustine, doxorubicin, vinblastine, prednisolone, vincristine, bleomycin and etoposide

ABVD Regimen: Chemotherapy with doxorubicin, bleomycin, vinblastine and dacarbazine

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

Mustine, doxorubicin, vinblastine, prednisolone, vincristine, bleomycin, etoposide and dacarbazine.

Primary outcome(s)

Not provided at time of registration

Key secondary outcome(s))

Not provided at time of registration

Completion date

30/06/2008

Eligibility

Key inclusion criteria

1. Histologically confirmed Hodgkin's disease
2. Clinical stage IIB, IIIA, IIIB or IV
3. Aged 18 - 60 years
4. No previous history of malignancy, except for basal cell or squamous cell carcinoma of the skin
5. Normal values for Full Blood Count (FBC), hepatic and renal function, unless directly attributable to involvement by Hodgkin's disease
6. Written informed consent
7. All patients must be assessed by the treating haematologist/medical oncologist and radiation oncologist TOGETHER prior to study entry. This is an absolute requirement for the study eligibility

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

60 years

Sex

All

Key exclusion criteria

1. Previous therapy for Hodgkin's disease
2. Clinical evidence of infection with the Human Immunodeficiency Virus (HIV)
3. Pre-existing cardiac or pulmonary disease

Date of first enrolment

01/01/2001

Date of final enrolment

30/06/2008

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Lymphoma Trials Office
London
United Kingdom
NW1 2DA

Sponsor information

Organisation
British National Lymphoma Investigation (BNLI) (UK)

Funder(s)

Funder type
Research organisation

Funder Name
British National Lymphoma Investigation (BNLI) (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	10/11/2009		Yes	No
Other publications	questionnaire-based audit	01/10/2007		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Plain English results				No	Yes