Effect of L-thyroxine on progression of Carotid Atherosclerosis in Subclinical Hypothyroidism

Submission date	Recruitment status	Prospectively registered
21/05/2008	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
21/05/2008	Completed	Results
Last Edited	Condition category	[] Individual participant data
04/03/2009	Circulatory System	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr John David Spence

Contact details

Stroke Prevention & Atherosclerosis Research Centre (SPARC)
Robarts Research Institute
Siebens-Drake Research Bldg.
1400 Western Rd
London, Ontario
Canada
N6G 2V2
+1 519 663 3113
dspence@robarts.ca

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

MCT-79197

Study information

Scientific Title

Effect of L-thyroxine on progression of Carotid Atherosclerosis in Subclinical Hypothyroidism: a single centre, randomised, two-arm parallel, placebo-controlled drug intervention

Acronym

T4CASH

Study objectives

Primary hypothesis:

That treatment of patients with subclinical hypothyroidism with L-thyroxine to achieve a target thyroid stimulating hormone (TSH) less than 2 and greater than 0.4 mIU/L and free thyroxine (FT4) greater than 4 and less than 25 pmol/L, will slow progression of carotid plaque volume and carotid intima-media thickness (IMT), compared to placebo.

Secondary hypotheses:

That treatment with L-thyroxine will improve control of plasma lipids and homocysteine, and reduce levels of C-reactive protein, weight, waist circumference and insulin resistance.

Please note that as of 04/03/2009 the anticipated end date in this record was amended; the previous date at the time of registration was:

Initial anticipated end date: 31/07/2007

Ethics approval required

Old ethics approval format

Ethics approval(s)

Research Ethics Board of the University of Western Ontario gave approval on the 24th June 2004 (ref: 10458).

Study design

Single centre, randomised, double blind (participant, investigator, caregiver, outcome assessor and data analyst), two-arm parallel, placebo-controlled drug intervention

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Atherosclerosis, subclinical hypothyroidism

Interventions

1. L-thyroxine, beginning at 0.0125 and gradually increased to achieve target TSH level greater than 0.4 and less than 2 and FT4 greater than 4 and less than 25 pmol/L for 18 months

2. Matching placebo, with dummy dose adjustments for 18 months

Contact for public queries:

Tisha Mabb

Lab Manager

Stroke Prevention & Atherosclerosis Research Centre (SPARC)

Robarts Research Institute

Siebens-Drake Research Bldg.

1400 Western Rd

London, Ontario

N6G 2V2

Canada

Email: tisha@robarts.ca

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

L-thyroxine

Primary outcome measure

Rate of progression of carotid plaque volume (mm³) measured by 3-D ultrasound at baseline and 18 months.

Secondary outcome measures

- 1. Intima-media thickness (IMT), measured at baseline and 18 months
- 2. Weight, waist circumference, measured every 6 months
- 3. Insulin resistance by Homeostasis Model Assessment (HOMA), homocysteine, lipids, measured at baseline and 19 months

Overall study start date

01/07/2006

Completion date

31/07/2010

Eligibility

Key inclusion criteria

Male or female patients:

1. With subclinical hypothyroidism, defined as a TSH of 3 to 10 mU/L, and a normal serum free thyroxine level (FT4 11 to 25 pmol/L), and

- 2. Measurable plaque in the carotid arteries
- 3. Aged greater than or equal to 45 years

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

254

Key exclusion criteria

- 1. Unwilling to give informed consent
- 2. History of atrial fibrillation
- 3. Unstable angina, or angina requiring nitroglycerine as often as once a month
- 4. Unlikely to adhere to the protocol
- 5. Unlikely to survive 18 months because of severe illness such as cancer
- 6. At risk of pregnancy

Date of first enrolment

01/07/2006

Date of final enrolment

31/07/2010

Locations

Countries of recruitment

Canada

Study participating centre

Stroke Prevention & Atherosclerosis Research Centre (SPARC)

London, Ontario Canada N6G 2V2

Sponsor information

Organisation

Robarts Research Institute (Canada)

Sponsor details

P.O. Box 5015 100 Perth Drive London, Ontario Canada N6A 5K8 +1 519 663 5777 lchapman@robarts.ca

Sponsor type

Research organisation

Website

http://www.robarts.ca/home.php

ROR

https://ror.org/01e36dv41

Funder(s)

Funder type

Research organisation

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - http://www.cihr-irsc.gc.ca (ref: MCT-79197)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration