

# The efficacy and cost-effectiveness of confrontational counselling for smoking cessation in smokers with previously undiagnosed COPD: a randomised controlled trial

<b>Submission date</b> 28/04/2006	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 28/04/2006	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 15/06/2010	<b>Condition category</b> Respiratory	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Daniel Kotz

**Contact details**  
University Maastricht (UM)  
CAPHRI Research Institute  
Department of General Practice  
P.O. Box 616  
Maastricht  
Netherlands  
6200 MD  
+31 (0)43 3882893  
d.kotz@hag.unimaas.nl

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

NTR627

## **Study information**

**Scientific Title**

**Acronym**

COSMO

**Study objectives**

Confrontation with the results from spirometry as part of counselling for smoking cessation in smokers with not earlier diagnosed mild to moderate COPD is effective with regard to prolonged abstinence from smoking during a period of 12 months.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Ethics approval received from the local medical ethics committee

**Study design**

Randomised controlled trial

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Not specified

**Study type(s)**

Treatment

**Participant information sheet**

**Health condition(s) or problem(s) studied**

Tobacco addiction, chronic obstructive pulmonary disease (COPD)

**Interventions**

### Intervention groups:

1. Confrontational counselling delivered by a pulmonary nurse and pharmacotherapy
2. Health education and promotion delivered by a pulmonary nurse and pharmacotherapy
3. 'Care as usual' delivered by the general practitioner

### Intervention Type

Other

### Phase

Not Specified

### Primary outcome measure

The primary outcome of this study is biochemically validated prolonged abstinence from smoking during a period of 12 months. A smoker is defined as prolonged abstinent if he/she is a non-smoker (no cigarette smoked during the preceding seven days), at the end of the intervention (day 50), and at the follow-up visits after 6 months (day 197) and 12 months (day 380). Non-smoking is verified by urine cotinine. Participants with a cotinine-value of >50 ng/ml are regarded as smokers as well as participants who are lost to follow-up.

### Secondary outcome measures

1. Point prevalence of non-smokers (no cigarette smoked during the preceding seven days) at the end of the intervention period and at 6- and 12-month follow-up
2. Self-reported number of quit attempts and temporary or complete relapse
3. Attitudes, social norms and self-efficacy with regard to smoking cessation
4. Lung function (FEV1 post-bronchodilatory and FEV1/FVC) at baseline and at 12-month follow-up
5. Anthropometry: physical height and weight at baseline, at the end of the intervention period and at 6- and 12-month follow-up
6. Perceived specific health-related complaints (impairments and functional disabilities in everyday life)
7. Health-related quality of life
8. Mental health (fear, depression)
9. Smoking related cognitions (risk perception, health concerns, self-exempting beliefs)
10. Number of unplanned visits to the general practitioner or specialized physician due to respiratory complaints and the number, severity and frequency of exacerbations (self-reported and/or reported by the general practitioner or specialist)

### Overall study start date

01/02/2004

### Completion date

01/02/2008

## Eligibility

### Key inclusion criteria

1. Age between 35 through 70 years
2. Smoker with a smoking history of >10 pack years of cigarettes
3. Motivated to stop smoking
4. One or more of the following symptoms are present: cough, progressive persistent shortness

- of breath (worse during exercise or respiratory infections) or sputum production
5. Bronchus obstruction detected by spirometry: FEV1/FVC-ratio <70% and postbronchodilatory FEV1 >50% predicted (= mild or moderate COPD/GOLD I or II)
6. Competent enough in speaking the Dutch language

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

350

**Key exclusion criteria**

1. Known by the general practitioner or in the second-line medical care with the diagnosis asthma or COPD (e.g. chronic bronchitis, lung emphysema)
2. Spirometry performed during the preceding 12 months
3. FEV1 <50% predicted (= severe or very severe COPD/GOLD III or IV)
4. Contraindications for the intake of the medication such as an acute myocardial infarction and hypersensitiveness for nortriptyline
5. Current use of antidepressants
6. Quit smoking attempt(s) using nortriptyline or bupropion during the preceding 6 months
7. Co-morbidity: hypersensitiveness towards nortriptyline, tuberculosis, porphyria, epilepsy, Parkinson's disease, glaucoma, bronchial carcinoma or any other life threatening disease

**Date of first enrolment**

01/02/2004

**Date of final enrolment**

01/02/2008

**Locations****Countries of recruitment**

Netherlands

**Study participating centre**

University Maastricht (UM)

Maastricht

Netherlands

6200 MD

**Sponsor information**

**Organisation**

University Maastricht (UM), CAPHRI Research Institute (The Netherlands)

**Sponsor details**

P.O. Box 616  
Maastricht  
Netherlands  
6200 MD  
+31 (0)43 3882446  
e.habets@caphri.unimaas.nl

**Sponsor type**

University/education

**ROR**

<https://ror.org/02jz4aj89>

**Funder(s)****Funder type**

University/education

**Funder Name**

University Hospital Maastricht (AZM) (Netherlands)

**Funder Name**

Partners in Care Solutions (PICASSO) (Netherlands)

**Funder Name**

Dutch Asthma Foundation (Netherlands)

**Results and Publications****Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan**

## IPD sharing plan summary

Not provided at time of registration

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/11/2010		Yes	No