

Robotic-assisted versus open removal of the bladder in bladder cancer patients

Submission date 01/10/2019	Recruitment status Recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 04/11/2019	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 17/12/2024	Condition category Cancer	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Bladder cancer is one of the most common forms of cancer. For most patients, the bladder cancer is superficial and can be treated locally in the bladder. However, for about one in five patients, cancer invades the bladder muscle and for these the most common curative treatment is cystectomy. Cystectomy can be performed either as open surgery, with a longer incision in the lower abdomen, or as a robot-assisted laparoscopic procedure. Robot-assisted laparoscopic cystectomy has been developed during the last 15 years and is routinely performed at some centres around the world. However, it is not entirely certain that a laparoscopic cystectomy is as safe as open surgery, or that there are sufficient advantages to offset the higher procedure-related costs. The aim of this study is, therefore, to find out whether robot-assisted laparoscopic cystectomy is oncologically as safe as open cystectomy, whether there is a difference in rates of complications, and if it is cost-effective

Who can participate?

Men and women with urinary bladder cancer planned to undergo cystectomy

What does the study involve?

Participants are randomly allocated to either robot-assisted laparoscopic cystectomy (experimental group) or open cystectomy (control group)

What are the possible benefits and risks of participating?

Robot-assisted laparoscopic cystectomy may lead to less blood loss, faster recovery, and fewer complications than open cystectomy. It is possible, however, that there may be an increased risk of recurrence or progression from bladder cancer

Where is the study run from?

University of Gothenburg/Sahlgrenska University Hospital (Sweden)

When is the study starting and how long is it expected to run for?

September 2019 to September 2028

Who is funding the study?

The study is funded by the Agreement concerning research and education of doctors in Sweden

Who is the main contact?

Dr Henrik Kjölhede

henrik.kjolhede@vgregion.se

Contact information

Type(s)

Scientific

Contact name

Dr Henrik Kjölhede

ORCID ID

<https://orcid.org/0000-0001-6441-4729>

Contact details

Department of Urology

Institute of Clinical Sciences

University of Gothenburg

Bruna stråket 11B

Sahlgrenska University Hospital

Gothenburg

Sweden

413 45

+46 31 342 10 00

henrik.kjolhede@vgregion.se

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

2019-00717 / 844-18

Study information

Scientific Title

Robotic-Assisted radical Cystectomy vErsus open Radical cystectomy: a randomised non-inferiority trial comparing 3-year recurrence-free survival in patients undergoing cystectomy for urothelial carcinoma of the bladder

Acronym

RACER

Study objectives

Robot assisted laparoscopic cystectomy is non-inferior to open cystectomy with regards to 3-year progression-free survival

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 07/01/2019, Swedish Ethical Review Authority (Etikprövningsmyndigheten, Box 2110, 750 02 Uppsala, Sweden; +46-10-4750800), ref: 2019-00717 / 844-18

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Bladder cancer

Interventions

For both the treatment arms, eligible patients will be offered neo-adjuvant or induction chemotherapy. Randomisation to robot assisted or open surgery will be performed after any preoperative chemotherapy has been finished. Randomisation will be performed through an electronic data capture system, with a computer-generated list. Treatment arms will be stratified by participating centre and muscle-invasive or non-muscle-invasive disease. Both arms will undergo surgery with an identical enhanced recovery protocol.

Participating subjects will receive either robot assisted laparoscopic cystectomy (experimental arm) or open cystectomy (control arm). The urinary deviation can be performed either entirely intra-corporeally or through a small incision extra-corporeally. The type of deviation is at the discretion of the surgeon and patient. All surgeons participating in the study is mandated to have performed at least 10 cystectomies of the relevant type (open or robotic) in the preceding year.

Follow-up will be performed with CT thorax and abdomen, which will be assessed by RECIST criteria, according to the Swedish National Guidelines for urothelial cancer. This follow-up scheme is dependent on the final pathology report, and may be subject to change according to updates in the National Guidelines.

Intervention Type

Other

Primary outcome(s)

Progression-free survival at three years following cystectomy

Key secondary outcome(s)

Surgical outcomes:

1. Rate of complications
2. Blood loss
3. Rate of transfusion
4. Time of stay

Long-term outcomes:

5. Rate of unplanned readmission
6. Quality of life measured by questionnaires FACT-G/Bl-Cys and WHODAS2.0 at baseline, and 3, 6 and 12 months and EQ-5D-5L at baseline and weekly for the first 4 postoperative weeks
7. Total health-care related costs following cystectomy, calculated for the direct and indirect costs related to the surgery for the first 6 months

Completion date

01/09/2028

Eligibility**Key inclusion criteria**

1. Histologically verified urothelial cancer
2. cT1-4a or BCG-unresponsive CIS
3. N0-1, or N2-3 if response on pre-operative chemotherapy to N0-1
4. Able to sign informed consent

Participant type(s)

All

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Distant metastasis (M1)
2. Unfit for either type of procedure
3. Extant ileo- or colostomy
4. Duplicated ureters

Date of first enrolment

04/11/2019

Date of final enrolment

31/12/2025

Locations

Countries of recruitment

Sweden

Study participating centre

University of Gothenburg/Sahlgrenska University Hospital

Department of Urology

Bruna Stråket 11B

Gothenburg

Sweden

413 45

Study participating centre

University of Lund/Skåne University Hospital

Department of Urology

Jan Waldenströms Gata 5

Malmö

Sweden

214 28

Sponsor information

Organisation

Sahlgrenska University Hospital

ROR

<https://ror.org/04vgqjj36>

Funder(s)

Funder type

Government

Funder Name

The study is funded by the Agreement concerning research and education of doctors in Sweden.

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available as ethical approval does not allow it.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet		10/10/2019	04/11/2019	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes