

# Prospective, randomised evaluation of efficacy of antidepressant monotherapy and antidepressant combinations in the treatment of patients with resistant depression

<b>Submission date</b> 03/09/2010	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 27/09/2010	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 27/09/2010	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr Martin Bares

### Contact details

Prague Psychiatric Center  
Ústavní 91  
Prague 8-Bohnice  
Czech Republic  
181 03

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

### Secondary identifying numbers

IGA MZ CR NS 10368-3

# Study information

## Scientific Title

Randomised, open-label study comparing the efficacy of antidepressant monotherapy and antidepressant combinations in the acute treatment of patients with resistant depression

## Acronym

D-REZ-KOMB

## Study objectives

The aim of our study is to examine efficacy of antidepressant monotherapy and combinations of antidepressants in the treatment of resistant depression in common clinical practice. We will test the following null hypotheses:

1. That the reduction of depressive symptoms does not differ between monotherapy and combinations groups
2. The number of responders in both groups does not differ

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

The Ethics Committee of Prague Psychiatric Centre approved on the 21st May 2008

## Study design

Single centre two arm open label randomised active controlled parallel group trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Not specified

## Study type(s)

Treatment

## Participant information sheet

Not available in web format, please use contact details below to request a patient information sheet

## Health condition(s) or problem(s) studied

Resistant depressive disorder

## Interventions

Patients will be randomly allocated to antidepressant monotherapy group or combination of antidepressant group. Specific antidepressant will be chosen by attending psychiatrist according to clinical judgment and with regard to the history of previous treatment. Antidepressant or

combinations of antidepressant will be used in the dose cited in Summary of Products. Both interventions are defined as a new intervention not augmentation of previous antidepressant treatments. We will administer combinations that are commonly used in clinical practice. The total duration of treatment is 6 weeks. We will evaluate stability of outcome in responders after 2 months.

**Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome measure**

The change in the Montgomery-Asberg Depression Rating Scale (MADRS). Patients will be rated at baseline, week 2 week 4 and the end of acute treatment and responders after 2 month of treatment.

**Secondary outcome measures**

Response to treatment that is defined as a greater than or equal to 50% reduction of MADRS score from baseline to the end of treatment.

**Overall study start date**

01/01/2009

**Completion date**

30/06/2011

## Eligibility

**Key inclusion criteria**

1. Patients suffering from major depressive disorder (recurrent or single episode) diagnosed according to Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM IV) criteria, confirmed using the Mini-International Neuropsychiatric Interview (MINI) Czech version 5.0.0.
2. Patients fulfilling at least Stage I criteria for resistant depression according to Thase and Rush, 1997
3. The mental ability to understand and sign Informed Consent Form
4. The score in Montgomery-Asberg Depression Rating Scale greater than or equal to 25 and the score in Clinical Global Impression greater than or equal to 4
5. Outpatients and inpatients
6. Age between 18 and 65 years

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

60 participants

**Key exclusion criteria**

1. Contraindications of used antidepressant treatments according to Summary of product (SPC)
2. The use of antidepressants as a monotherapy or as a part of combination that were ineffective in the treatment of current episode
3. The use of antipsychotics, thymostabilizers and other biological treatment of depression (ECT, rTMS, sleep deprivation etc.) during the study (anxiolytics and hypnotics for case of severe anxiety and insomnia are permitted) as well as formal psychotherapy
4. Comorbidity of Axis I and II (DSM IV) disorders in the 6 month before enrollment to the study
5. Severe or uncontrolled somatic disorders, likely to cause depressive symptoms or interfere with the conduct of the study

**Date of first enrolment**

01/01/2009

**Date of final enrolment**

30/06/2011

**Locations****Countries of recruitment**

Czech Republic

**Study participating centre**

**Prague Psychiatric Center**

Prague 8-Bohnice

Czech Republic

181 03

**Sponsor information****Organisation**

Ministry of Health (Czech Republic) - Internal Grant Agency

**Sponsor details**

Palackeho nam.4

Prague 2

Czech Republic

128 01

**Sponsor type**

Government

**Website**

<http://www.mzcr.cz>

**ROR**

<https://ror.org/00y6khe77>

**Funder(s)****Funder type**

Government

**Funder Name**

Ministry of Health (Czech Republic) - Internal Grant Agency (ref: No. NS 10368-3)

**Results and Publications****Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration